

# MMP FOR MODULI OF SHEAVES ON K3S VIA WALL-CROSSING: NEF AND MOVABLE CONES, LAGRANGIAN FIBRATIONS

AREND BAYER AND EMANUELE MACRÌ

**ABSTRACT.** We use wall-crossing with respect to Bridgeland stability conditions to systematically study the birational geometry of a moduli space  $M$  of stable sheaves on a K3 surface  $X$ :

- (a) We describe the nef cone, the movable cone, and the effective cone of  $M$  in terms of the Mukai lattice of  $X$ .
- (b) We establish a long-standing conjecture that predicts the existence of a birational Lagrangian fibration on  $M$  whenever  $M$  admits an integral divisor class  $D$  of square zero (with respect to the Beauville-Bogomolov form).

These results are proved using a natural map from the space of Bridgeland stability conditions  $\text{Stab}(X)$  to the cone  $\text{Mov}(X)$  of movable divisors on  $M$ ; this map relates wall-crossing in  $\text{Stab}(X)$  to birational transformations of  $M$ . In particular, every minimal model of  $M$  appears as a moduli space of Bridgeland-stable objects on  $X$ .

## CONTENTS

1. Introduction	2
2. Review: derived categories of K3 surfaces, Bridgeland stability conditions, moduli spaces, projectivity	7
3. Review: basic facts about hyperkähler varieties	13
4. Harder-Narasimhan filtrations in families	15
5. The hyperbolic lattice associated to a wall	18
6. Totally semistable non-isotropic walls	21
7. Divisorial contractions in the non-isotropic case	28
8. Isotropic walls are Uhlenbeck walls	31
9. Flopping walls	37
10. Main theorems	39
11. Application 1: Lagrangian fibrations	45
12. Application 2: Mori cone, nef cone, movable cone, effective cone	47
13. Examples of nef cones and movable cones	50
14. The geometry of flopping contractions	54
15. Le Potier's Strange Duality for isotropic classes	56
References	58

---

*Date:* January 30, 2013.

*2010 Mathematics Subject Classification.* 14D20 (Primary); 18E30, 14J28, 14E30 (Secondary).

*Key words and phrases.* Bridgeland stability conditions, Derived category, Moduli spaces of sheaves and complexes, Hyperkähler manifolds, Minimal Model Program, Cone Conjecture, Lagrangian fibrations.

## 1. INTRODUCTION

By our previous work [BM12], a moduli space of Bridgeland stable objects comes equipped with a numerically positive determinant line bundle, depending only on the stability condition. This provides a direct link between wall-crossing for stability conditions and birational transformations of the moduli space. In the present paper, we exploit this link to systematically study the most important birational properties of a moduli space  $M$  of stable sheaves (or Bridgeland stable complexes) on a K3 surface via wall-crossing.

The motivation for our work comes from several directions:

- Given the recent success [BCHM10] of the minimal model program (MMP), there has been a lot of interest to relate the MMP for moduli spaces to the underlying moduli problem; we refer [FS10] for a survey of the case of the moduli space  $\overline{M}_{g,n}$  of stable curves, which is known as the Hassett-Keel program. Ideally, one would like a moduli interpretation for every chamber of the base locus decomposition of the movable or effective cone; in our situation, we will show that this moduli interpretation comes naturally as a moduli of Bridgeland stable objects.
- In light of Verbitsky's recent proof in [Ver09] of a global Torelli statement for hyperkähler manifolds it has become particularly interesting to understand their nef cones: two hyperkähler varieties  $X_1, X_2$  are isomorphic if and only if there exists an isomorphism of integral Hodge structures  $H^2(X_1) \rightarrow H^2(X_2)$ , that is induced by parallel transport in a family, and that maps the nef cone of  $X_1$  to the nef cone of  $X_2$  (see [Huy11, Mar11]).
- According to a long-standing conjecture, the existence of a (birational) Lagrangian torus fibration  $M \rightarrow Y$  can be detected by an integral divisor class  $D \in \text{NS}(M)$  that has square zero with respect to the Beauville-Bogomolov pairing. Birationality of wall-crossing allows us to prove this conjecture by reducing it to the well-known case of a moduli space of torsion sheaves, first studied by Beauville in [Bea91].

**Birationality of wall-crossing and the map to the movable cone.** Consider a projective K3 surface  $X$ , and a primitive algebraic class  $\mathbf{v}$  in the Mukai lattice, whose self-intersection with respect to the Mukai pairing satisfies  $\mathbf{v}^2 > 0$ . Let  $\sigma, \tau$  be two stability conditions<sup>1</sup> on  $X$ , and assume that they are *generic* with respect to  $\mathbf{v}$ . By [BM12, Theorem 1.3], the moduli spaces  $M_\sigma(\mathbf{v})$  and  $M_\tau(\mathbf{v})$  of stable objects  $\mathcal{E} \in \text{D}^b(X)$  with Mukai vector  $\mathbf{v}(\mathcal{E}) = \mathbf{v}$  exist as a smooth projective variety. Choosing a path from  $\sigma$  to  $\tau$  in  $\text{Stab}(X)$  relates them by a series of wall-crossings. Our first result, based on a detailed analysis of the possible wall-crossings, is the following:

**Theorem 1.1.** *Let  $\sigma, \tau$  be generic stability conditions with respect to  $\mathbf{v}$ .*

- The two moduli spaces  $M_\sigma(\mathbf{v})$  and  $M_\tau(\mathbf{v})$  of Bridgeland-stable objects are birational to each other.*
- More precisely, there is a birational map induced by a derived (anti-)autoequivalence  $\Phi$  of  $\text{D}^b(X)$  in the following sense: there exists a common open subset  $U \subset M_\sigma(\mathbf{v})$ ,  $U \subset M_\tau(\mathbf{v})$ , with complements of codimension at least two, such that for any  $u \in U$ , the corresponding objects  $\mathcal{E}_u \in M_\sigma(\mathbf{v})$  and  $\mathcal{F}_u \in M_\tau(\mathbf{v})$  are related via  $\mathcal{F}_u = \Phi(\mathcal{E}_u)$ .*

<sup>1</sup>Throughout this introduction we will tacitly assume that all stability conditions on  $X$  are contained in the component of the space of stability conditions constructed by Bridgeland in [Bri08].

As a consequence, we can canonically identify the Néron-Severi groups of  $M_\sigma(\mathbf{v})$  and  $M_\tau(\mathbf{v})$ .

Now consider the chamber decomposition of the space  $\text{Stab}(X)$  of stability conditions with respect to  $\mathbf{v}$ : for each open chamber  $\mathcal{C} \subset \text{Stab}(X)$ , there is a moduli space  $M_{\mathcal{C}}(\mathbf{v})$  of  $\sigma$ -stable object of class  $\mathbf{v}$  for every  $\sigma \in \mathcal{C}$ . The main result of [BM12] gives a natural map

$$(1) \quad \ell_{\mathcal{C}}: \mathcal{C} \rightarrow \text{NS}(M_{\mathcal{C}}(\mathbf{v}))$$

to the Néron-Severi group of the moduli space, whose image is contained in the ample cone of  $M_{\mathcal{C}}(\mathbf{v})$ .

**Theorem 1.2.** *Fix a base point  $\sigma \in \text{Stab}(X)$ .*

(a) *Under the identification of the Néron-Severi groups induced by the birational maps of Theorem 1.1, the maps  $\ell_{\mathcal{C}}$  of (1) glue to a piece-wise analytic continuous map*

$$(2) \quad \ell: \text{Stab}(X) \rightarrow \text{NS}(M_{\sigma}(\mathbf{v})).$$

(b) *The image of  $\ell$  is the intersection of the movable cone with the big cone of  $M_{\sigma}(\mathbf{v})$ .*

(c) *The map  $\ell$  is compatible, in the sense that for any generic  $\sigma' \in \text{Stab}(X)$ , the moduli space  $M_{\sigma'}(\mathbf{v})$  is the birational model corresponding to  $\ell(\sigma')$ . In particular, every smooth  $K$ -trivial birational model of  $M_{\sigma}(\mathbf{v})$  appears as a moduli space  $M_{\mathcal{C}}(\mathbf{v})$  of Bridgeland stable objects for some chamber  $\mathcal{C} \subset \text{Stab}(X)$ .*

(d) *For a chamber  $\mathcal{C} \subset \text{Stab}(X)$ , we have  $\ell(\mathcal{C}) = \text{Amp}(M_{\mathcal{C}}(\mathbf{v}))$ .*

In other words, we can run the MMP of the moduli space via wall-crossing for Bridgeland stability conditions. Divisorial contractions appear as coarse moduli spaces for stability conditions on a wall.

The image  $\ell(\tau)$  of a stability condition  $\tau$  is determined by its central charge; see Theorem 10.2 for a precise statement.

In Theorem 5.7, we describe the correspondence between walls in the space of stability conditions and birational modifications of the moduli space  $M_{\sigma}(\mathbf{v})$ . For each wall  $\mathcal{W} \subset \text{Stab}(X)$ , there is a rank two sublattice  $\mathcal{H}_{\mathcal{W}}$  of the Mukai lattice that gets mapped to a line via the central charge  $Z$ , for every stability  $\tau = (Z, \mathcal{P}) \in \mathcal{W}$ ; Theorem 5.7 determines the wall-crossing behavior of the wall  $\mathcal{W}$  completely in terms of  $\mathcal{H}_{\mathcal{W}}$ .

The proof of Theorem 5.7 takes up Sections 5 to 9, and can be considered the heart of this paper. Our main technique is a detailed analysis of the possible configurations in  $\mathcal{H}_{\mathcal{W}}$  of Mukai vectors appearing the Jordan-Hölder filtrations of strictly semistable objects, and Harder-Narasimhan filtration in families; the latter allows us to compare the contraction morphism induced by the nef line bundle  $\ell(\sigma_0)$  for a stability condition  $\sigma_0$  on a wall with rational maps induced by the Harder-Narasimhan filtrations in families.

We can state the following consequence of Verbitsky's Torelli Theorem for hyperkähler manifolds, Mukai-Orlov's Derived Torelli Theorem for K3 surfaces, and Theorem 1.1. It completes Mukai's program, started in [Muk81, Muk87b], to understand birational maps between moduli spaces of sheaves via Fourier-Mukai transforms:

**Corollary 1.3.** *Let  $X$  and  $X'$  be smooth projective K3 surfaces. Let  $\mathbf{v} \in H_{\text{alg}}^*(X, \mathbb{Z})$  and  $\mathbf{v}' \in H_{\text{alg}}^*(X', \mathbb{Z})$  be primitive Mukai vectors. Let  $H$  (resp.,  $H'$ ) be a generic polarization with respect to  $\mathbf{v}$  (resp.,  $\mathbf{v}'$ ). Then  $M_H(\mathbf{v})$  is birational to  $M_{H'}(\mathbf{v}')$  if and only if there exists a derived (anti-)equivalence  $\Phi: \text{D}^b(X) \xrightarrow{\sim} \text{D}^b(X')$  with  $\Phi(\mathbf{v}) = \mathbf{v}'$ . In this case, we can choose  $\Phi$  such that it maps a generic object  $E \in M_H(\mathbf{v})$  to an object  $\Phi(E) \in M_{H'}(\mathbf{v}')$ .*

By [Tod08], stability is an open property in families; therefore such an equivalence  $\Phi$  induces a birational map  $M_{H'}(\mathbf{v}') \dashrightarrow M_H(\mathbf{v})$ . We will prove the Corollary at the end of Section 10.

**Cones of curves and divisors.** As an application, we can use Theorems 1.2 and 5.7 to determine the cones of effective, movable, and nef divisors (and thus dually the Mori cone of curves) of any moduli space of Gieseker-stable sheaves completely in terms of the algebraic Mukai lattice of  $X$ ; as an example we will state here our description of the nef cone:

Assume that  $H$  is a polarization that is generic with respect to  $\mathbf{v}$ , and let  $M_H(\mathbf{v})$  be the Gieseker-moduli space of stable sheaves with Mukai vector  $\mathbf{v}$ . By [Yos01], the construction of determinant line bundles gives an isomorphism  $\theta: \mathbf{v}^\perp \rightarrow \text{NS}(M_H(\mathbf{v}))$ , where  $\mathbf{v}^\perp$  denotes the orthogonal complement of  $\mathbf{v}$  inside the algebraic Mukai lattice  $H_{\text{alg}}^*(X, \mathbb{Z})$ , and  $\text{NS}$  the Néron-Severi group of the moduli space; it identifies the Mukai pairing  $(\_, \_)$  in  $H_{\text{alg}}^*(X, \mathbb{Z})$  with the Beauville-Bogomolov pairing on  $\text{NS}(M_H(\mathbf{v}))$ . Let  $\text{Pos}(M_H(\mathbf{v}))$  denote the cone of strictly positive classes  $D$  with respect to the Beauville-Bogomolov pairing, satisfying  $(D, D) > 0$  and  $(A, D) > 0$  for a fixed ample class  $A \in \text{NS}(M_H(\mathbf{v}))$ . We let  $\overline{\text{Pos}}(M_H(\mathbf{v}))$  denote its closure, and call it the *positive cone*.

**Theorem 12.1.** *Consider the chamber decomposition of the closed positive cone  $\overline{\text{Pos}}(M_H(\mathbf{v}))$  whose walls are given by linear subspaces of the form*

$$\theta(\mathbf{v}^\perp \cap \mathbf{a}^\perp),$$

*for all  $\mathbf{a} \in H_{\text{alg}}^*(X, \mathbb{Z})$  satisfying  $\mathbf{a}^2 \geq -2$  and  $0 \leq (\mathbf{v}, \mathbf{a}) \leq \frac{\mathbf{v}^2}{2}$ . Then the nef cone of  $M_H(\mathbf{v})$  is one of the chambers of this chamber decomposition.*

*In other words, given an ample class  $A \in \text{NS}(M_H(\mathbf{v}))$ , a class  $D \in \overline{\text{Pos}}(M_H(\mathbf{v}))$  is nef if and only if  $(D, \theta(\mathbf{a})) \geq 0$  for all classes  $\mathbf{a}$  as above that also satisfy  $(A, \theta(\mathbf{a})) > 0$ .*

We refer to Section 12 for similar statements for the movable and effective cone. Note that the intersection of the movable cone with the positive cone has been determined in general by Markman in [Mar11, Lemma 6.22] for any hyperkähler variety; the pseudo-effective can also easily be deduced from Markman's results. In these cases, our method gives an alternative wall-crossing proof; the new result is the description of the boundary, due to the proof of the Lagrangian conjecture discussed below.

However, there was no known description of the nef cone except for specific examples, even in the case of the Hilbert scheme of points. A general conjecture by Hassett and Tschinkel, [HT10, Thesis 1.1], suggested that the nef cone (or better, its dual, the Mori cone) of a hyperkähler variety  $M$  depends only on the lattice of algebraic cycles in  $H_2(M, \mathbb{Z})$ . In small dimension, their conjecture has been verified in [HT01, HT09, HT10, HHT12, BJ11]. The original conjecture turned out to be incorrect, already for Hilbert schemes (see [BM12, Remark 9.4] and [CK12, Remark 8.10]). However, Theorem 12.1 is in fact very closely related to the Hassett-Tschinkel Conjecture: we will explain this precisely in Section 12, in particular Proposition 12.6 and Remark 12.7. In Section 13, we give many explicit examples of nef cone and movable cones.

**Existence of Lagrangian fibrations.** The geometry of a hyperkähler variety  $M$  is particularly rigid. For example, Matsushita proved in [Mat01] that any map  $f: M \rightarrow Y$  with connected fibers and  $\dim(Y) < \dim(M)$  is a Lagrangian fibration; further, Hwang proved in [Hwa08] that if  $Y$  is smooth, it must be isomorphic to projective space.

It then becomes a natural question to ask when such a fibration exists, or when it exists birationally. According to a long-standing conjecture, this can be detected purely in terms of the quadratic Beauville-Bogomolov form on the Néron-Severi group of  $M$ :

**Conjecture 1.4** (Tyurin-Bogomolov-Hassett-Tschinkel-Huybrechts-Sawon). *Let  $M$  be a compact hyperkähler manifold of dimension  $2m$ , and let  $q$  denote its Beauville-Bogomolov form.*

- (a) *There exists an integral divisor class  $D$  with  $q(D) = 0$  if and only if there exists a birational hyperkähler manifold  $M'$  admitting a Lagrangian fibration.*
- (b) *If in addition,  $M$  admits a nef integral divisor class  $D$  with  $q(D) = 0$ , then there exists a Lagrangian fibration  $f: M \rightarrow \mathbb{P}^m$  induced by the complete linear system of a multiple of  $D$ .*

In the literature, it was first suggested by Hassett-Tschinkel in [HT01] for symplectic four-folds, and, independently, by Huybrechts [GHJ03] and Sawon [Saw03] in general; see [Ver10] for more remarks on the history of the Conjecture.

Based on the birationality of wall-crossing, we can prove this conjecture for moduli spaces of sheaves on a K3 surface:

**Theorem 1.5.** *Let  $X$  be a smooth projective K3 surface. Let  $\mathbf{v} \in H_{\text{alg}}^*(X, \mathbb{Z})$  be a primitive Mukai vector with  $\mathbf{v}^2 > 0$  and let  $H$  be a generic polarization with respect to  $\mathbf{v}$ . Then Conjecture 1.4 holds for the moduli space  $M_H(\mathbf{v})$  of  $H$ -Gieseker-stable sheaves.*

The basic idea of our proof is the following: as we recalled above, the Néron-Severi group of  $M_H(\mathbf{v})$ , along with its Beauville-Bogomolov form, is isomorphic to the orthogonal complement  $\mathbf{v}^\perp \subset H_{\text{alg}}^*(X, \mathbb{Z})$  of  $\mathbf{v}$  in the algebraic Mukai lattice of  $X$ , along with the restriction of the Mukai pairing. The existence of an integral divisor  $D = c_1(L)$  with  $q(D) = 0$  is thus equivalent to the existence of an isotropic class  $\mathbf{w} \in \mathbf{v}^\perp$ : a class with  $(\mathbf{w}, \mathbf{w}) = 0$  and  $(\mathbf{v}, \mathbf{w}) = 0$ . The moduli space  $Y = M_H(\mathbf{w})$  is a smooth K3 surface, and the associated Fourier-Mukai transform  $\Phi$  sends sheaves of class  $\mathbf{v}$  on  $X$  to complexes of rank 0 on  $Y$ . While these complexes on  $Y$  are typically not sheaves—not even for a generic object in  $M_H(\mathbf{v})$ —, we can arrange them to be Bridgeland-stable complexes with respect to a Bridgeland-stability condition  $\tau$  on  $D^b(Y)$ . We then deform  $\tau$  along a path with endpoint  $\tau'$ , such that  $\tau'$ -stable complexes of class  $\Phi_*(\mathbf{w})$  are Gieseker-stable sheaves, necessarily of rank zero. In other words, the Bridgeland-moduli space  $M_{\tau'}(\Phi_*(\mathbf{v}))$  is a moduli space of sheaves  $\mathcal{F}$  with support  $|\mathcal{F}|$  on a curve of fixed degree. As is well-known, the map  $\mathcal{F} \mapsto |\mathcal{F}|$  defines a map from  $M_{\tau'}(\Phi_*(\mathbf{v}))$  to the linear system of the associated curve, and this map is a Lagrangian fibration. On the other hand, birationality of wall-crossing shows that  $M_{\tau'}(\Phi_*(\mathbf{v})) = M_H(\mathbf{v})$  is birational to  $M_{\tau'}(\Phi_*(\mathbf{v}))$ .

The idea to use a Fourier-Mukai transform to prove Conjecture 1.4 was used previously by Markushevich [Mar06] and Sawon [Saw07] for specific family of examples of Hilbert schemes on K3 surfaces of Picard rank one; their assumptions implied that the Fourier-Mukai transform of a generic ideal sheaf is a stable torsion sheaf. Birationality of wall-crossing makes such a claim unnecessary.

**Remark 1.6.** By [MM12], Hilbert schemes of  $n$  points on projective K3 surfaces are Zariski-dense in the moduli space of hyperkähler varieties of  $K3^{[n]}$ -type.

Conjecture 1.4 has been proved independently by Markman [Mar13] for a general hyperkähler variety  $M$  of  $K3^{[n]}$ -type, under the assumption that the two dimensional subspace  $H^{2,0}(M) \oplus H^{0,2}(M)$  of  $H^2(M, \mathbb{C})$  does not contain any integral class. These varieties are

dense in the moduli space with respect to the analytic topology. His proof is completely different from ours, based on Verbitski’s Torelli Theorem, and a way to associate a K3 surface (purely lattice theoretically) to certain hyperkähler manifolds with a square-zero divisor class.

A related result has recently been announced by Matsushita [Mat12b].

**Remark 1.7.** Theorems 12.1 and 1.5 hold for every moduli space  $M_\sigma(\mathbf{v})$  of Bridgeland-stable object, for  $\mathbf{v}$  primitive and  $\sigma$  generic. Similarly, all main results hold for twisted K3 surfaces.

**Further applications.** We conclude the paper with two additional concrete applications of our methods. In Section 14, we explain how to deduce basic properties of the geometry of flopping contractions; in particular, we give examples where the contracted locus has either arbitrarily many connected components, or arbitrarily many irreducible components all intersecting in one point. In Section 15 we apply Theorem 1.5 to study Le Potier’s Strange Duality (in the case where one of the two classes involved has square zero).

**Relation to previous work on wall-crossing.** Various authors have previously studied examples of the relation between wall-crossing and the birational geometry of the moduli space induced by the chamber decomposition of its cone of movable divisors: the first examples (for moduli of torsion sheaves on  $K$ -trivial surfaces) were studied in [AB13], and moduli on abelian surfaces were considered (in varying generality) in [MM11, Mac12, MYY11a, MYY11b, YY12, Yos12].

Several of our results have analogues for abelian surfaces that have been obtained previously by Yoshioka, or by Minamide, Yanagida and Yoshioka: the birationality of wall-crossing has been established in [MY11a, Theorem 4.3.1]; the ample cone of the moduli spaces is described in [MY11b, Section 4.3]; statements related to Theorem 1.2 can be found in [Yos12]; the analogue of Corollary 1.3 is contained in [Yos09, Theorem 0.1]; and Conjecture 1.4 is proved in [Yos09, Proposition 3.4 and Corollary 3.5] with the same basic approach.

The crucial difference between abelian surfaces and K3 surfaces is the existence spherical objects on the latter. They are responsible for the existence of *totally semistable walls* (walls for which there are no strictly stable objects) that are harder to control; in particular, these can correspond to any possible type of birational transformation (isomorphism, divisorial contraction, flop). The spherical classes are the main reason our wall-crossing analysis in Sections 5–9 is fairly involved.

A somewhat different behavior was established in [ABCH13] in many cases for the Hilbert scheme of points on  $\mathbb{P}^2$  (extended to torsion-free sheaves in [Hui12, BMW13], and to Hirzebruch surfaces in [BC12]): the authors show that the chamber decomposition in the space of stability conditions corresponds to the base locus decomposition of the *effective* cone. In particular, while the map  $\ell_C$  of equation (1) exists similarly in their situation, it will behave differently across walls corresponding to a divisorial contraction: in our case, the map “bounces back” into the ample cone, while in their case, it will extend across the wall.

**Acknowledgments.** Conversations with Ralf Schiffler dissuaded us from pursuing a failed approach to the birationality of wall-crossing, and we had extremely useful discussions with Daniel Huybrechts. Tom Bridgeland pointed us towards Corollary 1.3, and Dragos Oprea towards the results in Section 15. We also received helpful comments from Daniel Greb, Antony Maciocia, Alina Marian, Eyal Markman, Daisuke Matsushita, Misha Verbitsky, Kōta Yoshioka, and we would like to thank all of them.



The authors were visiting the Max-Planck-Institut Bonn, respectively the Hausdorff Center for Mathematics in Bonn, while working on this paper, and would like to thank both institutes for their hospitality and stimulating atmosphere.

A. B. is partially supported by NSF grant DMS-1101377. E. M. is partially supported by NSF grant DMS-1001482/DMS-1160466, Hausdorff Center for Mathematics, Bonn, and by SFB/TR 45.

**Notation and Convention.** For an abelian group  $G$  and a field  $k(= \mathbb{Q}, \mathbb{R}, \mathbb{C})$ , we denote by  $G_k$  the  $k$ -vector space  $G \otimes k$ .

Throughout the paper,  $X$  will be a smooth projective K3 surface over the complex numbers. For a (locally-Noetherian) scheme (or algebraic space)  $S$ , we will use the notation  $D^b(S)$  for its bounded derived category of coherent sheaves, and  $D_{qc}(S)$  for the unbounded derived category of quasi-coherent sheaves.

We will abuse notation and denote all derived functors as if they were underived. We denote by  $p_S$  and  $p_X$  the two projections from  $S \times X$  to  $S$  and  $X$ , respectively. Given  $\mathcal{E} \in D_{qc}(S \times X)$ , we denote the Fourier-Mukai functor associated to  $\mathcal{E}$  by

$$\Phi_{\mathcal{E}}(\_) := (p_X)_*(\mathcal{E} \otimes p_S^*(\_)).$$

Given  $E, F \in D^b(X)$ , we denote the Euler characteristic by

$$\chi(E, F) = \sum_p (-1)^p \text{ext}^p(E, F).$$

We denote by  $\text{NS}(X)$  the Néron-Severi group of  $X$ . The space of full numerical stability conditions on  $D^b(X)$  will be denoted by  $\text{Stab}(X)$ .

The skyscraper sheaf at a point  $x \in X$  is denoted by  $k(x)$ . For a complex number  $z \in \mathbb{C}$ , we denote its real and imaginary part by  $\Re z$  and  $\Im z$ , respectively.

For a K3 surface  $X$ , we denote the Mukai vector of an object  $E \in D^b(X)$  by  $\mathbf{v}(E)$ . We will often write it as  $\mathbf{v}(E) = (r, c, s)$ , where  $r$  is the rank of  $E$ ,  $c \in \text{NS}(X)$  and  $s$  the degree of  $\mathbf{v}(E)$ .

An object with  $\text{Hom}(E, E) = \mathbb{C}$  will be called a *Schur object*. By *simple object* in an abelian category we will denote an object that has no non-trivial subobjects. An object with  $\text{Ext}^1(E, E) = 0$  will be called *rigid*. A rigid Schur object will be called *spherical*.

For a spherical object  $S$  we denote the spherical twist at  $S$  by  $\text{ST}_S(\_)$ , defined in [ST01] by the exact triangle, for all  $E \in D^b(X)$ ,

$$\text{Hom}^\bullet(S, E) \otimes S \rightarrow E \rightarrow \text{ST}_S(E).$$

We will write *stable* (in italics) whenever we are considering strictly stable objects in a context where there exist strictly semistable objects: for a non-generic stability condition, or objects with non-primitive Mukai vector.

## 2. REVIEW: DERIVED CATEGORIES OF K3 SURFACES, BRIDGELAND STABILITY CONDITIONS, MODULI SPACES, PROJECTIVITY

In this section, we give a review of stability conditions K3 surfaces, and their moduli spaces of stable complexes. The main references are [Bri07, Bri08, Tod08, Yos01, BM12].

**Bridgeland stability conditions.** Let  $\mathcal{D}$  be a triangulated category.

**Definition 2.1.** A slicing  $\mathcal{P}$  of the category  $\mathcal{D}$  is a collection of full extension-closed subcategories  $\mathcal{P}(\phi)$  for  $\phi \in \mathbb{R}$  with the following properties:

- (a)  $\mathcal{P}(\phi + 1) = \mathcal{P}(\phi)[1]$ .
- (b) If  $\phi_1 > \phi_2$ , then  $\text{Hom}(\mathcal{P}(\phi_1), \mathcal{P}(\phi_2)) = 0$ .
- (c) For any  $E \in \mathcal{D}$ , there exists a collection of real numbers  $\phi_1 > \phi_2 > \dots > \phi_n$  and a sequence of triangles

$$(3) \quad 0 = E_0 \longrightarrow E_1 \longrightarrow E_2 \longrightarrow \dots \longrightarrow E_{n-1} \longrightarrow E_n = E$$

with  $A_i \in \mathcal{P}(\phi_i)$ .

The collection of exact triangles in (3) is called the *Harder-Narasimhan filtration* of  $E$ . Each subcategory  $\mathcal{P}(\phi)$  is extension-closed and abelian. Its nonzero objects are called semistable of phase  $\phi$ , and its simple objects are called stable.

We will write  $\phi_{\min}(E) := \phi_n$  and  $\phi_{\max}(E) := \phi_1$ . By  $\mathcal{P}(\phi - 1, \phi]$  we denote the full subcategory of objects with  $\phi_{\min}(E) > \phi - 1$  and  $\phi_{\max}(E) \leq \phi$ . This is the heart of a bounded t-structure  $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$  given by

$$\mathcal{D}^{\leq 0} = \mathcal{P}( > \phi - 1) = \{E \in \mathcal{D} : \phi_{\min} > \phi - 1\} \quad \text{and} \quad \mathcal{D}^{\geq 0} = \mathcal{P}(\leq \phi) = \{E \in \mathcal{D} : \phi_{\max} \leq \phi\}.$$

The associated truncation functors will be denoted by

$$\tau_{>\phi} : \mathcal{D} \rightarrow \mathcal{P}( > \phi) \quad \text{and} \quad \tau_{\leq\phi} : \mathcal{D} \rightarrow \mathcal{P}(\leq \phi)$$

**Definition 2.2.** A Bridgeland stability condition on  $\mathcal{D}$  is a triple  $(\Lambda, Z, \mathcal{P})$ , where

- $\Lambda$  is a lattice of finite rank together with a surjective map  $\mathbf{v} : K(\mathcal{D}) \twoheadrightarrow \Lambda$ ,
- $Z : \Lambda \rightarrow \mathbb{C}$  is a group homomorphism,
- $\mathcal{P}$  is a slicing of  $Z$ ,

satisfying the following compatibilities:

- (a)  $\frac{1}{\pi} \arg Z(E) = \phi$ , for all non-zero  $E \in \mathcal{P}(\phi)$ ;
- (b) given a norm  $\|\_ \|$  on  $\Lambda_{\mathbb{R}}$ , there exists a constant  $C > 0$  such that

$$|Z(E)| \geq C \|\mathbf{v}(E)\|,$$

for all  $E \in \mathcal{P}$ .

A stability condition is called *algebraic* if  $\text{Im}(Z) \subset \mathbb{Q} \oplus \mathbb{Q}\sqrt{-1}$ .

We let  $\text{Stab}_{\Lambda}(\mathcal{D})$  the set of stability conditions with fixed lattice  $\Lambda$ . The main theorem in [Bri07, KS08] shows that it is a complex manifold of dimension given by the rank of  $\Lambda$ .

**Remark 2.3.** By [Bri07, Lemma 8.2], we have a left action on  $\text{Stab}_{\Lambda}(\mathcal{D})$  by the autoequivalence group  $\text{Aut}(\mathcal{D})$ , and a right action by  $\widetilde{\text{GL}}_2(\mathbb{R})$ , the universal cover of the matrices in  $\text{GL}_2(\mathbb{R})$  with positive determinant. The first action is defined, for  $\Phi \in \text{Aut}(\mathcal{D})$ , by  $\Phi(Z, \mathcal{P}) = (Z \circ \phi_*^{-1}, \Phi(\mathcal{P}))$ , where  $\phi_*$  is the automorphism induced by  $\Phi$  at the level of Grothendieck groups. The second one is the lift of the action of  $\text{GL}_2(\mathbb{R})$  on  $\text{Hom}(K(\mathcal{D}), \mathbb{C})$  (by identifying  $\mathbb{C} \cong \mathbb{R}^2$ ).



**The algebraic Mukai lattice.** We let  $X$  be a smooth projective K3 surface.

**Definition 2.4.** Let  $H_{\text{alg}}^*(X, \mathbb{Z}) := H^0(X, \mathbb{Z}) \oplus \text{NS}(X) \oplus H^4(X, \mathbb{Z})$ .

(a) We denote by  $\mathbf{v}: K(X) \rightarrow H_{\text{alg}}^*(X, \mathbb{Z})$  the *Mukai vector*, given by

$$\mathbf{v}(E) := \text{ch}(E)\sqrt{\text{td}(X)}.$$

(b) The *Mukai pairing*  $(\_, \_)$  is defined on  $H_{\text{alg}}^*(X, \mathbb{Z})$  by

$$((r, c, s), (r', c', s')) := cc' - rs' - sr' \in \mathbb{Z}.$$

It is an even pairing of signature  $(2, \rho(X))$ , satisfying  $(\mathbf{v}(E), \mathbf{v}(F)) = -\chi(E, F)$ , for  $E, F \in K(X)$ .

(c) The *algebraic Mukai lattice* is defined to be the pair  $(H_{\text{alg}}^*(X, \mathbb{Z}), (\_, \_))$ .

Recall that an embedding  $i: V \rightarrow L$  of a lattice  $V$  into a lattice  $L$  is *primitive* if  $L/i(V)$  is a free abelian group. In particular, we call a non-zero vector  $\mathbf{v} \in H_{\text{alg}}^*(X, \mathbb{Z})$  *primitive* if it is not divisible in  $H_{\text{alg}}^*(X, \mathbb{Z})$ . Throughout the paper  $\mathbf{v}$  will often denote a primitive class with  $\mathbf{v}^2 > 0$ .

Given a Mukai vector  $\mathbf{v} \in H_{\text{alg}}^*(X, \mathbb{Z})$ , we denote its orthogonal complement by  $\mathbf{v}^\perp$ .

**Remark 2.5.** Let  $\alpha \in \text{Br}(X)$  be a torsion class in the Brauer group of  $X$ . The above definitions can be extended to the *twisted K3 surface*  $(X, \alpha)$ , as explained in [HS05]. In particular, we will denote the algebraic Mukai lattice in the twisted case by  $(H_{\text{alg}}^*(X, \alpha, \mathbb{Z}), (\_, \_))$ . For the basic theory of twisted K3 surfaces, we refer the reader to [Că100, HS05, Yos06, Lie07].

**Stability conditions on K3 surfaces.** Let  $X$  be a K3 surface.

**Definition 2.6.** A (full, numerical) *stability condition* on  $X$  is a Bridgeland stability condition on  $\text{D}^b(X)$ , whose the lattice  $\Lambda$  is given by the Mukai lattice  $H_{\text{alg}}^*(X, \mathbb{Z})$ .

In [Bri08], Bridgeland describes a connected component of the space of numerical stability conditions on  $X$ . These results have been extended to twisted K3 surfaces in [HMS08]. We will briefly summarize his main results in the following.

Let  $\beta, \omega \in \text{NS}(X)_{\mathbb{R}}$  be two real divisor classes, with  $\omega$  being ample. In [Bri08, Lemma 6.1], Bridgeland constructs a heart  $\mathcal{A}_{\omega, \beta}$  by tilting at a torsion pair in  $\text{Coh } X$ : its objects are two-term complexes  $E^{-1} \xrightarrow{d} E$  such that the  $\text{Ker } d$  is a torsion-free sheaf, all of whose HN filtration factors with respect to  $\mu_\omega$ -slope stability have slope  $\mu_\omega \leq \omega \cdot \beta$ , and such that the torsion-free part of  $\text{Cok } d$  only has HN filtration factors of slope  $\mu_\omega > \omega \cdot \beta$ .

**Theorem 2.7** ([Bri08, Sections 10, 11]). *Let  $\sigma = (Z, \mathcal{P})$  be a stability condition such that all skyscraper sheaves  $k(x)$  of points are  $\sigma$ -stable. Then there are real divisor classes  $\omega, \beta \in \text{NS}(X)_{\mathbb{R}}$  with  $\omega$  ample, such that, up to the  $\text{GL}_2$ -action,  $\sigma$  is determined by  $\mathcal{P}((0, 1]) = \mathcal{A}_{\omega, \beta}$  and*

$$Z(E) = \left( e^{i\omega + \beta}, \mathbf{v}(E) \right).$$

We will denote this stability condition by  $\sigma_{\omega, \beta}$ , and the open subset of  $\text{Stab}(X)$  consisting of such stability conditions by  $U(X)$ .

Using the Mukai pairing, we can identify any central charge  $Z \in \text{Hom}(H_{\text{alg}}^*(X, \mathbb{Z}), \mathbb{C})$  with a vector in  $H_{\text{alg}}^*(X, \mathbb{Z}) \otimes \mathbb{C}$ . We will next describe the set of central charges using this identification: Let  $\mathcal{P}(X) \subset H_{\text{alg}}^*(X, \mathbb{Z}) \otimes \mathbb{C}$  be the set of vectors  $\Omega$  such that  $\Im \Omega, \Re \Omega$  span a

positive definite 2-plane in  $H_{\text{alg}}^*(X, \mathbb{Z}) \otimes \mathbb{R}$ . The subset  $\mathcal{P}_0(X)$  is defined as the orthogonal complement of all spherical classes:

$$\mathcal{P}_0(X) = \{ \Omega \in \mathcal{P}(X) : (\Omega, \mathbf{s}) \neq 0, \text{ for all } \mathbf{s} \in H_{\text{alg}}^*(X, \mathbb{Z}) \text{ with } \mathbf{s}^2 = -2 \}.$$

Finally,  $\mathcal{P}_0(X)$  has two connected components (corresponding to the orientation induced on the plane spanned by  $\Im \Omega, \Re \Omega$ ), and we let  $\mathcal{P}_0^+(X)$  be the component containing vectors of the form  $e^{i\omega+\beta}$  for  $\omega$  ample,  $\omega^2 \gg 0$ .

**Theorem 2.8** ([Bri08, Section 8]). *Let  $\text{Stab}^\dagger(X)$  be the connected component of the space of stability conditions containing  $U(X)$ . Let  $\mathcal{Z}: \text{Stab}^\dagger(X) \rightarrow H_{\text{alg}}^*(X, \mathbb{Z}) \otimes \mathbb{C}$  be the map sending a stability conditions  $(Z, \mathcal{P})$  to  $\Omega_Z$ , where  $\mathcal{Z}(\_) = (\Omega_Z, \_)$ .*

*Then  $\mathcal{Z}$  is a covering map of  $\mathcal{P}_0^+(X)$ .*

**Remark 2.9.** Theorem 2.7 and Theorem 2.8 have analogous statements, though slightly more technical, in the context of twisted K3 surfaces  $(X, \alpha)$ , as proved in [HMS08, Section 3.1]. The analogue objects in the twisted case will be denoted by  $U(X, \alpha)$ ,  $\text{Stab}^\dagger(X, \alpha)$ ,  $\mathcal{P}_0^+(X, \alpha)$ , etc.

We will need the following observation:

**Proposition 2.10.** *The stability conditions  $\sigma_{\omega, \beta}$  on  $U(X, \alpha)$  and  $\sigma_{\omega, -\beta}$  on  $U(X, -\alpha)$  are dual to each other in the following sense: An object  $E \in D^b(X, \alpha)$  is  $\sigma_{\omega, \beta}$ -(semi)stable of phase  $\phi$  if and only if its derived dual  $E^\vee[2] \in D^b(X, -\alpha)$  is  $\sigma_{\omega, -\beta}$ -(semi)stable of phase  $-\phi$ .*

*Proof.* This follows as in [BMT11, Proposition 4.3.6].  $\square$

**Walls.** One of the main properties of Bridgeland stability conditions is that the space of stability conditions carries a well-behaved wall and chamber structure. This is due to Bridgeland and Toda (the precise statement we need is in [BM12, Proposition 2.3]).

Let  $(X, \alpha)$  be a twisted K3 surface and let  $\mathbf{v} \in H_{\text{alg}}^*(X, \alpha, \mathbb{Z})$  be a Mukai vector. Then there exists a locally finite set of *walls* (real codimension one submanifolds with boundary) in  $\text{Stab}(X, \alpha)$ , depending only on  $\mathbf{v}$ , with the following properties:

- (a) When  $\sigma$  varies within a chamber, the sets of  $\sigma$ -semistable and  $\sigma$ -stable objects of class  $\mathbf{v}$  does not change.
- (b) When  $\sigma$  lies on a single wall  $\mathcal{W} \subset \text{Stab}(X, \alpha)$ , then there is a  $\sigma$ -semistable object that is unstable in one of the adjacent chambers, and semistable in the other adjacent chamber.
- (c) When we restrict to an intersection of finitely many walls  $\mathcal{W}_1, \dots, \mathcal{W}_k$ , we obtain a wall-and-chamber decomposition on  $\mathcal{W}_1 \cap \dots \cap \mathcal{W}_k$  with the same properties, where the walls are given by the intersections  $\mathcal{W} \cap \mathcal{W}_1 \cap \dots \cap \mathcal{W}_k$  for any of the walls  $\mathcal{W} \subset \text{Stab}(X, \alpha)$  with respect to  $\mathbf{v}$ .

Moreover, if  $\mathbf{v}$  is primitive, then  $\sigma$  lies on a wall if and only if there exists a strictly  $\sigma$ -semistable object of class  $\mathbf{v}$ . The Jordan-Hölder filtration of  $\sigma$ -semistable objects does not change when  $\sigma$  varies within a chamber.

**Definition 2.11.** Let  $\mathbf{v} \in H_{\text{alg}}^*(X, \alpha, \mathbb{Z})$ . A stability condition is called *generic* with respect to  $\mathbf{v}$  if it does not lie on a wall.

**Remark 2.12.** Given a polarization  $H$  that is generic with respect to  $\mathbf{v}$ , there is always a Gieseker chamber  $\mathcal{C}$ : for  $\sigma \in \mathcal{C}$ , the moduli space  $M_\sigma(\mathbf{v})$  of Bridgeland stable objects is exactly the moduli space of  $H$ -Gieseker stable sheaves; see [Bri08, Proposition 14.2].

**Moduli spaces and projectivity.** Let  $(X, \alpha)$  be a twisted K3 surface and let  $\mathbf{v} \in H_{\text{alg}}^*(X, \alpha, \mathbb{Z})$ . Following [Tod08], by fixing  $\sigma = (Z, \mathcal{P}) \in \text{Stab}^\dagger(X, \alpha)$  and  $\phi \in \mathbb{R}$  such that  $Z(\mathbf{v}) \in \mathbb{R}_{>0} \cdot e^{\pi\phi\sqrt{-1}}$ , we can define

$$\mathfrak{M}_\sigma(\mathbf{v}, \phi) \quad (\text{resp. } \mathfrak{M}_\sigma^{\text{st}}(\mathbf{v}, \phi))$$

as the moduli stack of  $\sigma$ -semistable (resp.  $\sigma$ -stable) objects with phase  $\phi$  and Mukai vector  $\mathbf{v}$ . This is an Artin stack of finite type over  $\mathbb{C}$ . We will often omit  $\phi$  or  $\mathbf{v}$  from the notation, when it is clear from the context.

By [BM12, Theorem 1.3(a)] (which generalizes [MY11b, Theorem 0.0.2]), if  $\sigma \in \text{Stab}^\dagger(X, \alpha)$  is generic with respect to  $\mathbf{v}$ , then there exists a coarse moduli space  $M_\sigma(\mathbf{v})$  of  $\sigma$ -semistable objects with Mukai vector  $\mathbf{v}$ . It is a normal projective irreducible variety with  $\mathbb{Q}$ -factorial singularities. If  $\mathbf{v}$  is primitive, then  $M_\sigma(\mathbf{v}) = M_\sigma^{\text{st}}(\mathbf{v})$  is a smooth projective hyperkähler manifold (see Section 3).

By results of Yoshioka and Toda, there is a very precise criterion for non-emptiness of a moduli space, and it always has expected dimension:

**Theorem 2.13.** *Let  $\mathbf{v} = m\mathbf{v}_0 \in H_{\text{alg}}^*(X, \alpha, \mathbb{Z})$  be a vector with  $\mathbf{v}_0$  primitive and  $m > 0$ , and let  $\sigma \in \text{Stab}^\dagger(X, \alpha)$  be a generic stability condition with respect to  $\mathbf{v}$ .*

- (a) *The coarse moduli space  $M_\sigma(\mathbf{v})$  is non-empty if and only if  $\mathbf{v}_0^2 \geq -2$ .*
- (b) *Either  $\dim M_\sigma(\mathbf{v}) = \mathbf{v}^2 + 2$  and  $M_\sigma^{\text{st}}(\mathbf{v}) \neq \emptyset$ , or  $m > 1$  and  $\mathbf{v}_0^2 \leq 0$ .*

In other words, when  $\mathbf{v}^2 \neq 0$  and the dimension of the moduli space is positive, then it is given by  $\dim M_\sigma(\mathbf{v}) = \mathbf{v}^2 + 2$ .

*Proof.* This is well-known: we provide a proof for completeness. First of all, claim (a) follows from results of Yoshioka and Toda (see [BM12, Theorem 5.7]). Since  $\sigma$  is generic with respect to  $\mathbf{v}$ , we know that  $M_\sigma(\mathbf{v})$  exists as a projective variety, parameterizing  $S$ -equivalence classes of semistable objects. Moreover, if  $E \in M_\sigma(\mathbf{v})$ , and we let  $F \hookrightarrow E$  be such that  $\phi_\sigma(F) = \phi_\sigma(E)$ , then  $\mathbf{v}(F) = m'\mathbf{v}_0$ , for some  $m' > 0$ . Hence, the locus of properly semistable objects in  $M_\sigma(\mathbf{v})$  coincides with the image of the natural map

$$\text{SSL}: \coprod_{m_1+m_2=m} (M_\sigma(m_1\mathbf{v}_0) \times M_\sigma(m_2\mathbf{v}_0)) \longrightarrow M_\sigma(\mathbf{v}).$$

If we assume  $\mathbf{v}_0^2 > 0$  (and so  $\geq 2$ ), then we can proceed by induction on  $m$ . For  $m = 1$ ,  $M_\sigma^{\text{st}}(\mathbf{v}_0) = M_\sigma(\mathbf{v}_0)$  and the conclusion follows from the Riemann-Roch Theorem and [Muk87a]. If  $m > 1$ , then by the inductive assumption, we deduce that the image of the map SSL has dimension equal to the maximum of  $(m_1^2 + m_2^2)\mathbf{v}_0^2 + 4$ , for  $m_1 + m_2 = m$ .

We claim that we can construct a semistable object with vector  $\mathbf{v}$  which is also Schur. Indeed, again by the inductive assumption, we can consider a  $\sigma$ -stable object  $E(m-1)$  with vector  $(m-1)\mathbf{v}_0$ . Let  $F \in M_\sigma(\mathbf{v}_0)$ . Then, again by the Riemann-Roch Theorem,  $\text{Ext}^1(F, E(m-1)) \neq 0$ . We can take any non-trivial extension

$$0 \rightarrow E(m-1) \rightarrow E(m) \rightarrow F \rightarrow 0.$$

Since both  $E(m-1)$  and  $F$  are Schur, and they have no morphism between each other,  $E(m)$  is Schur as well.

Again by the Riemann-Roch Theorem and [Muk84], we deduce that the dimension of  $M_\sigma(\mathbf{v})$  is equal to  $\text{ext}^1(E(m), E(m)) = m^2\mathbf{v}_0^2 + 2$ . This shows that  $M_\sigma^{\text{st}}(\mathbf{v}) \neq \emptyset$ , as we wanted. The last part of the statement follows from [BM12, Lemma 6.1 & Lemma 6.2].  $\square$

**Line bundles on moduli spaces.** In this final section we recall the main result of [BM12]: it shows that every moduli space of Bridgeland-stable objects comes equipped with a numerically positive line bundle, naturally associated to the stability condition. To avoid complicating the notation, we state the results only in the untwisted case: the extension to the twisted case works analogously.

Let  $X$  be a K3 surface. Let  $S$  be a proper algebraic space of finite type over  $\mathbb{C}$ , let  $\sigma = (Z, \mathcal{P}) \in \text{Stab}(X)$ , and let  $\mathcal{E} \in D^b(S \times X)$  be a family of semistable objects of class  $\mathbf{v}$  and phase  $\phi$ , namely, for all closed point  $s \in S$ ,  $\mathcal{E}_s \in \mathcal{P}(\phi)$  with  $\mathbf{v}(\mathcal{E}_s) = \mathbf{v}$ .

We define a numerical divisor class  $\ell_\sigma \in \text{NS}(S)_{\mathbb{R}}$  on  $S$  as follows: To every curve  $C \subset S$ , we associate

$$\ell_\sigma: C \mapsto \ell_\sigma.C := \Im \left( -\frac{Z(\Phi_{\mathcal{E}} \mathcal{O}_C)}{Z(v)} \right),$$

and prove that it extends linearly to all curve classes.

**Remark 2.14.** The linearity can, for example, be seen by comparison with the classical determinant line bundle construction. First of all, we recall the Donaldson morphism ([HL10, Section 8.1]):

$$\lambda_{\mathcal{E}}: \mathbf{v}^{\#} \rightarrow \text{NS}(S),$$

given as the composition

$$\mathbf{v}^{\#} \xrightarrow{p_X^*} K_{\text{num}}(S \times X) \xrightarrow{[\mathcal{E}]} K_{\text{num}}(S \times X) \xrightarrow{(p_S)^*} K_{\text{num}}(S) \xrightarrow{\det} \text{NS}(S),$$

where  $K_{\text{num}}$  denotes the Grothendieck group of a variety modulo numerical equivalence, and

$$\mathbf{v}^{\#} := \{\mathbf{w} \in K_{\text{num}}(X) : \chi(\mathbf{v} \otimes \mathbf{w}) = 0\}.$$

When  $X$  is a K3 surface, it is more convenient to use a dual version, called the Mukai morphism,

$$\theta_{\mathcal{E}}: v^{\perp} \rightarrow \text{NS}(S), \quad \theta_{\mathcal{E}}(\mathbf{w}) := -\lambda_{\mathcal{E}}(\mathbf{v}^{-1}(\mathbf{w})^*).$$

If we assume  $Z(\mathbf{v}) = -1$ , and write  $Z(\_) = (\Omega_Z, \_)$  as above, then

$$(4) \quad \ell_\sigma = \theta_{\mathcal{E}}(\Im \Omega_Z).$$

**Theorem 2.15** ([BM12]). *The following are the main properties of  $\ell_\sigma$ :*

- (a)  $\ell_\sigma$  is a nef divisor class on  $S$ . Additionally, for a curve  $C \subset S$ , we have  $\ell_\sigma.C = 0$  if and only if, for two general closed points  $c, c' \in C$ , the corresponding objects  $\mathcal{E}_c, \mathcal{E}_{c'} \in D^b(X)$  are  $S$ -equivalent.
- (b) For any Mukai vector  $\mathbf{v} \in H_{\text{alg}}^*(X, \mathbb{Z})$ ,  $\ell_\sigma$  induces a divisor class on the coarse moduli space  $M_\sigma(\mathbf{v})$ , when the stability condition  $\sigma \in \text{Stab}^\dagger(X)$  is generic with respect to  $\mathbf{v}$ . Moreover, in such a case,  $\ell_\sigma$  is an ample divisor on  $M_\sigma(\mathbf{v})$ .

For any chamber  $\mathcal{C} \subset \text{Stab}^\dagger(X)$ , we thus get a map

$$(5) \quad \ell_{\mathcal{C}}: \mathcal{C} \rightarrow \text{Amp}(M_{\mathcal{C}}(\mathbf{v})),$$

where we used the notation  $M_{\mathcal{C}}(\mathbf{v})$  to denote the coarse moduli space  $M_\sigma(\mathbf{v})$ , independent of the choice  $\sigma \in \mathcal{C}$ . The main goal of this paper is to understand the global behavior of this map.

We recall one more result from [BM12]. Let  $\mathbf{v} \in H_{\text{alg}}^*(X, \mathbb{Z})$  be a *primitive* vector with  $\mathbf{v}^2 \geq -2$ . Let  $\mathcal{W}$  be a wall for  $\mathbf{v}$  and let  $\sigma_0 \in \mathcal{W}$  be a generic stability condition on the wall, namely it does not belong to any other wall (see also Remark 5.6). We denote by  $\sigma_+$  and  $\sigma_-$  two generic stability conditions nearby  $\mathcal{W}$  in opposite chambers. Then all  $\sigma_{\pm}$ -semistable

objects are also  $\sigma_0$ -semistable. Hence,  $\ell_{\sigma_0}$  induces two nef divisors  $\ell_{\sigma_0,+}$  and  $\ell_{\sigma_0,-}$  on  $M_{\sigma_+}(\mathbf{v})$  and  $M_{\sigma_-}(\mathbf{v})$  respectively.

**Theorem 2.16** ([BM12, Theorem 1.4(a)]). *The divisors  $\ell_{\sigma_0,\pm}$  are big and nef on  $M_{\sigma_{\pm}}(\mathbf{v})$ . In particular, they are semi-ample, and induce birational contractions*

$$\pi^{\pm}: M_{\sigma_{\pm}}(\mathbf{v}) \rightarrow \overline{M}_{\pm},$$

where  $\overline{M}_{\pm}$  are normal irreducible projective varieties. The curves contracted by  $\pi^{\pm}$  are precisely curves of objects that are  $S$ -equivalent with respect to  $\sigma_0$ .

**Definition 2.17.** We call a wall  $\mathcal{W}$ :

- (a) a *fake wall*, if there are no curves in  $M_{\sigma_{\pm}}(\mathbf{v})$  of objects that are  $S$ -equivalent to each other with respect to  $\sigma_0$ ;
- (b) a *totally semistable wall*, if  $M_{\sigma_0}^{st}(\mathbf{v}) = \emptyset$ ;
- (c) a *flopping wall*, if we can identify  $\overline{M}_+ = \overline{M}_-$  and the induced morphism  $M_{\sigma_+}(\mathbf{v}) \dashrightarrow M_{\sigma_-}(\mathbf{v})$  induces a flopping contraction;
- (d) a *divisorial wall*, if the morphisms  $\pi^{\pm}: M_{\sigma_{\pm}}(\mathbf{v}) \rightarrow \overline{M}_{\pm}$  are both divisorial contractions.

By [BM12, Theorem 1.4(b)], if  $\mathcal{W}$  is not a fake wall and  $M_{\sigma_0}^{st}(\mathbf{v}) \subset M_{\sigma_{\pm}}(\mathbf{v})$  has complement of codimension at least two, then  $\mathcal{W}$  is a flopping wall. We will classify walls in Theorem 5.7.

### 3. REVIEW: BASIC FACTS ABOUT HYPERKÄHLER VARIETIES

In this section we give a short review on hyperkähler manifolds. The main references are [Bea83, GHJ03, Mar11].

**Definition 3.1.** A *projective hyperkähler manifold* is a simply connected smooth projective variety  $M$  such that  $H^0(M, \Omega_M^2)$  is one-dimensional, spanned by an everywhere non-degenerate holomorphic 2-form.

The Néron-Severi group of a hyperkähler manifold carries a natural bilinear form, called the *Fujiki-Beauville-Bogomolov form*. It is induced by a quadratic form on the whole second cohomology group  $q: H^2(M, \mathbb{Z}) \rightarrow \mathbb{Z}$ , which is primitive of signature  $(3, b_2(M) - 3)$ . It satisfies the Fujiki relation

$$(6) \quad \int_M \alpha^{2n} = F_M \cdot q(\alpha)^n, \quad \alpha \in H^2(M, \mathbb{Z}),$$

where  $2n = \dim M$  and  $F_M$  is the *Fujiki constant*, which depends only on the deformation type of  $M$ . We will mostly use the notation  $(\_, \_) := q(\_, \_)$  for the induced bilinear form on  $\text{NS}(M)$ .

The Hodge structure  $(H^2(M, \mathbb{Z}), q)$  behaves similarly to the case of a K3 surface. For example, by [Ver09], there is a weak global Hodge theoretic Torelli theorem for (deformation equivalent) hyperkähler manifolds.

Moreover, some positivity properties of divisors on  $M$  can be rephrased in terms of  $q$ . We first recall a few basic definitions on cones of divisors.

**Definition 3.2.** An integral divisor  $D \in \text{NS}(M)$  is called

- *big*, if its Iitaka dimension is maximal;
- *movable*, if its stable base-locus has codimension  $\geq 2$ ;

- *strictly positive*, if  $(D, D) > 0$  and  $(D, A) > 0$  for a fixed ample class  $A$  on  $M$ .

The real (not necessarily closed) cone generated by big (resp., movable, strictly positive, effective) integral divisors will be denoted by  $\text{Big}(M)$  (resp.,  $\text{Mov}(M)$ ,  $\text{Pos}(M)$ ,  $\text{Eff}(M)$ ). We have the following inclusions:

$$\begin{aligned} \text{Pos}(M) &\subset \text{Big}(M) \subset \text{Eff}(M) \\ \text{Nef}(M) &\subset \overline{\text{Mov}}(M) \subset \overline{\text{Pos}}(M) \subset \overline{\text{Big}}(M) = \overline{\text{Eff}}(M). \end{aligned}$$

The only non-trivial inclusion is  $\text{Pos}(M) \subset \text{Big}(M)$ , which follows from [Huy99, Corollary 3.10]. Divisors in  $\overline{\text{Pos}}(M)$  are called *positive*.

**Definition 3.3.** Let  $M$  be a projective hyperkähler manifold of dimension  $2n$ . A *Lagrangian fibration* is a surjective morphism with connected fibers  $h: M \rightarrow B$ , where  $B$  is a smooth projective variety, such that the generic fiber is Lagrangian with respect to the symplectic form  $\omega \in H^0(M, \Omega_M^2)$ .

By the Arnold-Liouville Theorem, any smooth fiber of a Lagrangian fibration is an abelian variety of dimension  $n$ . Moreover:

**Theorem 3.4** ([Mat99, Mat01] and [Hwa08]). *Let  $M$  be a projective hyperkähler manifold of dimension  $2n$ . Let  $B$  be a smooth projective variety of dimension  $0 < \dim B < 2n$  and let  $h: M \rightarrow B$  be a surjective morphism with connected fibers. Then  $h$  is a Lagrangian fibration, and  $B \cong \mathbb{P}^n$ .*

This result explains the importance of Conjecture 1.4. There has been recent series of articles on Lagrangian fibrations with a slightly different perspective. They treat the following question (see [Bea10, Question 1.6]): Given a torus  $L$  which is a Lagrangian subvariety of a HK manifold, does there exist a Lagrangian fibration with  $L$  as a fiber? This has been addressed in [GLR11a] for non-projective compact hyperkähler manifolds, and in general in [HW12, Mat12a] (based on previous results in [Ame11, GLR11b]).

The examples of hyperkähler manifolds we will consider are moduli spaces of stable complexes, as explained by the theorem below. It has been proven for moduli of sheaves in [Yos01, Sections 7 & 8], and generalized to Bridgeland stability conditions in [BM12, Theorem 5.9]):

**Theorem 3.5** (Huybrechts-O’Grady-Yoshioka). *Let  $(X, \alpha)$  be a twisted K3 surface and let  $\mathbf{v} \in H_{\text{alg}}^*(X, \alpha, \mathbb{Z})$  be a primitive vector with  $\mathbf{v}^2 \geq -2$ . Let  $\sigma \in \text{Stab}^\dagger(X, \alpha)$  be a generic stability condition with respect to  $\mathbf{v}$ . Then:*

- $M_\sigma(\mathbf{v})$  is a projective hyperkähler manifold, deformation-equivalent to the Hilbert scheme of points on any K3 surface.*
- The Mukai morphism induces an isomorphism*
  - $\bullet \theta_{\sigma, \mathbf{v}}: \mathbf{v}^\perp \xrightarrow{\sim} \text{NS}(M_\sigma(\mathbf{v})), \text{ if } \mathbf{v}^2 > 0;$
  - $\bullet \theta_{\sigma, \mathbf{v}}: \mathbf{v}^\perp / \mathbf{v} \xrightarrow{\sim} \text{NS}(M_\sigma(\mathbf{v})), \text{ if } \mathbf{v}^2 = 0.$

*Under this isomorphism, the quadratic Beauville-Bogomolov form for  $\text{NS}(M_\sigma(\mathbf{v}))$  coincides with the quadratic form of the Mukai pairing on  $(X, \alpha)$ .*

Here  $\theta_{\sigma, \mathbf{v}}$  is the Mukai morphism as in Remark 2.14, induced by a (quasi-)universal family. We will often drop  $\sigma$  or  $\mathbf{v}$  from the notation. It extends to an isomorphism of Hodge structures, if we take the orthogonal complement  $\mathbf{v}^\perp$  inside the whole cohomology  $H^*(X, \alpha, \mathbb{Z})$  (and not only in the algebraic part); it becomes identified with  $H^2(M, \mathbb{Z})$ . For  $\mathbf{v}^2 > 0$  the embedding  $H^2(M, \mathbb{Z}) \cong \mathbf{v}^\perp \hookrightarrow H^*(X, \alpha, \mathbb{Z})$  of integral Hodge structures determines the birational class of  $M_\sigma(\mathbf{v})$  via Verbitsky’s Torelli Theorem, see [Mar11, Example 9.6].



We will also need the following special case of a result by Namikawa and Wierzbka:

**Theorem 3.6** ([Wie03, Theorem 1.2 (ii)] and [Nam01, Proposition 1.4]). *Let  $M$  be a projective hyperkähler manifold of dimension  $2n$ , and let  $\overline{M}$  be a projective normal variety. Let  $\pi: M \rightarrow \overline{M}$  be a birational projective morphism. We denote by  $S_i$  the set of points  $p \in M$  such that  $\dim \pi^{-1}(p) = i$ . Then  $\dim S_i \leq 2n - 2i$ .*

*In particular, if  $\pi$  contracts a divisor  $D \subset M$ , we must have  $\dim \pi(D) = m - 2$ .*

Consider a non-primitive vector  $\mathbf{v}$ . As shown by O’Grady and Kaledin-Lehn-Sorger, the moduli space  $M_\sigma(\mathbf{v})$  can still be thought of as a singular hyperkähler manifold, in the following sense:

**Definition 3.7.** A normal projective variety  $M$  is said to have *symplectic singularities* if

- The smooth part  $M_{\text{reg}} \subset M$  admits a symplectic 2-form  $\omega$ ;
- For any resolution  $f: \tilde{M} \rightarrow M$ , the pull-back of  $\omega$  to  $f^{-1}(M_{\text{reg}})$  extends to a holomorphic form on  $\tilde{M}$ .

Our results in [BM12] reduce the following theorem to the case of moduli of sheaves:

**Theorem 3.8** ([O’G99] and [KLS06]). *Let  $(X, \alpha)$  be a twisted K3 surface and let  $\mathbf{v} = m\mathbf{v}_0 \in H_{\text{alg}}^*(X, \alpha, \mathbb{Z})$  be a Mukai vector with  $\mathbf{v}_0$  primitive and  $\mathbf{v}_0^2 \geq 2$ . Let  $\sigma \in \text{Stab}^\dagger(X, \alpha)$  be a generic stability condition with respect to  $\mathbf{v}$ . Then  $M_\sigma(\mathbf{v})$  has symplectic singularities.*

Given a hyperkähler manifold  $M$  and a dominant rational map  $M \dashrightarrow \overline{M}$ , where  $\overline{M}$  is a normal projective variety with symplectic singularities, then it follows from the definitions that  $\dim(M) = \dim(\overline{M})$ .

#### 4. HARDER-NARASIMHAN FILTRATIONS IN FAMILIES

In this section, we will show that results by Abramovich-Polishchuk and Toda imply the existence of Harder-Narasimhan filtrations in families, see Theorem 4.3.

Let  $Y$  be a smooth projective variety over  $\mathbb{C}$ , and let  $\sigma$  be a Bridgeland stability condition on  $D^b(Y)$ . The results we present will work as well in the twisted context: for simplicity of notation we only prove and state them in the untwisted context.

**Definition 4.1.** We say  $\sigma$  satisfies *openness of stability* if the following condition holds: for any scheme  $S$  of finite type over  $\mathbb{C}$ , and for any  $\mathcal{E} \in D^b(S \times Y)$  such that its derived restriction  $\mathcal{E}_s$  is a  $\sigma$ -semistable object of  $D^b(Y)$  for some  $s \in S$ , there exists an open neighborhood  $U \subset S$  of  $s$ , such that  $\mathcal{E}_{s'}$  is  $\sigma$ -semistable for all  $s' \in U$ .

**Theorem 4.2** ([Tod08, Section 3]). *Openness of stability holds when  $Y$  is a K3 surface and  $\sigma$  is a stability condition in the connected component  $\text{Stab}^\dagger(Y)$ .*<sup>2</sup>

**Theorem 4.3.** *Let  $\sigma = (Z, \mathcal{A}) \in \text{Stab}(Y)$  be an algebraic stability condition satisfying openness of stability. Assume we are given an irreducible variety  $S$  over  $\mathbb{C}$ , and an object  $\mathcal{E} \in D^b(S \times Y)$ . Then there exists a system of maps*

$$(7) \quad 0 = \mathcal{E}^0 \rightarrow \mathcal{E}^1 \rightarrow \mathcal{E}^2 \rightarrow \cdots \rightarrow \mathcal{E}^m = \mathcal{E}$$

<sup>2</sup>Note that in [Tod08, Section 3], this Theorem is only stated for families  $\mathcal{E}$  satisfying  $\text{Ext}^{<0}(\mathcal{E}_s, \mathcal{E}_s) = 0$  for all  $s \in S$ . However, Toda’s proof in Lemma 3.13 and Proposition 3.18 never uses that assumption.

in  $D^b(S \times Y)$ , and an open subset  $U \subset S$  with the following property: for any  $s \in U$ , the derived restriction of the system of maps (7)

$$0 = \mathcal{E}_s^0 \rightarrow \mathcal{E}_s^1 \rightarrow \mathcal{E}_s^2 \rightarrow \cdots \rightarrow \mathcal{E}_s^m = \mathcal{E}_s$$

is the Harder-Narasimhan filtration of  $\mathcal{E}_s$ .

The proof is based on the notion of constant family of t-structures due to Abramovich and Polishchuk, constructed in [AP06] (in case  $S$  is smooth) and [Pol07] (in general).

Throughout the remainder of this section, we will assume that  $\sigma$  and  $S$  satisfy the assumptions of Theorem 4.3. A t-structure is called *close to Noetherian* if it can be obtained via tilting from a t-structure whose heart is Noetherian. For  $\phi \in \mathbb{R}$ , the category  $\mathcal{P}((\phi - 1, \phi]) \subset D^b(Y)$  is the heart of a close to Noetherian bounded t-structure on  $Y$  given by  $\mathcal{D}^{\leq 0} = \mathcal{P}((\phi - 1, +\infty))$  and  $\mathcal{D}^{\geq 0} = \mathcal{P}((-\infty, \phi])$  (see the example discussed at the end of [Pol07, Section 1]). In this situation, Abramovich and Polishchuk's work induces a bounded t-structure  $(\mathcal{D}_S^{\leq 0}, \mathcal{D}_S^{\geq 0})$  on  $D^b(S \times Y)$ ; we paraphrase their main results as follows:

**Theorem 4.4** ([AP06, Pol07]). *Let  $\mathcal{A}$  be the heart of a close to Noetherian bounded t-structure  $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$  on  $D^b(Y)$ . Denote by  $\mathcal{A}_{qc} \subset D_{qc}(Y)$  the closure of  $\mathcal{A}$  under infinite coproducts in the derived category of quasi-coherent sheaves.*

- (a) *For any scheme  $S$  of finite type of  $\mathbb{C}$  there is a close to Noetherian bounded t-structure  $(\mathcal{D}_S^{\leq 0}, \mathcal{D}_S^{\geq 0})$  on  $D^b(S \times Y)$ , whose heart  $\mathcal{A}_S$  is characterized by*

$$\mathcal{E} \in \mathcal{A}_S \Leftrightarrow (p_Y)_*(\mathcal{E}|_{Y \times U}) \in \mathcal{A}_{qc} \quad \text{for every open affine } U \subset S$$

- (b) *The above construction defines a sheaf of t-structures over  $S$ : when  $S = \bigcup_i U_i$  is an open covering of  $S$ , then  $\mathcal{E} \in \mathcal{A}_S$  if and only if  $\mathcal{E}|_{Y \times U_i} \in \mathcal{A}_{U_i}$  for every  $i$ . In particular, for  $i: U \subset S$  open, the restriction functor  $i^*$  is t-exact.*
- (c) *When  $i: S' \subset S$  is a closed subscheme, then  $i_*$  is t-exact, and  $i^*$  is t-right exact.*

We briefly comment on the statements that are not explicitly mentioned in [Pol07, Theorem 3.3.6]: From part (i) of [Pol07, Theorem 3.3.6], it follows that the t-structure constructed there on  $D(S \times Y)$  descends to a bounded t-structure on  $D^b(S \times Y)$ . To prove that the push-forward in claim (c) is t-exact, we first use the sheaf property to reduce to the case where  $S$  is affine; in this case, the claim follows by construction. By adjointness, it follows that  $i^*$  is t-right exact.

For an algebraic stability condition  $\sigma = (Z, \mathcal{P})$  on  $D^b(Y)$  and a phase  $\phi \in \mathbb{R}$ , we will from now on denote its associated t-structure by  $\mathcal{P}( > \phi ) = \mathcal{D}^{\leq -1}$ ,  $\mathcal{P}( \leq \phi ) = \mathcal{D}^{\geq 0}$ , and the associated truncation functors by  $\tau^{>\phi}, \tau^{\leq\phi}$ . By [Pol07, Lemma 2.1.1], it induces a t-structure on  $D_{qc}(Y)$ , which we denote by  $\mathcal{P}_{qc}( > \phi ), \mathcal{P}_{qc}( \leq \phi )$ . For the t-structure on  $D^b(S \times Y)$  induced via Theorem 4.4, we will similarly write  $\mathcal{P}_S( > \phi ), \mathcal{P}_S( \leq \phi )$ , and  $\tau_S^{>\phi}, \tau_S^{\leq\phi}$ .

We start with a technical observation:

**Lemma 4.5.** *The t-structures on  $D^b(S \times Y)$  constructed via Theorem 4.4 satisfy the following compatibility relation:*

$$(8) \quad \bigcap_{\epsilon > 0} \mathcal{P}_S( \leq \phi + \epsilon ) = \mathcal{P}_S( \leq \phi ).$$

*Proof.* Assume  $\mathcal{E}$  is in the intersection of the left-hand side of (8). By the sheaf property, we may assume that  $S$  is affine. The assumption implies  $(p_Y)_*\mathcal{E} \in \mathcal{P}_{qc}( \leq \phi + \epsilon )$  for all  $\epsilon > 0$ .

By [Pol07, Lemma 2.1.1], we can describe  $\mathcal{P}_{qc}(\leq \phi + \epsilon) \subset D_{qc}(Y)$  as the right orthogonal complement of  $\mathcal{P}(> \phi + \epsilon) \subset D^b(Y)$  inside  $D_{qc}(Y)$ ; thus we obtain

$$\bigcap_{\epsilon > 0} \mathcal{P}_{qc}(\leq \phi + \epsilon) = \bigcap_{\epsilon > 0} (\mathcal{P}(> \phi + \epsilon))^\perp = \left( \bigcup_{\epsilon > 0} \mathcal{P}(> \phi + \epsilon) \right)^\perp = (\mathcal{P}(> \phi))^\perp = \mathcal{P}_{qc}(\leq \phi).$$

Hence  $(p_Y)_* \mathcal{E} \in \mathcal{P}_{qc}(\leq \phi)$ , proving the lemma.  $\square$

We next observe that the truncation functors  $\tau_S^{>\phi}, \tau_S^{\leq\phi}$  induce a slicing on  $D^b(S \times Y)$ . (See Definition 2.1 for the notion of slicing on a triangulated category.)

**Lemma 4.6.** *Assume that  $\sigma = (Z, \mathcal{P})$  is an algebraic stability condition, and  $\mathcal{P}_S(> \phi), \mathcal{P}_S(\leq \phi)$  are as defined above. There is a slicing  $\mathcal{P}_S$  on  $D^b(S \times Y)$  defined by*

$$\mathcal{P}_S(\phi) = \mathcal{P}_S(\leq \phi) \cap \bigcap_{\epsilon > 0} \mathcal{P}_S(> \phi - \epsilon).$$

Note that  $\mathcal{P}_S(\phi)$  cannot be characterized by the analogue of Theorem 4.4, part (a). For example, consider the case where  $Y$  is a curve and  $(Z, \mathcal{P})$  the standard stability condition corresponding to ordinary slope-stability in  $\text{Coh } Y$ . Then  $\mathcal{P}(1) \subset \text{Coh } Y$  is the category of torsion sheaves, and  $\mathcal{P}_S(1) \subset \text{Coh } S \times Y$  is the category of sheaves  $\mathcal{F}$  that are torsion relative over  $S$ . However, for  $U \subset S$  affine and a non-trivial family  $\mathcal{F}$ , the push-forward  $(p_Y)_* \mathcal{F}|_U$  is never a torsion sheaf.

*Proof.* By standard arguments, it is sufficient to consider the case  $\mathcal{E} \in \mathcal{A}_S := \mathcal{P}_S(0, 1]$ . In particular, since  $\sigma$  is algebraic, we can assume that both  $\mathcal{A} := \mathcal{P}(0, 1]$  and  $\mathcal{A}_S$  are Noetherian. For any  $\phi \in (0, 1]$ , we have  $\mathcal{P}_S(\phi, \phi + 1] \subset \langle \mathcal{A}_S, \mathcal{A}_S[1] \rangle$ . By [Pol07, Lemma 1.1.2], this induces a torsion pair  $(\mathcal{T}_\phi, \mathcal{F}_\phi)$  on  $\mathcal{A}_S$  with

$$\mathcal{T}_\phi = \mathcal{A}_S \cap \mathcal{P}_S(\phi, \phi + 1] \quad \text{and} \quad \mathcal{F}_\phi = \mathcal{A}_S \cap \mathcal{P}_S(\phi - 1, \phi].$$

Let  $T_\phi \hookrightarrow \mathcal{E} \twoheadrightarrow F_\phi$  be the induced short exact sequence in  $\mathcal{A}_S$ . Assume  $\phi < \phi'$ ; since  $\mathcal{F}_\phi \subset \mathcal{F}_{\phi'}$ , the surjection  $\mathcal{E} \twoheadrightarrow F_\phi$  factors via  $\mathcal{E} \twoheadrightarrow F_{\phi'} \twoheadrightarrow F_\phi$ . Since  $\mathcal{A}_S$  is Noetherian, the set of induced quotients  $\{F_\phi : \phi \in (0, 1]\}$  of  $\mathcal{E}$  must be finite. In addition, if  $F_\phi \cong F_{\phi'}$ , we must also have  $F_{\phi''} \cong F_\phi$  for any  $\phi'' \in (\phi, \phi')$ .

Thus, there exist real numbers  $\phi_0 = 1 > \phi_1 > \phi_2 > \dots > \phi_l > \phi_{l+1} = 0$  such that  $F_\phi$  is constant for  $\phi \in (\phi_{i+1}, \phi_i)$ , but such that  $F_{\phi_i - \epsilon} \neq F_{\phi_i + \epsilon}$ . Let us assume for simplicity that  $F_{\phi_1 + \epsilon} \cong \mathcal{E}$ ; the other case is treated similarly by setting  $F^1 = F_{\phi_1 + \epsilon}$ , and shifting all other indices by one. For  $i = 1, \dots, l$  we set

- $F^i := F_{\phi_i - \epsilon}$ ,
- $\mathcal{E}^i := \text{Ker}(\mathcal{E} \twoheadrightarrow F^i)$ , and
- $A^i = \mathcal{E}^i / \mathcal{E}^{i-1}$ .

We have  $\mathcal{E}^i \in \mathcal{P}_S(> \phi_i - \epsilon)$  and  $\mathcal{E}^{i-1} = \tau_S^{>\phi_i + \epsilon} \mathcal{E}^i$  for all  $\epsilon > 0$ . Hence the quotient  $A^i$  satisfies, for all  $\epsilon > 0$ ,

- $A^i \in \mathcal{P}_S(> \phi_i - \epsilon)$ ,
- $A^i \in \mathcal{P}_S(\leq \phi_i + \epsilon)$ .

The latter implies  $A^i \in \mathcal{P}_S(\leq \phi_i)$  by Lemma 4.5. By definition, we obtain  $A^i \in \mathcal{P}_S(\phi)$ . Finally, we have  $F^l \in \mathcal{P}_S(0, 1] \cap \mathcal{P}_S(\leq \epsilon)$  for all  $\epsilon > 0$ . Using Lemma 4.5 again, we obtain  $F^l = 0$ , and thus  $\mathcal{E}^l = \mathcal{E}$ . Thus the  $\mathcal{E}^i$  induce a Harder-Narasimhan filtration as claimed.  $\square$

The following lemma is an immediate extension of [AP06, Proposition 3.5.3]:

**Lemma 4.7.** *Assume that  $\mathcal{E} \in \mathcal{P}_S(\phi)$  for some  $\phi \in \mathbb{R}$ , and that  $\mathcal{E}_s \neq 0$  for  $s \in S$  generic. Then there exists a dense subset  $Z \subset S$ , such that  $\mathcal{E}_s$  is semistable of phase  $\phi$  for all  $s \in Z$ .*

*Proof.* By [AP06, Proposition 3.5.3], applied to the smooth locus of  $S$ , there exists a dense subset  $Z \subset S$  such that  $\mathcal{E}_s \in \mathcal{P}((\phi - 1, \phi])$ . Since  $\mathcal{E} \in \mathcal{P}(>\phi - \epsilon)$  for all  $\epsilon > 0$ , and since  $i_s^*$  is t-right exact, we also have  $\mathcal{E}_s \in \mathcal{P}(>\phi - \epsilon)$  for all  $\epsilon > 0$ . Considering the HN filtration of  $\mathcal{E}_s$ , this shows that  $\mathcal{E}_s \in \mathcal{P}(\phi)$  for all  $s \in Z$ .  $\square$

*Proof of Theorem 4.3.* The statement now follows easily from the above two lemmata: First of all, under the assumption of openness of stability, the dense subset  $Z$  of Lemma 4.7 may of course be taken to be open.

Given any  $\mathcal{E} \in D^b(S \times Y)$ , let

$$(9) \quad 0 = \mathcal{E}^0 \rightarrow \mathcal{E}^1 \rightarrow \dots \rightarrow \mathcal{E}^m = \mathcal{E}$$

be the Harder-Narasimhan filtration with respect to the slicing of Lemma 4.6, and let  $A^j$  be the HN filtration quotients fitting in the exact triangle  $\mathcal{E}^{j-1} \rightarrow \mathcal{E}^j \rightarrow A^j$ . Let  $j_1, \dots, j_l$  be the indices for which the generic fiber  $i_s^* A^j$  does not vanish, and let  $\phi_i$  be the phase of  $A^{j_i}$ . Then we claim that

$$(10) \quad 0 = \mathcal{E}^0 \rightarrow \mathcal{E}^{j_1} \rightarrow \mathcal{E}^{j_2} \rightarrow \dots \rightarrow \mathcal{E}^m = \mathcal{E}$$

has the desired property. Indeed, there is an open subset  $U$  such that for all  $s \in U$ , the fibers  $A_s^{j_i}$  are semistable for all  $i = 1, \dots, l$ , and such that  $A_s^j = 0$  for all  $j \notin \{j_1, \dots, j_l\}$ . Then, for each such  $s$ , the restriction of the sequence of maps (10) via  $i_s^*$  induces a sequence of maps that satisfies all properties of a HN filtration.  $\square$

## 5. THE HYPERBOLIC LATTICE ASSOCIATED TO A WALL

Our second main tool will be a rank two hyperbolic lattice associated to any wall. Let  $(X, \alpha)$  be a twisted K3 surface. Fix a primitive vector  $\mathbf{v} \in H_{\text{alg}}^*(X, \alpha, \mathbb{Z})$  with  $\mathbf{v}^2 > 0$ , and a wall  $\mathcal{W}$  of the chamber decomposition with respect to  $\mathbf{v}$ .

**Proposition 5.1.** *To each such wall, let  $\mathcal{H}_{\mathcal{W}} \subset H_{\text{alg}}^*(X, \alpha, \mathbb{Z})$  be the set of classes*

$$\mathbf{w} \in \mathcal{H}_{\mathcal{W}} \quad \Leftrightarrow \quad \Im \frac{Z(\mathbf{w})}{Z(\mathbf{v})} = 0 \quad \text{for all } \sigma = (Z, \mathcal{P}) \in \mathcal{W}.$$

*Then  $\mathcal{H}_{\mathcal{W}}$  has the following properties:*

- (a) *It is a primitive sublattice of rank two and of signature  $(1, -1)$  (with respect to the restriction of the Mukai form).*
- (b) *Let  $\sigma_+, \sigma_-$  be two sufficiently close and generic stability conditions on opposite sides of the wall  $\mathcal{W}$ , and consider any  $\sigma_+$ -stable object  $E \in M_{\sigma_+}(\mathbf{v})$ . Then any Harder-Narasimhan filtration factor  $A_i$  of  $E$  with respect to  $\sigma_-$  has Mukai vector  $\mathbf{v}(A_i)$  contained in  $\mathcal{H}_{\mathcal{W}}$ .*
- (c) *If  $\sigma_0$  is a generic stability condition on the wall  $\mathcal{W}$ , the conclusion of the previous claim also holds for any  $\sigma_0$ -semistable object  $E$  of class  $\mathbf{v}$ .*
- (d) *Similarly, let  $E$  be any object with  $\mathbf{v}(E) \in \mathcal{H}_{\mathcal{W}}$ , and assume that it is  $\sigma_0$ -stable for a generic stability condition  $\sigma_0 \in \mathcal{W}$ . Then every Jordan-Hölder factors of  $E$  with respect to  $\sigma_0$  will have Mukai vector contained in  $\mathcal{H}_{\mathcal{W}}$ .*

The precise meaning of “sufficiently close” will become apparent in the proof.

*Proof.* The first two claims of (a) are evident. To verify the claim on the signature, first note that by assumption on  $\mathbf{v}$ , the lattice  $\mathcal{H}_{\mathcal{W}}$  is either hyperbolic or positive (semi-)definite. On the other hand, consider a stability condition  $\sigma = (Z, \mathcal{A})$  with  $Z(\mathbf{v}) = -1$ . Since  $(\Im Z)^2 > 0$  by Theorem 2.8, since  $\mathcal{H}_{\mathcal{W}}$  is contained in the orthogonal complement of  $\Im Z$ , and since the algebraic Mukai lattice has signature  $(2, \rho(X))$ , this leaves the hyperbolic case as the only possibility.

In order to prove the remaining claims, consider an  $\epsilon$ -neighborhood  $B_\epsilon(\tau)$  of a generic stability conditions  $\tau \in \mathcal{W}$ , with  $0 < \epsilon \ll 1$ . Let  $\mathfrak{S}_{\mathbf{v}}$  be the set of objects  $E$  with  $\mathbf{v}(E) = \mathbf{v}$ , and that are semistable for some stability condition in  $B_\epsilon(\tau)$ . Let  $\mathfrak{U}_{\mathbf{v}}$  be the set of classes  $\mathbf{u} \in H_{\text{alg}}^*(X, \alpha, \mathbb{Z})$  that can appear as Mukai vectors of Jordan-Hölder factors of  $E \in \mathfrak{S}_{\mathbf{v}}$ , for any stability condition  $(Z', \mathcal{A}') \in B_\epsilon(\tau)$ . As shown in the proof of local finiteness of walls (see [Bri08, Proposition 9.3] or [BM11, Proposition 3.3]), the set  $\mathfrak{U}_{\mathbf{v}}$  is finite; indeed, such a class would have to satisfy  $|Z'(\mathbf{u})| < |Z'(\mathbf{v})|$ . Hence, the union of all walls for all classes in  $\mathfrak{U}_{\mathbf{v}}$  is still locally finite.

To prove claim (b), we may assume that  $\mathcal{W}$  is the only wall separating  $\sigma_+$  and  $\sigma_-$ , among all walls for classes in  $\mathfrak{U}_{\mathbf{v}}$ . Let  $\sigma_0 \in \mathcal{W}$  be a generic stability condition in the wall separating the chambers of  $\sigma_+, \sigma_-$ . It follows that both  $E$ , and each  $A_i$ , is  $\sigma_0$ -semistable. Since this argument works for generic  $\sigma_0$ , we must have  $v(A_i) \in \mathcal{H}_{\mathcal{W}}$  by the definition of  $\mathcal{H}_{\mathcal{W}}$ .

Claim (c) follows from the same discussion, and (d) similarly by considering the set of all walls for the classes  $\mathfrak{U}_{\mathbf{v}(E)}$  instead of  $\mathfrak{U}_{\mathbf{v}}$ .  $\square$

Our main approach is to characterize which hyperbolic lattices  $\mathcal{H} \subset H_{\text{alg}}^*(X, \alpha, \mathbb{Z})$  correspond to a wall, and to determine the type of wall purely in terms of  $\mathcal{H}$ . We start by making the following definition:

**Definition 5.2.** Let  $\mathcal{H} \subset H_{\text{alg}}^*(X, \alpha, \mathbb{Z})$  be a primitive rank two hyperbolic sublattice containing  $\mathbf{v}$ . A *potential wall*  $\mathcal{W}$  associated to  $\mathcal{H}$  is a connected component of the real codimension one submanifold of stability conditions  $\sigma = (Z, \mathcal{P})$  with the condition that  $Z(\mathcal{H})$  is contained in a line.

**Remark 5.3.** The statements of Proposition 5.1 are still valid when  $\mathcal{W}$  is a potential wall as in the previous definition.

**Definition 5.4.** Given any hyperbolic lattice  $\mathcal{H} \subset H_{\text{alg}}^*(X, \alpha, \mathbb{Z})$  of rank two containing  $\mathbf{v}$ , we will denote by  $P_{\mathcal{H}} \subset \mathcal{H} \otimes \mathbb{R}$  the cone generated by integral classes  $\mathbf{u} \in \mathcal{H}$  with  $\mathbf{u}^2 \geq 0$  and  $(\mathbf{v}, \mathbf{u}) > 0$ . We will call  $P_{\mathcal{H}}$  the *positive cone* of  $\mathcal{H}$ , and a class in  $P_{\mathcal{H}} \cap \mathcal{H}$  will be called a *positive class*.

We note that the condition  $(\mathbf{v}, \mathbf{u}) > 0$  just picks out one of the two components of the set of real classes with  $\mathbf{u}^2 > 0$ . Also observe that  $P_{\mathcal{H}}$  can be an open or closed cone, depending on whether the lattice contains integral classes  $\mathbf{w}$  that are isotropic:  $\mathbf{w}^2 = 0$ .

**Proposition 5.5.** Let  $\mathcal{W}$  be a potential wall associated to a hyperbolic rank two sublattice  $\mathcal{H} \subset H_{\text{alg}}^*(X, \alpha, \mathbb{Z})$ . For any  $\sigma = (Z, \mathcal{P}) \in \mathcal{W}$ , let  $C_\sigma \subset \mathcal{H} \otimes \mathbb{R}$  be the cone generated by classes  $\mathbf{u} \in \mathcal{H}$  satisfying the two conditions

$$\mathbf{u}^2 \geq -2 \quad \text{and} \quad \Re \frac{Z(\mathbf{u})}{Z(\mathbf{v})} > 0.$$

This cone does not depend on the choice of  $\sigma \in \mathcal{W}$ , and it contains  $P_{\mathcal{H}}$ .

If  $\mathbf{u} \in C_\sigma$ , then there exists a semistable object of class  $\mathbf{u}$  for every  $\sigma' \in \mathcal{W}$ . If  $\mathbf{u} \notin C_\sigma$ , then there does not exist a semistable object of class  $\mathbf{u}$  for generic  $\sigma' \in \mathcal{W}$ .

From here on, we will write  $C_{\mathcal{W}}$  instead of  $C_{\sigma}$ , and call it the cone of effective classes in  $\mathcal{H}$ . Given two different walls  $\mathcal{W}_1, \mathcal{W}_2$ , the corresponding effective cones  $C_{\mathcal{W}_1}, C_{\mathcal{W}_2}$  will only differ by spherical classes.

*Proof.* If  $\mathbf{u}^2 \geq -2$ , then by Theorem 2.13 there exists a  $\sigma$ -semistable object of class  $\mathbf{u}$  for every  $\sigma = (Z, \mathcal{P}) \in \mathcal{W}$ . Hence  $Z(\mathbf{u}) \neq 0$ , i.e., we cannot simultaneously have  $\mathbf{u} \in \mathcal{H}$  (which implies  $\Im \frac{Z(\mathbf{u})}{Z(\mathbf{v})} = 0$ ) and  $\Re \frac{Z(\mathbf{u})}{Z(\mathbf{v})} = 0$ . Therefore, the condition  $\Re \frac{Z(\mathbf{u})}{Z(\mathbf{v})} > 0$  is invariant under deforming a stability condition inside  $\mathcal{W}$ , and  $C_{\sigma}$  does not depend on the choice of  $\sigma \in \mathcal{W}$ .

Now assume for contradiction that  $P_{\mathcal{H}}$  is not contained in  $C_{\mathcal{W}}$ . Since  $\mathbf{v} \in C_{\mathcal{W}}$ , this is only possible if there is a real class  $\mathbf{u} \in P_{\mathcal{H}}$  with  $\Re \frac{Z(\mathbf{u})}{Z(\mathbf{v})} = 0$ ; after deforming  $\sigma \in \mathcal{W}$  slightly, we may assume  $\mathbf{u}$  to be integral. As above, this implies  $Z(\mathbf{u}) = 0$ , in contradiction to the existence of a  $\sigma$ -semistable object of class  $\mathbf{u}$ .

The statements about existence of semistable objects follow directly from Theorem 2.13.  $\square$

**Remark 5.6.** Note that by construction,  $C_{\mathcal{W}} \subset \mathcal{H} \otimes \mathbb{R}$  is strictly contained in a half-plane. In particular, there are only finitely many classes in  $C_{\mathcal{W}} \cap \mathbf{v} - C_{\mathcal{W}} \cap \mathcal{H}$  (in other words, effective classes  $\mathbf{u}$  such that  $\mathbf{v} - \mathbf{u}$  is also effective).

We will use this observation throughout in order to freely make genericity assumptions: a generic stability condition  $\sigma_0 \in \mathcal{W}$  will be a stability condition that does not lie on any additional wall (other than  $\mathcal{W}$ ) for any of the above-mentioned classes. Similarly, by stability conditions  $\sigma_+, \sigma_-$  nearby  $\sigma_0$  we will mean stability conditions that lie in the two chambers adjacent to  $\sigma_0$  for the wall-and-chamber decompositions with respect to any of the classes in  $C_{\mathcal{W}} \cap \mathbf{v} - C_{\mathcal{W}} \cap \mathcal{H}$ .

The behavior of the potential wall  $\mathcal{W}$  is completely determined by the lattice  $\mathcal{H}$  and its effective cone  $C_{\mathcal{W}}$ :

**Theorem 5.7.** *Let  $\mathcal{H} \subset H_{\text{alg}}^*(X, \alpha, \mathbb{Z})$  be a primitive hyperbolic rank two sublattice containing  $\mathbf{v}$ . Let  $\mathcal{W} \subset \text{Stab}(X, \alpha)$  be a potential wall associated to  $\mathcal{H}$  (see Definition 5.2).*

*The set  $\mathcal{W}$  is a totally semistable wall if and only if there exists either an isotropic class  $\mathbf{w} \in \mathcal{H}$  with  $(\mathbf{v}, \mathbf{w}) = 1$ , or an effective spherical class  $\mathbf{s} \in C_{\mathcal{W}} \cap \mathcal{H}$  with  $(\mathbf{s}, \mathbf{v}) < 0$ . In addition:*

- (a) *The set  $\mathcal{W}$  is a wall inducing a divisorial contraction if one of the following three conditions hold:*
  - (Brill-Noether):** *there exists a spherical class  $\mathbf{s} \in \mathcal{H}$  with  $(\mathbf{s}, \mathbf{v}) = 0$ , or*
  - (Hilbert-Chow):** *there exists an isotropic class  $\mathbf{w} \in \mathcal{H}$  with  $(\mathbf{w}, \mathbf{v}) = 1$ , or*
  - (Li-Gieseker-Uhlenbeck):** *there exists an isotropic class  $\mathbf{w} \in \mathcal{H}$  with  $(\mathbf{w}, \mathbf{v}) = 2$ .*
- (b) *Otherwise, if  $\mathbf{v}$  can be written as the sum  $\mathbf{v} = \mathbf{a} + \mathbf{b}$  of two positive<sup>3</sup> classes, or if there exists a spherical class  $\mathbf{s} \in \mathcal{H}$  with  $0 < (\mathbf{s}, \mathbf{v}) \leq \frac{\mathbf{v}^2}{2}$ , then  $\mathcal{W}$  is a wall corresponding to a flopping contraction.*
- (c) *In all other cases,  $\mathcal{W}$  is either a fake wall (if it is a totally semistable wall), or it is not a wall.*

The Gieseker-Uhlenbeck morphism from the moduli space of Gieseker semistable sheaves to slope-semistable vector bundle was constructed in [Li93]. Many papers deal with birational transformations between moduli spaces of twisted Gieseker semistable sheaves, induced by

<sup>3</sup>In the sense of Definition 5.4.



variations of the polarization. In particular, we refer to [Tha96, DH98] for the general theory of variation of GIT quotients and [EG95, FQ95, MW97] for the case of sheaves on surfaces. Theorem 5.7 can be thought as a generalization and completion of these results in the case of K3 surfaces.

The proof of the above theorem will be broken into four sections. We will distinguish two cases, depending on whether  $\mathcal{H}$  contains isotropic classes:

**Definition 5.8.** We say that  $\mathcal{W}$  is an *isotropic wall* if  $\mathcal{H}_{\mathcal{W}}$  contains an isotropic class.

In Section 6, we will analyze totally semistable non-isotropic walls, and Section 7 will describe non-isotropic walls corresponding to divisorial contractions. In Section 8, we will use a Fourier-Mukai transform to reduce the treatment of isotropic walls to the well-known behavior of the Li-Gieseker-Uhlenbeck morphism from the Gieseker moduli space to the Uhlenbeck space, and Section 9 will describe which of the remaining cases correspond to flopping walls, to fake walls, or to no wall at all.

## 6. TOTALLY SEMISTABLE NON-ISOTROPIC WALLS

In this section, we will analyze *totally semistable walls*; while some of our intermediate results hold in general, we will focus on the case where  $\mathcal{H}$  does not contain an isotropic class. The relevance of this follows from Theorem 2.13: in this case, if the dimension of a moduli space  $M_{\sigma}(\mathbf{u})$  is positive, then it is given by  $\mathbf{u}^2 + 2$ .

We will first describe the possible configurations of effective spherical classes in  $C_{\mathcal{W}}$ , and of spherical objects  $S \in \mathcal{P}(1)$  with  $\mathbf{v}(S) \in \mathcal{H}_{\mathcal{W}}$ .

We start with the following classical argument of Mukai (cfr. [Bri08, Lemma 5.2]). Given an exact sequence in the heart of a bounded t-structure  $\mathcal{A}$

$$0 \rightarrow A \rightarrow E \rightarrow B \rightarrow 0,$$

we assume that  $\mathrm{Hom}(A, B) = 0$ .

**Lemma 6.1** (Mukai). *We have an inequality*

$$\mathrm{ext}^1(E, E) \geq \mathrm{ext}^1(A, A) + \mathrm{ext}^1(B, B).$$

The following is a well-known consequence of Mukai's lemma (cfr. [HMS08, Section 2]):

**Lemma 6.2.** *Assume that  $S$  is a semistable rigid object with respect to a given stability condition. Then any Jordan-Hölder filtration factor of  $S$  is also spherical.*

*Proof.* Let  $S$  be any semistable object with  $\mathrm{Ext}^1(S, S) = 0$ . Pick any stable subobject  $T \subset S$  of the same phase. Then there exists a short exact sequence

$$\tilde{T} \hookrightarrow S \twoheadrightarrow R$$

with the following two properties:

- (a) The object  $\tilde{T}$  is an iterated extension of  $T$ .
- (b)  $\mathrm{Hom}(T, R) = 0$ .

Indeed, this can easily be constructed inductively: we let  $R_1 = S/T$ . If  $\mathrm{Hom}(T, S/T) = 0$ , the subobject  $\tilde{T} = T$  already has the desired properties. Otherwise, any non-zero morphism  $T \rightarrow R_1$  is necessarily injective; if we let  $R_2$  be its quotient, then the kernel of  $S \twoheadrightarrow R_2$  is a self-extension of  $T$ , and we can proceed inductively.

It follows that  $\text{Hom}(\tilde{T}, R) = 0$ , and we can apply Lemma 6.1 to conclude that  $\text{Ext}^1(\tilde{T}, \tilde{T}) = 0$ . Hence  $(\mathbf{v}(\tilde{T}), \mathbf{v}(\tilde{T})) < 0$ , which also implies  $(\mathbf{v}(T), \mathbf{v}(T)) < 0$ . Thus  $\mathbf{v}(T)$  is spherical, too.

The lemma follows by induction on the length of  $S$ .  $\square$

**Proposition 6.3.** *Let  $\mathcal{W}$  be a potential wall associated to the primitive hyperbolic lattice  $\mathcal{H}$ , and let  $\sigma_0 = (Z_0, \mathcal{P}_0) \in \mathcal{W}$  be a generic stability condition with  $Z_0(\mathcal{H}) \subset \mathbb{R}$ . Then  $\mathcal{H}$  and  $\sigma_0$  satisfy one of the following mutually exclusive conditions:*

- (a) *The lattice  $\mathcal{H}$  does not admit a spherical class.*
- (b) *The lattice  $\mathcal{H}$  admits, up to sign, a unique spherical class, and there exists a unique  $\sigma_0$ -stable object  $S \in \mathcal{P}_0(1)$  with  $\mathbf{v}(S) \in \mathcal{H}$ .*
- (c) *The lattice  $\mathcal{H}$  admits infinitely many spherical classes, and there exist exactly two  $\sigma_0$ -stable spherical objects  $S, T \in \mathcal{P}_0(1)$  with  $\mathbf{v}(S), \mathbf{v}(T) \in \mathcal{H}$ . In this case,  $\mathcal{H}$  is not isotropic.*

*Proof.* Given any spherical class,  $\mathbf{s} \in \mathcal{H}_{\mathcal{W}}$ , then by Theorem 2.13, there exists a  $\sigma$ -semistable object  $S$  with  $\mathbf{v}(S) = \mathbf{s}$  and  $S \in \mathcal{P}_0(1)$ . If  $\mathcal{H}$  admits a unique spherical class, then by Proposition 5.1 and Lemma 6.2,  $S$  must be stable.

Hence it remains to consider the case where  $\mathcal{H}$  admits two linearly independent spherical classes. If we consider the Jordan-Hölder filtrations of  $\sigma_0$ -semistable objects of the corresponding class, and apply Proposition 5.1 and Lemma 6.2, we see that there must be two  $\sigma_0$ -stable objects  $S, T$  whose Mukai vectors are linearly independent.

Now assume that there are three stable spherical objects  $S_1, S_2, S_3 \in \mathcal{P}(1)$ , and let  $\mathbf{s}_i = \mathbf{v}(S_i)$ . Since they are stable of the same phase, we have  $\text{Hom}(S_i, S_j) = 0$  for all  $i \neq j$ , as well as  $\text{Ext}^k(S_i, S_j) = 0$  for  $k < 0$ . Combined with Serre duality, this implies  $(\mathbf{s}_i, \mathbf{s}_j) = \text{ext}^1(S_i, S_j) \geq 0$ .


However, a rank two lattice of signature  $(1, -1)$  can never contain three spherical classes  $\mathbf{s}_1, \mathbf{s}_2, \mathbf{s}_3$  with  $(\mathbf{s}_i, \mathbf{s}_j) \geq 0$  for  $i \neq j$ : We may assume that  $\mathbf{s}_1, \mathbf{s}_2$  are linearly independent. Let  $m := (\mathbf{s}_1, \mathbf{s}_2) \geq 0$ ; since  $\mathcal{H}$  has signature  $(1, -1)$ , we have  $m \geq 3$ . If we write  $\mathbf{s}_3 = x\mathbf{s}_1 + y\mathbf{s}_2$ , we get the following implications:

$$\begin{aligned} (\mathbf{s}_1, \mathbf{s}_3) \geq 0 &\Rightarrow y \geq \frac{2}{m}x \\ (\mathbf{s}_2, \mathbf{s}_3) \geq 0 &\Rightarrow y \leq \frac{m}{2}x \\ (\mathbf{s}_3, \mathbf{s}_3) = -2 &\Rightarrow -2x^2 + 2mxy - 2y^2 < 0 \end{aligned}$$

However, by solving the quadratic equation for  $y$ , it is immediate that the term in the last inequality is positive in the range  $\frac{2}{m}x \leq y \leq \frac{m}{2}x$  (see also Figure 1).

Finally, if  $\mathcal{H}$  admits two linearly independent spherical class  $\mathbf{s}, \mathbf{t}$ , then the group generated by the associated reflections  $\rho_{\mathbf{s}}, \rho_{\mathbf{t}}$  is infinite; the orbit of  $\mathbf{s}$  consists of infinitely many spherical classes. Additionally, an isotropic class would be a rational solution of  $-2x^2 + 2mxy - 2y^2 = 0$ , but the discriminant  $m^2 - 4$  can never be a square when  $m$  is an integer  $m \geq 3$ .  $\square$

Whenever we are in case (c), we will denote the two  $\sigma_0$ -stable spherical objects by  $S, T$ . We may assume that  $S$  has smaller phase than  $T$  with respect to  $\sigma_+$ ; conversely,  $S$  has bigger phase than  $T$  with respect to  $\sigma_-$ . We will also write  $\mathbf{s} := \mathbf{v}(S), \mathbf{t} = \mathbf{v}(T)$ , and  $m := (\mathbf{s}, \mathbf{t}) > 2$ . We identify  $\mathbb{R}^2$  with  $\mathcal{H}_{\mathcal{W}} \otimes \mathbb{R}$  by sending the standard basis to  $(\mathbf{s}, \mathbf{t})$ ; under this identification, the ordering of phases in  $\mathbb{R}^2$  will be consistent with the ordering induced by  $\sigma_+$ . We denote by  $Q(x, y) = -2x^2 + 2mxy - 2y^2$  the pull-back of the quadratic form induced

FIGURE 1. The plane  $\mathcal{H}_W$ , oriented compatibly with  $\sigma_+$ 

by the Mukai pairing on  $\mathcal{H}_W$ . Let  $r_1 < r_2$  be the two solutions of  $-2r^2 + 2mr - 2 = 0$ ; they are both positive and irrational (as  $m^2 - 4$  cannot be a square for  $m \geq 3$  integral). The positive cone  $P_{\mathcal{H}}$  is thus the cone between the two lines  $y = r_i x$ , and the effective cone  $C_W$  is the upper right quadrant  $x, y \geq 0$ .

We will first prove that the condition for the existence of totally semistable walls given in Theorem 5.7 is necessary in the case of non-isotropic walls. We start with an easy numerical observation:

**Lemma 6.4.** *Given  $l > 1$  positive classes  $\mathbf{a}_1, \dots, \mathbf{a}_l \in P_{\mathcal{H}}$  with  $\mathbf{a}_i^2 > 0$ , set  $\mathbf{a} = \mathbf{a}_1 + \dots + \mathbf{a}_l$ . Then*

$$\sum_{i=1}^l (\mathbf{a}_i^2 + 2) < \mathbf{a}^2.$$

*Proof.* Since the  $\mathbf{a}_i$  are integral classes, and  $\mathcal{H}_W$  is an even lattice, we have  $\mathbf{a}_i^2 \geq 2$ . If  $\mathbf{a}_i \neq \mathbf{a}_j$ , then  $\mathbf{a}_i, \mathbf{a}_j$  span a lattice of signature  $(1, -1)$ , which gives  $(\mathbf{a}_i, \mathbf{a}_j) > \sqrt{\mathbf{a}_i^2 \mathbf{a}_j^2} \geq 2$ . Hence  $\mathbf{a}^2 > \sum_{i=1}^l \mathbf{a}_i^2 + 2l(l-1) \geq \sum_{i=1}^l \mathbf{a}_i^2 + 2l$ .  $\square$

**Lemma 6.5.** *Assume that the potential wall  $\mathcal{W}$  associated to  $\mathcal{H}$  satisfies the following conditions:*

- (a) *The wall is non-isotropic.*
- (b) *There does not exist an effective spherical class  $\mathbf{s} \in C_W$  with  $(\mathbf{s}, \mathbf{v}) < 0$ .*

*Then  $\mathcal{W}$  cannot be a totally semistable wall.*


In other words, there exists a  $\sigma_0$ -stable object of class  $\mathbf{v}$ .

*Proof.* We will consider two maps from the moduli space  $M_{\sigma_+}(\mathbf{v})$ : On the one hand, by Theorem 2.16, the line bundle  $\ell_{\sigma_0}$  on  $M_{\sigma_+}(\mathbf{v})$  induces a birational morphism

$$\pi^+ : M_{\sigma_+}(\mathbf{v}) \rightarrow \overline{M}.$$

The curves contracted by  $\pi^+$  are exactly curves of S-equivalent objects.

For the second map, first assume for simplicity that  $M_{\sigma_+}(\mathbf{v})$  is a fine moduli space, and let  $\mathcal{E}$  be a universal family. Consider the relative Harder-Narasimhan filtration for  $\mathcal{E}$  with respect to  $\sigma_-$  given by Theorem 4.3. Let  $\mathbf{a}_1, \dots, \mathbf{a}_m$  be the Mukai vectors of the semistable HN filtration quotients of a generic fiber  $\mathcal{E}_m$  for  $m \in M_{\sigma_+}(\mathbf{v})$ . On the open subset  $U$  of the Theorem 4.3,

FIGURE 2.  $\mathbf{a}^2 \leq \mathbf{v}^2$ 

the filtration quotients  $\mathcal{E}^i/\mathcal{E}^{i-1}$  are flat families of  $\sigma_-$ -semistable objects of class  $\mathbf{a}_i$ ; thus we get an induced rational map

$$\mathrm{HN}: M_{\sigma_+}(\mathbf{v}) \dashrightarrow M_{\sigma_-}(\mathbf{a}_1) \times \cdots \times M_{\sigma_-}(\mathbf{a}_m).$$

Let  $I \subset \{1, 2, \dots, m\}$  be the subset of indices  $i$  with  $\mathbf{a}_i^2 > 0$ , and let  $\mathbf{a} = \sum_{i \in I} \mathbf{a}_i$ .

Our first claim is  $\mathbf{a}^2 \leq \mathbf{v}^2$ , with equality if and only if  $\mathbf{a} = \mathbf{v}$ , i.e., if there are no classes with  $\mathbf{a}_i^2 < 0$ : Let us consider case (c) of Proposition 6.3; case (b) is simpler, and (a) trivial. Let  $v_x, v_y$  be the coordinates of  $\mathbf{v}$  in  $\mathbb{R}^2$ ; the conditions  $(\mathbf{s}, \mathbf{v}) \geq 0$  and  $(\mathbf{t}, \mathbf{v}) \geq 0$  imply  $\frac{2}{m} \leq \frac{v_y}{v_x} \leq \frac{m}{2}$ . The intersections of the lines  $y = \frac{2}{m}x$  and  $y = \frac{m}{2}x$  with the hyperbola  $Q(x, y) = \mathbf{v}^2$  are the points with horizontal and vertical tangent lines, respectively. (Indeed, the line  $y = \frac{2}{m}x$  as well as the hyperbola are fixed under the reflection  $\rho_s$ , and similarly  $y = \frac{m}{2}x$  for the reflection  $\rho_t$ .) Thus  $\mathbf{v}$  lies in between these two points, where the hyperbola has negative slope. If  $a_x, a_y$  are the coordinates of  $\mathbf{a}$ , then  $0 \leq a_x \leq v_x$  and  $0 \leq a_y \leq v_y$ , as  $\mathbf{v} - \mathbf{a}$  is an effective class in  $C_{\mathcal{W}}$ . The claim thus follows from Figure 2, which shows that  $\mathbf{a}$  cannot lie in the interior  $Q(x, y) > \mathbf{v}^2$  of the hyperbola.

Lemma 6.4 then implies

$$(11) \quad \mathbf{v}^2 + 2 \geq \mathbf{a}^2 + 2 \geq \sum_{i \in I} (\mathbf{a}_i^2 + 2),$$

with equality if and only if  $|I| = 1$ . By Theorem 2.13, part (b), this says that the target of the rational map HN has at most the dimension of the source:

$$(12) \quad \dim M_{\sigma_+}(\mathbf{v}) \geq \sum_{i=1}^m \dim M_{\sigma_-}(\mathbf{a}_i).$$

However, if  $\mathrm{HN}(E_1) = \mathrm{HN}(E_2)$ , then  $E_1, E_2$  are S-equivalent: indeed, they admit Jordan-Hölder filtrations that are refinements of their Harder-Narasimhan filtrations with respect to  $\sigma_-$ , which have the same filtration quotients.

It follows that any curve contracted by HN is also contracted by  $\pi^+$ ; therefore

$$\sum_{i=1}^m \dim M_{\sigma_-}(\mathbf{a}_i) \geq \dim \overline{M} = \dim M_{\sigma_+}(\mathbf{v})$$

Hence we have equality in each step of the above inequalities, the relative Harder-Narasimhan filtration is trivial, and the generic fiber  $\mathcal{E}_m$  is  $\sigma_-$ -stable.

In case  $M_{\sigma_+}(\mathbf{v})$  does not admit a universal family, we can construct HN by first passing to an étale neighborhood  $f: U \rightarrow M_{\sigma_+}(\mathbf{v})$  admitting a universal family; the induced rational map from  $U$  induced by the relative Harder-Narasimhan filtration will then factor via  $f$ .  $\square$

We recall some theory of Pell's equation in the language of spherical reflections of the hyperbolic lattice  $\mathcal{H}$ :

**Proposition and Definition 6.6.** *Let  $G_{\mathcal{H}} \subset \text{Aut } \mathcal{H}$  be the group generated by spherical reflections  $\rho_s$  for effective spherical classes  $s \in C_{\mathcal{W}}$ . Given a positive class  $\mathbf{v} \in P_{\mathcal{H}} \cap \mathcal{H}$ , the  $G_{\mathcal{H}}$ -orbit  $G_{\mathcal{H}} \cdot \mathbf{v}$  contains a unique class  $\mathbf{v}_0$  such that  $(\mathbf{v}_0, s) \geq 0$  for all effective spherical classes  $s \in C_{\mathcal{W}}$ .*

*We call  $\mathbf{v}_0$  the minimal class of the orbit  $G_{\mathcal{H}} \cdot \mathbf{v}$ .*

Note that the notion of minimal class depends on the potential wall  $\mathcal{W}$ , not just on the lattice  $\mathcal{H}$ .

*Proof.* Again, we only treat the case (c) of Proposition 6.3, the other cases being trivial. It is sufficient to prove that  $(\mathbf{v}_0, s) \geq 0$  and  $(\mathbf{v}_0, t) \geq 0$ . Assume  $(\mathbf{v}, s) < 0$ . Then  $\rho_s(\mathbf{v}) = \mathbf{v} - |(\mathbf{v}, s)| \cdot s$  is still in the upper right quadrant, with smaller  $x$ -coordinate than  $\mathbf{v}$ , and with the same  $y$ -coordinate. Similarly if  $(\mathbf{v}, t) < 0$ . If we proceed inductively, the procedure has to terminate, thus reaching  $\mathbf{v}_0$ .

The uniqueness follows from Proposition 6.7 below.  $\square$

Assume additionally that  $\mathcal{H}$  admits infinitely many spherical classes, so we are in case (c) of Proposition 6.3. The hyperbola  $\mathbf{v}^2 = -2$  intersects the upper right quadrant  $x, y \geq 0$  in two branches, starting at  $s$  and  $t$ , respectively. Let  $s_0 = s, s_{-1}, s_{-2}, \dots$  be the integral spherical classes on the lower branch starting at  $s$ , and  $t_1 = t, t_2, t_3, \dots$  be those on the upper branch starting at  $t$ , see also Figure 3. The  $s_i$  can be defined recursively by  $s_{-1} = \rho_s(t)$ , and  $s_{k-1} = \rho_{s_k}(s_{k+1})$  for  $k \leq -1$ ; similarly for the  $t_i$ .

**Proposition 6.7.** *Given a minimal class  $\mathbf{v}_0$  of a  $G_{\mathcal{H}}$ -orbit, define  $\mathbf{v}_i, i \in \mathbb{Z}$  via  $\mathbf{v}_i = \rho_{t_i}(\mathbf{v}_{i-1})$  for  $i > 0$ , and  $\mathbf{v}_i = \rho_{s_{i+1}}(\mathbf{v}_{i+1})$  for  $i < 0$ . Then the orbit  $G \cdot \mathbf{v}_0$  is given by  $\{\mathbf{v}_i : i \in \mathbb{Z}\}$ , where the latter are ordered according to their slopes in  $\mathbb{R}^2$ .*

Note that these classes may coincide pairwise, in case  $\mathbf{v}_0$  is orthogonal to  $s$  or  $t$ .

*Proof.* The group  $G_{\mathcal{H}}$  is the free product  $\mathbb{Z}_2 \star \mathbb{Z}_2$ , generated by  $\rho_s$  and  $\rho_t$ . It is straightforward to check that with  $\mathbf{v}_i$  defined as above, we have

$$\mathbf{v}_{-1} = \rho_s(\mathbf{v}_0), \quad \mathbf{v}_{-2} = \rho_s \rho_t(\mathbf{v}_0), \quad \mathbf{v}_{-3} = \rho_s \rho_t \rho_s(\mathbf{v}_0), \dots,$$


and similarly  $\mathbf{v}_1 = \rho_t(\mathbf{v}_0)$  and so on. This list contains  $g(\mathbf{v}_0)$  for all  $g \in \mathbb{Z}_2 \star \mathbb{Z}_2$ . That the  $\mathbf{v}_i$  are ordered by slopes is best seen by drawing a picture; see also Figure 3.  $\square$

For  $i > 0$ , let  $T_i^+ \in \mathcal{P}_0(1)$  be the unique  $\sigma_+$ -stable object with  $\mathbf{v}(T_i^+) = t_i$ ; similarly for  $S_i^+$  with  $\mathbf{v}(S_i^+) = s_i$  for  $i \leq 0$ . We also write  $T_i^-$  and  $S_i^-$  for the corresponding  $\sigma_-$ -stable objects.

**Proposition 6.8.** *Let  $\mathcal{W}$  be a potential wall, and assume there is an effective spherical class  $\tilde{s} \in C_{\mathcal{W}}$  with  $(\mathbf{v}, \tilde{s}) < 0$ . Then  $\mathcal{W}$  is a totally semistable wall.*

*Additionally, let  $\mathbf{v}_0$  be the minimal class in the orbit  $G_{\mathcal{H}} \cdot \mathbf{v}$ , and write  $\mathbf{v} = \mathbf{v}_l$  as in Proposition 6.7. If  $\phi^+(\mathbf{v}) > \phi^+(\mathbf{v}_0)$ , then*

$$\text{ST}_{T_l^+} \circ \text{ST}_{T_{l-1}^+} \circ \dots \circ \text{ST}_{T_1^+}(E_0)$$

FIGURE 3. The orbit of  $\mathbf{v}_0$ 

is  $\sigma_+$ -stable of class  $\mathbf{v}$ , for every  $\sigma_0$ -stable object  $E_0$  of class  $\mathbf{v}_0$ .

Similarly, if  $\phi^+(\mathbf{v}) < \phi^+(\mathbf{v}_0)$ , then

$$\mathrm{ST}_{S_{-l+1}^+}^{-1} \circ \mathrm{ST}_{S_{-l+2}^+}^{-1} \circ \cdots \circ \mathrm{ST}_{S_0^+}^{-1}(E_0)$$

is  $\sigma_+$ -stable of class  $\mathbf{v}$  for every  $\sigma_0$ -stable object of class  $\mathbf{v}_0$ .

The analogous statement holds for  $\sigma_-$ .

Note that when we are in case (b) of Proposition 6.3, the above sequence of stable spherical objects will consist of just one object.

Before the proof, we recall the following statement (see [BM11, Lemma 5.9]):

**Lemma 6.9.** *Assume that  $A, B$  are simple objects in an abelian category. If  $E$  is an extension of the form*

$$A \hookrightarrow E \twoheadrightarrow B^{\oplus r}$$

*with  $\mathrm{Hom}(B, E) = 0$ , then any quotient of  $E$  is of the form  $B^{\oplus r'}$ . Similarly, given an extension*

$$A^{\oplus r} \hookrightarrow E \twoheadrightarrow B$$

*with  $\mathrm{Hom}(E, A) = 0$ , then any subobject of  $E$  is of the form  $A^{\oplus r'}$ .*

*Proof.* We consider the former case, i.e., an extension  $A \hookrightarrow E \twoheadrightarrow B^{\oplus r}$ ; the latter case follows by dual arguments. Let  $E \twoheadrightarrow N$  be any quotient of  $E$ . Since  $A$  is a simple object, the composition  $\psi: A \hookrightarrow E \twoheadrightarrow N$  is either injective, or zero.

If  $\psi = 0$ , then  $N$  is a quotient of  $B^{\oplus r}$ , and the claim follows. If  $\psi$  is injective, let  $M$  be the kernel of  $E \twoheadrightarrow N$ . Then  $M \cap A = 0$ , and so  $M$  is a subobject of  $B^{\oplus r}$ . Since  $B$  is a simple object,  $M$  is of the form  $B^{\oplus r'}$  for some  $r' < r$ ; since  $\mathrm{Hom}(B, E) = 0$ , this is a contradiction.  $\square$

*Proof of Proposition 6.8.* Consider the first claim. By Lemma 6.2, there is a  $\sigma_0$ -stable spherical object  $\tilde{S}$  with  $(\mathbf{v}(\tilde{S}), \mathbf{v}) < 0$ . If  $E$  is a  $\sigma_0$ -stable object of class  $\mathbf{v}$ , then  $\mathrm{Hom}(\tilde{S}, E) = \mathrm{Hom}(E, \tilde{S}) = 0$ ; hence  $(\mathbf{v}(\tilde{S}), \mathbf{v}) = \mathrm{ext}^1(\tilde{S}, E) \geq 0$ , a contradiction.

To prove the construction of  $\sigma_+$ -stable objects, let us assume that we are in the case of infinitely many spherical classes. Let us also assume that  $\phi^+(\mathbf{v}) > \phi^+(\mathbf{v}_0)$ , the other case is analogous; in the notation of Proposition 6.7, this means  $\mathbf{v} = \mathbf{v}_l$  for some  $l > 0$ . We define  $E_i$  inductively by

$$E_i = \mathrm{ST}_{T_i^+}(E_{i-1}).$$



By the compatibility of the spherical twist  $\mathrm{ST}_{\tilde{S}}$  with the reflection  $\rho_{\mathbf{v}(\tilde{S})}$  and Proposition 6.7, we have  $\mathbf{v}(E_i) = \mathbf{v}_i$ . Lemma 6.9 shows that  $E_1$  is  $\sigma_+$ -stable; however, for the following induction steps, we cannot simply use Lemma 6.9 again, as neither  $E_i$  nor  $T_i^+$  are simple objects in  $\mathcal{P}_0(1)$ .

Instead, we will need a slightly stronger induction statement. Using Proposition 5.1, in particular part (b), we can define a torsion pair  $(\mathcal{T}_i, \mathcal{F}_i)$  in  $\mathcal{A}_0 := \mathcal{P}_0(1)$  as follows: we let  $\mathcal{T}_i$  be the extension closure of all  $\sigma_+$ -stable objects  $F \in \mathcal{A}_0$  with  $\phi^+(F) > \phi^+(\mathbf{t}_{i+1})$ ; by Theorem 2.13,  $\mathcal{T}_i$  is the extension-closure  $\mathcal{T}_i = \langle T_1^+, \dots, T_i^+ \rangle$ . Then let  $\mathcal{A}_i = \langle \mathcal{F}_i, \mathcal{T}_i[-1] \rangle$  (see Figure 4). We can also describe  $\mathcal{A}_{i+1}$  inductively as the tilt of  $\mathcal{A}_i$  at the torsion pair  $(\mathcal{T}, \mathcal{F})$  with  $\mathcal{T} = (T_{i+1}^+)^{\oplus k}$  and  $\mathcal{F} = (T_{i+1}^+)^{\perp}$ .

**Induction claim:** We have  $E_i \in \mathcal{F}_i$ , and both  $E_i$  and  $T_{i+1}^+$  are simple objects of  $\mathcal{A}_i$ .

By construction of the torsion pair  $(\mathcal{T}_i, \mathcal{F}_i)$ , this also shows that  $E_i$  is  $\sigma_+$ -stable.

The case  $i = 0$  follows by the assumption that  $E_0$  is  $\sigma_0$ -stable. To prove the induction step, we first consider  $T_{i+1}^+$ . By stability, we have  $T_{i+1}^+ \in \mathcal{T}_i^{\perp} = \mathcal{F}_i$ . Using stability again, we also see that any non-trivial quotient of  $T_{i+1}^+$  contained in  $\mathcal{T}_i$ , so  $T_{i+1}^+$  is a simple object of  $\mathcal{F}_i$ . Since  $T_{i+1}^+$  is stable of maximal slope in  $\mathcal{F}_i$ , there also cannot be a short exact sequence as in (13) below. Therefore, Lemma 6.10 shows that  $T_{i+1}^+$  is a simple object of  $\mathcal{A}_i$ .

Since  $E_i$  (by induction assumption) is also a simple object in  $\mathcal{A}_i$ , this shows  $\mathrm{Hom}(E_i, T_{i+1}^+) = \mathrm{Hom}(T_{i+1}^+, E_i) = 0$ . Therefore,  $\mathrm{RHom}(T_{i+1}^+, E_i) = \mathrm{Ext}^1(T_{i+1}^+, E_i)[-1]$ , and  $E_{i+1} = \mathrm{ST}_{T_{i+1}^+}^{-1}(E_i)$  fits into a short exact sequence

$$0 \rightarrow E_i \hookrightarrow E_{i+1} \twoheadrightarrow T_{i+1}^+ \otimes \mathrm{Ext}^1(T_{i+1}^+, E_i) \rightarrow 0.$$

In particular,  $E_{i+1}$  is also an object of  $\mathcal{A}_i$ . Note that

$$\mathrm{RHom}(T_{i+1}^+, E_{i+1}) = \mathrm{RHom}(\mathrm{ST}_{T_{i+1}^+}^{-1}(T_{i+1}^+), \mathrm{ST}_{T_{i+1}^+}^{-1}(E_{i+1})) = \mathrm{RHom}(T_{i+1}^+[1], E_i)$$

is concentrated in degree -2; this shows both that  $E_{i+1} \in (T_{i+1}^+)^{\perp} \subset \mathcal{A}_i$ , and that there are no extensions  $E_{i+1} \hookrightarrow F' \twoheadrightarrow T_{i+1}^{\oplus k}$ . Applying Lemma 6.10 via the inductive description of  $\mathcal{A}_{i+1}$  as a tilt of  $\mathcal{A}_i$ , this proves the induction claim.  $\square$

**Lemma 6.10.** *Let  $(\mathcal{T}, \mathcal{F})$  be a torsion pair in an abelian category  $\mathcal{A}$ , and let  $F \in \mathcal{F}$  be an object that is simple in the quasi-abelian category  $\mathcal{F}$ , and that admits no non-trivial short exact sequences*


$$(13) \quad 0 \rightarrow F \hookrightarrow F' \twoheadrightarrow T \rightarrow 0$$

*with  $F' \in \mathcal{F}$  and  $T \in \mathcal{T}$ . Then  $F$  is a simple object in the tilted category  $\mathcal{A}^{\sharp} = \langle \mathcal{F}, \mathcal{T}[-1] \rangle$ .*

*Proof.* Consider a short exact sequence  $A \hookrightarrow F \twoheadrightarrow B$  in  $\mathcal{A}^{\sharp}$ . The long exact cohomology sequence with respect to  $\mathcal{A}$  is

$$0 \rightarrow \mathcal{H}_{\mathcal{A}}^0(A) \hookrightarrow F \rightarrow F' \twoheadrightarrow \mathcal{H}_{\mathcal{A}}^1(A) \rightarrow 0$$

with  $\mathcal{H}_{\mathcal{A}}^0(A) \in \mathcal{F}$ ,  $F' \in \mathcal{F}$  and  $\mathcal{H}_{\mathcal{A}}^1(A) \in \mathcal{T}$ . Since  $F$  is a simple object in  $\mathcal{F}$ , we must have  $\mathcal{H}_{\mathcal{A}}^0(A) = 0$ . Thus we get a short exact sequence as in (13), a contradiction.  $\square$

FIGURE 4. The tilt categories  $\mathcal{A}_1$  and  $\mathcal{A}_2$ 

## 7. DIVISORIAL CONTRACTIONS IN THE NON-ISOTROPIC CASE

In this section we examine Theorem 5.7 in the case of divisorial contractions when the lattice  $\mathcal{H}_{\mathcal{W}}$  does not contain isotropic classes. The goal is to prove the following proposition.

**Proposition 7.1.** *Assume that the potential wall  $\mathcal{W}$  is non-isotropic. Then  $\mathcal{W}$  is a divisorial contraction if and only if there exists a spherical class  $\tilde{s} \in \mathcal{H}_{\mathcal{W}}$  with  $(\tilde{s}, \mathbf{v}) = 0$ . If we choose  $\tilde{s}$  to be effective, then the class of the contracted divisor  $D$  is given by  $D \equiv \theta(\tilde{s})$ .*

*If  $\tilde{S}$  is a stable spherical object of class  $\mathbf{v}(\tilde{S}) = \tilde{s}$ , then  $D$  can be described as a Brill-Noether divisor of  $\tilde{S}$ : it is given either by the condition  $\text{Hom}(\tilde{S}, \_) \neq 0$ , or by  $\text{Hom}(\_, \tilde{S}) \neq 0$ .*

One can use more general results of Markman in [Mar09] to show that a divisorial contraction implies the existence of an orthogonal spherical class in the non-isotropic case. We will instead give a categorical proof in our situation.

We first treat the case in which there exists a  $\sigma_0$ -stable object of class  $\mathbf{v}$ :

**Lemma 7.2.** *Assume that  $\mathcal{H}$  is non-isotropic, and that  $\mathcal{W}$  is a potential wall associated to  $\mathcal{H}$ . If  $\mathbf{v}$  is a minimal class of a  $G_{\mathcal{H}}$ -orbit, and if there is no spherical class  $\tilde{s} \in \mathcal{H}_{\mathcal{W}}$  with  $(\tilde{s}, \mathbf{v}) = 0$ , then the set of  $\sigma_0$ -stable objects in  $M_{\sigma_+}(\mathbf{v})$  has complement of codimension at least two.*

In particular,  $\mathcal{W}$  cannot induce a divisorial contraction.

*Proof.* The argument is similar to Lemma 6.5; an additional ingredient is Namikawa's and Wierzb'a's characterization of divisorial contractions recalled in Theorem 3.6.

For contradiction, assume that there is a divisor  $D \subset M_{\sigma_+}$  of objects that are strictly semistable with respect to  $\sigma_0$ . Let morphism  $\pi^+: M_{\sigma_+} \rightarrow \overline{M}$  be the morphism induced by  $\ell_{\sigma_0}$ ; it is either an isomorphism or a divisorial contraction. Let  $D \subset M_{\sigma_+}$  be an irreducible divisor of strictly  $\sigma_0$ -semistable objects. The morphism  $\pi^+$  may or may not be contracted by

$\pi^+$ ; by Theorem 3.6, we have  $\dim \pi^+(D) \geq \dim D - 1 = \dim M_{\sigma_+}(\mathbf{v}) - 2 = \mathbf{v}^2$  in either case.

On the other hand, consider the restriction of the universal family  $\mathcal{E}$  on  $M_{\sigma_+}(\mathbf{v})$  to the divisor  $D$ , and its relative Harder-Narasimhan filtration with respect to  $\sigma_-$ . As before, this induces a rational map

$$\mathrm{HN}_D: D \dashrightarrow M_{\sigma_-}(\mathbf{a}_1) \times \cdots \times M_{\sigma_-}(\mathbf{a}_l).$$

Again, let  $I \subset \{1, \dots, l\}$  be the subset of indices  $i$  with  $\mathbf{a}_i^2 > 0$ , and  $\mathbf{a} = \sum_{i \in I} \mathbf{a}_i$ . The proof based on Figure 2 still applies, and shows  $\mathbf{a}^2 \leq \mathbf{v}^2$ .

If  $I \neq \{1, \dots, l\}$ , there exists a class  $\mathbf{a}_j$  appearing in the Harder-Narasimhan filtration of the form  $\mathbf{a}_j = m\tilde{\mathbf{s}}$ ,  $\tilde{\mathbf{s}}^2 = -2$ . Under the assumptions, we now have the *strict* inequality  $(\tilde{\mathbf{s}}, \mathbf{v}) > 0$ , from which we obtain  $\mathbf{a}^2 \leq (\mathbf{v} - \tilde{\mathbf{s}})^2 \leq \mathbf{v}^2 - 3$ .

Otherwise, if  $I = \{1, \dots, l\}$ , we have  $|I| > 1$ , and we can apply Lemma 6.4; in either case we obtain

$$\sum_{i=1}^l \dim_{M_{\sigma_-}(\mathbf{a}_i)} = \sum_{i \in I} (\mathbf{a}_i^2 + 2) < \mathbf{v}^2 = \dim \pi^+(D).$$

As before, this is a contradiction to the observation that any curve contracted by  $\mathrm{HN}_D$  is also contracted by  $\pi^+$ .  $\square$

The case of totally semistable walls can be reduced to the previous one:

**Corollary 7.3.** *Assume that  $\mathcal{H}$  is non-isotropic, and that there does not exist a spherical class  $\tilde{\mathbf{s}} \in \mathcal{H}$  with  $(\tilde{\mathbf{s}}, \mathbf{v}) = 0$ . Then a potential wall associated to  $\mathcal{H}$  cannot induce a divisorial contraction.*

In fact, we will later see that all potential walls associated to  $\mathcal{H}$  are mapped to the same wall in the movable cone of the moduli space; thus they have to exhibit identical birational behavior.

*Proof.* As before, consider the minimal class  $\mathbf{v}_0$  of the orbit  $G_{\mathcal{H}} \cdot \mathbf{v}$ , in the sense of Definition 6.6. By Lemma 7.2, there is an open subset  $U \subset M_{\sigma_+}(\mathbf{v}_0)$  of objects that are  $\sigma_0$ -stable that has complement of codimension at least two.

Let  $\Phi$  be the composition of spherical twists given by Proposition 6.8, such that  $\Phi(E_0)$  is  $\sigma_+$ -stable of class  $\mathbf{v}$  for every  $[E_0] \in U$ . Observe that  $\Phi(E_0)$  has a Jordan-Hölder filtration such that  $E_0$  is one of its filtration factors (the other factors are stable spherical objects). Therefore, the induced map  $\Phi_*: U \rightarrow M_{\sigma_+}(\mathbf{v})$  is injective, and the image does not contain any curve of S-equivalent objects with respect to  $\sigma_0$ . Also,  $\Phi_*(U)$  has complement of codimension at least two (see e.g. [GHJ03, Proposition 21.6]). Since  $\ell_{\sigma_0}$  does not contract any curves in  $\Phi_*(U)$ , it cannot contract any divisors in  $M_{\sigma_+}(\mathbf{v})$ .  $\square$

The next step is to construct the divisorial contraction when there exists an orthogonal spherical class. To clarify the logic, we first treat the simpler case of a wall that is not totally semistable:

**Lemma 7.4.** *Assume  $\mathcal{H}$  is non-isotropic,  $\mathcal{W}$  a potential wall associated to  $\mathcal{H}$ , and that  $\mathbf{v}$  is a minimal class of a  $G_{\mathcal{H}}$ -orbit. If there exists a spherical class  $\tilde{\mathbf{s}} \in \mathcal{H}$  with  $(\tilde{\mathbf{s}}, \mathbf{v}) = 0$ , then  $\mathcal{W}$  induces a divisorial contraction.*

*If we assume that  $\tilde{\mathbf{s}}$  is effective, then the contracted divisor  $D \subset M_{\sigma_+}(\mathbf{v})$  has class  $\theta(\tilde{\mathbf{s}})$ . The HN filtration of a generic element  $[E] \in D$  with respect to  $\sigma_-$  is of the form*

$$0 \rightarrow \tilde{S} \hookrightarrow E \twoheadrightarrow F \rightarrow 0 \quad \text{or} \quad 0 \rightarrow F \hookrightarrow E \twoheadrightarrow \tilde{S} \rightarrow 0,$$

where  $\tilde{S}$  and  $F$  are  $\sigma_0$ -stable objects of class  $\tilde{s}$  and  $\mathbf{v} - \tilde{s}$ , respectively.

*Proof.* As before, we only treat the case when  $\mathcal{H}$  admits infinitely many spherical classes. In that case, we must have  $\tilde{s} = s$  or  $\tilde{s} = t$ ; we may assume  $\tilde{s} = s$ , and the other case will follow by dual arguments.

We first prove that  $\mathbf{v} - s$  is a minimal class in its  $G_{\mathcal{H}}$ -orbit by a straightforward computation. If  $\mathbf{v}^2 = 2$ , then  $(\mathbf{v} - s)^2 = 0$  in contradiction to the assumption; therefore  $\mathbf{v}^2 \geq 4$ . If we write  $\mathbf{v} = xs + yt$ , then  $(\mathbf{v}, s) = 0$  gives  $y = \frac{2}{m}x$ . Plugging in  $\mathbf{v}^2 \geq 4$  gives  $x^2(1 - \frac{4}{m^2}) \geq 2$ . Since  $m \geq 3$ , we obtain

$$x^2 \left(1 - \frac{4}{m^2}\right)^2 > x^2 \left(1 - \frac{4}{m^2}\right) \frac{1}{2} \geq 1,$$

and therefore

$$(t, \mathbf{v} - s) = m(x - 1) - 2\frac{2}{m}x = mx \left(1 - \frac{4}{m^2}\right) - m \geq 0.$$

Also,  $(s, \mathbf{v} - s) = 2 > 0$ , and therefore  $\mathbf{v} - s$  has positive pairing with every effective spherical class.

By Lemma 6.5, the generic element  $F \in M_{\sigma_+}(\mathbf{v} - s)$  is also  $\sigma_0$ -stable. Since  $(s, \mathbf{v} - s) = 2$  and  $\text{Hom}(F, S) = \text{Hom}(S, F) = 0$ , there is a family of extensions

$$0 \rightarrow S \hookrightarrow E_p \twoheadrightarrow F \rightarrow 0$$

parametrized by  $p \in \mathbb{P}^1 \cong \mathbb{P}(\text{Ext}^1(F, S))$ . By Lemma 6.9, they are  $\sigma_+$ -stable. Since all  $E_p$  are  $S$ -equivalent to each other, the contraction morphism  $\pi^+ : M_{\sigma_+}(\mathbf{v}) \rightarrow \overline{M}$  associated to  $\mathcal{W}$  will contract the image of this rational curve. Varying  $F \in M_{\sigma_0}^{st}(\mathbf{v} - s)$ , these span a family of dimension  $1 + (\mathbf{v} - s)^2 + 2 = \mathbf{v}^2 + 1$ ; this is a divisor in  $M_{\sigma_+}(\mathbf{v})$  contracted by  $\pi^+$ .

Since  $\pi^+$  has relative Picard-rank equal to one, it cannot contract any other component.  $\square$

The following lemma treats the general case, for which we will first set up notation. As before, we let  $\mathbf{v}_0$  be the minimal class in the  $G_{\mathcal{H}}$ -orbit of  $\mathbf{v}$ . By  $\tilde{s}_0$  we denote the effective spherical class with  $(\mathbf{v}_0, \tilde{s}_0) = 0$ ; we have  $\tilde{s}_0 = t$  or  $\tilde{s}_0 = s$ . Accordingly, in the list of the  $G_{\mathcal{H}}$ -orbit of  $\mathbf{v}$  given by Proposition 6.7, we have either  $v_{2i} = v_{2i+1}$ , or  $v_{2i} = v_{2i-1}$  for all  $i$ , since  $\mathbf{v}_0$  is fixed under the reflection  $\rho_{\tilde{s}_0}$  at  $\tilde{s}_0$ . We choose  $l$  such that  $\mathbf{v} = \mathbf{v}_l$ , and such that the corresponding sequence of reflections sends  $\tilde{s}_0$  to  $\tilde{s}$ :

$$\tilde{s} = \begin{cases} \rho_{t_l} \circ \rho_{t_{l-1}} \circ \cdots \circ \rho_{t_0}(\tilde{s}_0) & \text{if } l > 0 \\ \rho_{s_l} \circ \rho_{s_{l/1}} \circ \cdots \circ \rho_{s_{-1}}(\tilde{s}_0) & \text{if } l < 0 \end{cases}$$

Depending on the ordering of the slopes  $\phi^+(\mathbf{v}), \phi^+(\mathbf{v}_0)$ , we let  $\Phi$  be the composition of spherical twists appearing in Proposition 6.8.

**Lemma 7.5.** *Assume that  $\mathcal{H}$  is non-isotropic, and  $\mathcal{W}$  a corresponding potential wall. If there is an effective spherical  $\tilde{s} \in C_{\mathcal{W}}$  with  $(\mathbf{v}, \tilde{s}) = 0$ , then  $\mathcal{W}$  induces a divisorial contraction.*

*The contracted divisor  $D$  has class  $\theta(\tilde{s})$ . The Jordan-Hölder filtration of a generic element  $E \in D$  is a refinement of a short exact sequence of the form*

$$(14) \quad 0 \rightarrow \tilde{S} \hookrightarrow E \twoheadrightarrow F \rightarrow 0 \quad \text{or} \quad 0 \rightarrow F \hookrightarrow E \twoheadrightarrow \tilde{S} \rightarrow 0,$$

where  $F$  and  $\tilde{S}$  are  $\sigma_+$ -stable objects of class  $\mathbf{v} - \tilde{s}$  and  $\tilde{s}$ , respectively.

*In addition, there exists an open subset  $U^+ \subset M_{\sigma_+}(\mathbf{v}_0)$ , with complement of codimension two, such that  $\Phi(E_0)$  is  $\sigma_+$ -stable for every  $\sigma_+$ -stable object  $E_0 \in U$ .*

*Proof.* We rely on the construction in the proof of Proposition 6.8, and in particular on the induction claim proved therein.

Let  $\tilde{S}_0$  be the stable spherical object of class  $\tilde{s}_0$ ; we have  $\tilde{S}_0 = S$  or  $\tilde{S}_0 = T$ . As in the proof of Lemma 7.4, one shows that  $\mathbf{v}_0 - \tilde{s}_0$  is the minimal class in its  $G_{\mathcal{H}}$ -orbit.

Let  $F_0$  be a generic  $\sigma_0$ -stable object of class  $\mathbf{v}_0 - \tilde{s}_0$ . Applying Proposition 6.8 to the class  $\mathbf{v} - \tilde{s}$ , we see that  $F := \Phi(F_0)$  is  $\sigma_+$ -stable of that class.

Up to duality, we may again assume that  $\Phi$  is of the form  $\mathrm{ST}_{T_l}^+ \circ \cdots \circ \mathrm{ST}_{T_1}^+$ . Inductively, one shows that  $\Phi(S) = T_{l+1}^+$  and  $\Phi(T) = T_l[-1]$ . These are both simple objects of the category  $\mathcal{A}_l$  defined by tilting in the proof of Proposition 6.8; therefore,  $\tilde{S} := \Phi(\tilde{S}_0)$  is simple in  $\mathcal{A}_l$ . By the induction claim,  $F = \Phi(F_0)$  is also a simple object in this category. In particular,  $\mathrm{Hom}(\tilde{S}, F) = \mathrm{Hom}(F, \tilde{S}) = 0$  and  $\mathrm{ext}^1(\tilde{S}, F) = 2$ . Applying Lemma 6.9 again, and using the compatibility of  $\mathcal{A}_l$  with stability, we obtain a stable extension of the form (14).

This gives a divisor contracted by  $\pi^+$ , and we can proceed as in the previous lemma.

Let  $D_0 \subset M_{\sigma_+}(\mathbf{v}_0)$  be the contracted divisor for the class  $\mathbf{v}_0$ . The above proof also shows that for a generic object  $E_0 \in D_0$  (whose form is given by Lemma 7.4), the object  $\Phi(E_0)$  is a  $\sigma_+$ -stable (contained in the contracted divisor  $D$ ). Thus we can take  $U^+$  to be the union of all  $\sigma_0$ -stable objects in  $M_{\sigma_+}(\mathbf{v}_0)$ , with the open subset of  $D_0$  of objects of the form given in Lemma 7.4.  $\square$

*Proof of Proposition 7.1.* The statements follow from Corollary 7.3 and Lemma 7.5.  $\square$

## 8. ISOTROPIC WALLS ARE UHLENBECK WALLS

In this section, we study potential walls  $\mathcal{W}$  in the case where  $\mathcal{H}$  admits an isotropic class  $\mathbf{w} \in \mathcal{H}$ ,  $\mathbf{w}^2 = 0$ . Following an idea of Minamide, Yanagida, and Yoshioka [MY11b], we study the wall  $\mathcal{W}$  via a Fourier-Mukai transform after which  $\mathbf{w}$  becomes the class of a point. Then  $\sigma_+$  corresponds to Gieseker-stability and, as proven in [Lo12], the wall corresponds to the contraction to the Uhlenbeck compactification, as constructed by Jun Li in [Li93].

Parts of this section are well-known. In particular, [Yos99, Proposition 0.5] deals with the existence of stable locally-free sheaves. For other general results, see [Yos01].

**The Uhlenbeck compactification.** We start with the following observation:

**Lemma 8.1.** *Assume that there exists an isotropic class in  $\mathcal{H}$ . Then there are two effective, primitive, isotropic classes  $\mathbf{w}_0$  and  $\mathbf{w}_1$  in  $\mathcal{H}$ , such that, for a generic stability condition  $\sigma_0 \in \mathcal{W}$ , we have*

- (a)  $M_{\sigma_0}(\mathbf{w}_0) = M_{\sigma_0}^{st}(\mathbf{w}_0)$ , and
- (b) either  $M_{\sigma_0}(\mathbf{w}_1) = M_{\sigma_0}^{st}(\mathbf{w}_1)$ , or there exists a  $\sigma_0$ -stable spherical object  $S$ , with Mukai vector  $\mathbf{s}$ , such that  $(\mathbf{s}, \mathbf{w}_1) < 0$  and  $\mathcal{W}$  is a totally semistable wall for  $\mathbf{w}_1$ .

Any positive class  $\mathbf{v}' \in P_{\mathcal{H}}$  satisfies  $(\mathbf{v}', \mathbf{w}_i) \geq 0$  for  $i = 1, 2$ .

*Proof.* Let  $\tilde{\mathbf{w}} \in \mathcal{H}$  be primitive isotropic class; up to replacing  $\tilde{\mathbf{w}}$  by  $-\tilde{\mathbf{w}}$ , we may assume it to be effective. We complete  $\tilde{\mathbf{w}}$  to a basis  $\{\tilde{\mathbf{v}}, \tilde{\mathbf{w}}\}$  of  $\mathcal{H}_{\mathbb{Q}}$ . Then, for all  $(a, b) \in \mathbb{Q}$ , we have

$$(a\tilde{\mathbf{v}} + b\tilde{\mathbf{w}})^2 = a \cdot (a\tilde{\mathbf{v}}^2 + b(\tilde{\mathbf{v}}, \tilde{\mathbf{w}})).$$

This shows the existence of a second integral isotropic class. If we choose it to be effective, then the positive cone, generated by  $P_{\mathcal{H}}$ , is exactly the cone  $\mathbb{R}_{\geq 0} \cdot \mathbf{w}_0 + \mathbb{R}_{\geq 0} \cdot \mathbf{w}_1$ ; from this, the claim  $(\mathbf{v}', \mathbf{w}_i) \geq 0$  follows easily.

By Theorem 2.13, we have  $M_{\sigma_0}(\tilde{\mathbf{w}}) \neq \emptyset$ . If  $\mathcal{W}$  does not coincide with a wall for  $\tilde{\mathbf{w}}$ , then we can take  $\mathbf{w}_0 = \tilde{\mathbf{w}}$ , and claim (a) will be satisfied.

Otherwise, let  $\sigma \in \text{Stab}^\dagger(X, \alpha)$  be a generic stability condition nearby  $\mathcal{W}$ ; by [BM12, Lemma 6.2], we have  $M_\sigma(\tilde{\mathbf{w}}) = M_\sigma^{st}(\tilde{\mathbf{w}}) \neq \emptyset$ .

Up to applying a Fourier-Mukai equivalence, we may assume that  $\tilde{\mathbf{w}} = (0, 0, 1)$  is the Mukai vector of a point on a twisted K3 surface; then we can apply the classification of walls for isotropic classes in [Bri08, Theorem 12.1], extended to twisted surfaces in [HMS08]. If  $\mathcal{W}$  is a totally semistable wall for  $\tilde{\mathbf{w}}$ , then we are in cases  $(A^+)$  and  $(A^-)$  of [Bri08, Theorem 12.1]. Hence, there exists a spherical  $\sigma_0$ -stable twisted vector bundle  $S$  such that  $S$  or  $S[2]$  is a JH factor for the skyscraper sheaf  $k(x)$ , for any  $x \in M_\sigma(\tilde{\mathbf{w}})$ . Moreover, the other non-isomorphic JH factor is either  $\text{ST}_S(k(x))$ , or  $\text{ST}_S^{-1}(k(x))$ . In both cases, the Mukai vector  $\mathbf{w}_0$  of this last JH factor is primitive and isotropic, and  $\mathcal{W}$  is not a wall for  $\mathbf{w}_0$ .

Finally, if  $\mathcal{W}$  is a wall for  $\tilde{\mathbf{w}}$ , but not a totally semistable wall, it must be a wall of type  $(C_k)$ . However, the rank two lattice corresponding to a wall of type  $(C_k)$  is negative semi-definite; on the other hand, by Proposition 5.1, claim (d), it must coincide with  $\mathcal{H}$ , which has signature  $(1, -1)$ ; this is a contradiction.  $\square$

Let  $\mathbf{w}_0, \mathbf{w}_1 \in C_{\mathcal{W}}$  be the effective, primitive, isotropic classes given by the above lemma, and let  $Y := M_{\sigma_0}(\mathbf{w}_0)$ . Then  $Y$  is a K3 surface and, by [Muk87a, Că102, Yos06, HS06], there exist a class  $\alpha' \in \text{Br}(Y)$  and a Fourier-Mukai transform

$$\Phi: D^b(X, \alpha) \xrightarrow{\sim} D^b(Y, \alpha')$$

such that  $\Phi(\mathbf{w}_0) = (0, 0, 1)$ . By construction, skyscraper sheaves of points on  $Y$  are  $\Phi_*(\sigma_0)$ -stable. By Bridgeland's Theorem 2.7 (generalized to twisted K3 surfaces in [HMS08]), there exist divisor classes  $\omega, \beta \in \text{NS}(Y)_{\mathbb{Q}}$ , with  $\omega$  ample, such that up to the  $\text{GL}_2$ -action,  $\Phi_*(\sigma_0)$  is given by  $\sigma_{\omega, \beta}$ . In particular, the category  $\mathcal{P}_{\omega, \beta}(1)$  is the extension-closure of skyscraper sheaves of points, and the shifts  $F[1]$  of  $\mu_\omega$ -stable torsion-free sheaves  $F$  with slope  $\mu_{\omega, \beta}(F) = \omega \cdot \beta$ . Since  $\sigma_0$  by assumption does not lie on any other wall with respect to  $\mathbf{v}$ , the divisor  $\omega$  is generic with respect to  $\Phi_*(\mathbf{v})$ .

By abuse of notation, we will from now on write  $(X, \alpha)$  instead of  $(Y, \alpha')$ ,  $\mathbf{v}$  instead of  $\Phi_*(\mathbf{v})$ , and  $\sigma_0$  instead of  $\sigma_{\omega, \beta}$ . Let  $\sigma_+ = \sigma_{\omega, \beta - \epsilon}$  and  $\sigma_- = \sigma_{\omega, \beta + \epsilon}$ ; here  $\epsilon$  is a sufficiently small positive multiple of  $\omega$ .

**Proposition 8.2** ([Lo12, LQ11]). *An object of class  $\mathbf{v}$  is  $\sigma_+$ -stable if and only if it is the shift  $F[1]$  of a  $\beta$ -twisted Gieseker stable sheaf  $F$  on  $(X, \alpha)$ ; the shift  $[1]$  induces the following identification of moduli spaces:*

$$M_{\sigma_+}(\mathbf{v}) = M_{\omega, \beta}(-\mathbf{v}).$$

*Moreover, the contraction morphism  $\pi^+$  induced via Theorem 2.16 for generic  $\sigma_0 \in \mathcal{W}$  is the Li-Gieseker-Uhlenbeck morphism to the Uhlenbeck compactification.*

*Finally, an object  $F$  of class  $\mathbf{v}$  is  $\sigma_-$ -stable if and only if it is the shift  $F^\vee[2]$  of the derived dual of a  $(-\beta)$ -twisted Gieseker stable sheaf on  $(X, -\alpha)$ .*

*Proof.* The identification of  $M_{\sigma_+}(\mathbf{v})$  with the Gieseker-moduli space is well-known, and follows with the same arguments as in [Bri08, Proposition 14.2]. The identification of  $\pi^+$  with the morphism to the Uhlenbeck space follows from the combination of [Lo12, Theorem 3.1] with Theorem 2.16.

The claim about  $F^\vee[2]$  being  $\sigma_-$ -stable follows by combining Proposition 2.10 with the previous statements (see also see [MY11a, Proposition 2.2.7] in the case  $\alpha = 0$ ).  $\square$



In other words, the coarse moduli space  $M_{\sigma_0}(\mathbf{v})$  is isomorphic to the Uhlenbeck compactification ([Li93, Yos06]) of the moduli space of slope-stable vector bundles on  $(X, \alpha)$ . Given a (twisted) Gieseker-stable sheaf  $F \in M_{\omega, \beta}(-\mathbf{v})$ , the  $\sigma_+$ -stable object  $F[1]$  becomes strictly semistable with respect to  $\sigma_0$  if and only if  $F$  is not locally free, or if  $F$  is not slope-stable.

In particular, when the rank of  $-\mathbf{v}$  equals one, then the contraction morphism  $\pi_+$  is the Hilbert-Chow morphism  $\text{Hilb}^n(X) \rightarrow \text{Sym}^n(X)$ ; see also [BM12, Example 9.1].

**Totally semistable isotropic walls.** The goal of this section is to estimate the locus in  $M_{\sigma_+}(\mathbf{v}) = M_{\omega}(-\mathbf{v})$  of sheaves which are neither slope-semistable nor locally-free. We start with the existence of a unique spherical stable object in the case the wall is totally semistable:

**Lemma 8.3.** *Assume that  $\mathcal{W}$  is a totally semistable wall for  $\mathbf{v}$ .*

- (a) *There exists a unique spherical  $\sigma_0$ -stable object  $S \in \mathcal{P}_{\sigma_0}(1)$ .*
- (b) *Let  $E \in M_{\sigma_+}(\mathbf{v})$  be a generic object. Then its HN filtration with respect to  $\sigma_-$  has length 2 and takes the form*

$$(15) \quad S^{\oplus a} \rightarrow E \rightarrow F, \quad \text{or} \quad F \rightarrow E \rightarrow S^{\oplus a},$$

*with  $a \in \mathbb{Z}_{>0}$ . The  $\sigma_-$ -semistable object  $F$  is generic in  $M_{\sigma_-}(\mathbf{v}')$ , for  $\mathbf{v}' := \mathbf{v}(F)$ , and  $\dim M_{\sigma_-}(\mathbf{v}') = \dim M_{\sigma_+}(\mathbf{v}) = \mathbf{v}^2 + 2$ .*

The idea of the proof is very similar to the one in Lemma 6.5. The only difference is that we cannot use a completely numerical criterion like Lemma 6.4 and we will replace it by Mukai's Lemma 6.1.

*Proof of Lemma 8.3.* We first prove (a). We consider again the two maps

$$\begin{aligned} \pi^+ : M_{\sigma_+}(\mathbf{v}) &\rightarrow \overline{M}, \\ \text{HN} : M_{\sigma_+}(\mathbf{v}) &\dashrightarrow M_{\sigma_-}(\mathbf{a}_1) \times \cdots \times M_{\sigma_-}(\mathbf{a}_m). \end{aligned}$$

The first one is induced by  $\ell_{\sigma_0}$  and the second by the existence of relative HN filtrations. By [HL10, Section 4.5], we have, for all  $i = 1, \dots, m$  and for all  $A_i \in M_{\sigma_-}(\mathbf{a}_i)$ ,

$$\dim M_{\sigma_-}(\mathbf{a}_1) \leq \text{ext}^1(A_i, A_i).$$

Hence, by Mukai's Lemma 6.1, we deduce

$$(16) \quad \dim M_{\sigma_+}(\mathbf{v}) \geq \sum_{i=1}^m \dim M_{\sigma_-}(\mathbf{a}_i).$$

Equation (16) is the analogue of (12) in the non-isotropic case. Since any curve contracted by HN is also contracted by  $\pi^+$ , it follows that

$$\sum_{i=1}^m \dim M_{\sigma_-}(\mathbf{a}_i) \geq \dim \overline{M} = \dim M_{\sigma_+}(\mathbf{v}).$$

Therefore equality holds, and HN is a dominant map.

This shows that the projections

$$M_{\sigma_+} \dashrightarrow M_{\sigma_-}(\mathbf{a}_i)$$

are dominant. By Theorem 3.8,  $M_{\sigma_-}(\mathbf{a}_i)$  has symplectic singularities. Hence, we deduce that either  $M_{\sigma_-}(\mathbf{a}_i)$  is a point, or  $\dim M_{\sigma_-}(\mathbf{a}_i) = \dim M_{\sigma_+}(\mathbf{v}) = \mathbf{v}^2 + 2$ . Since  $m \geq 2$ , by Lemma 6.2 this shows the existence of a spherical  $\sigma_0$ -stable object in  $\mathcal{P}_{\sigma_0}(1)$ . By Proposition 6.3, there can only be one such spherical object.

To prove (b), we first observe that by uniqueness (and by Lemma 6.2 again), all  $\sigma_-$ -spherical objects appearing in a HN filtration of a generic element  $E \in M_{\sigma_+}(\mathbf{v})$  must be  $\sigma_0$ -stable as well. As a consequence, the length of a HN filtration of  $E$  with respect to  $\sigma_-$  must be 2 and have the form (15). Since the maps  $M_{\sigma_+} \dashrightarrow M_{\sigma_-}(\mathbf{a}_i)$  are dominant, the  $\sigma_-$ -semistable object  $F$  is generic.  $\square$

We can now prove the first implication for the characterization of totally semistable walls in the isotropic case. We let  $\mathbf{s} := \mathbf{v}(S)$ , where  $S$  is the unique  $\sigma_0$ -stable object in  $\mathcal{P}_{\sigma_0}(1)$ .

**Proposition 8.4.** *Let  $\mathcal{W}$  be a totally semistable wall for  $\mathbf{v}$ . Then either there exist an isotropic vector  $\mathbf{w}$  with  $(\mathbf{w}, \mathbf{v}) = 1$ , or the effective spherical class  $\mathbf{s}$  satisfies  $(\mathbf{s}, \mathbf{v}) < 0$ .*

*Proof.* We continue to use the notation of Lemma 8.3. In particular, consider  $\mathbf{v}' = \mathbf{v} - a\mathbf{s}$  with  $a > 0$ .

If  $(\mathbf{v}')^2 > 0$ , then by Lemma 8.3 and Theorem 2.13(b), we have  $(\mathbf{v}')^2 = \mathbf{v}^2$ . Since  $\mathbf{v}' = \mathbf{v} - a\mathbf{s}$ ,  $a > 0$ , this implies  $(\mathbf{s}, \mathbf{v}) < 0$ .

So we may assume  $\mathbf{v}'^2 = 0$ . Then  $\mathbf{v}^2 = 0 + 2a(\mathbf{v}', \mathbf{s}) - 2a^2$ , and it follows that  $(\mathbf{v}', \mathbf{s}) > 0$ . In the notation of Lemma 8.1, this means that  $\mathbf{v}'$  is a positive multiple of  $\mathbf{w}_0$ , which we can take to be the class of a point:  $\mathbf{v}' = c\mathbf{w}_0 = c(0, 0, 1)$ .

Then the coarse moduli space  $M_{\sigma_0}(\mathbf{v}')$  is the symmetric product  $\text{Sym}^c X$ ; if we define  $n$  by  $\mathbf{v}^2 = 2n - 2$ , then the equality of dimensions in Lemma 8.3 becomes  $c = n$ . Therefore

$$2n - 2 = \mathbf{v}^2 = (a\mathbf{s} + n\mathbf{w}_0)^2 = -2a^2 + 2an(\mathbf{s}, \mathbf{w}_0)$$

or, equivalently,

$$(17) \quad a(n(\mathbf{s}, \mathbf{w}_0) - a) = n - 1.$$

Recall that  $(\mathbf{s}, \mathbf{w}_0) > 0$ ; by the geometric-arithmetic mean inequality, this implies  $(\mathbf{s}, \mathbf{w}_0) = 1$ . (Concretely, this means the spherical object  $S$  is the shift of a line bundle.)

In this case, solving (17) for  $a$  gives the two solutions  $a = 1$  and  $a = n - 1$ . In the former case,  $(\mathbf{v}, \mathbf{w}_0) = 1$ . In the latter case, observe that  $\mathbf{w}_1 = \mathbf{w}_0 + \mathbf{s}$ , and  $(\mathbf{v}, \mathbf{w}_1) = 1$  follows directly.  $\square$

The converse statement follows from Proposition 6.8 above, and Lemma 8.5 below.

**Lemma 8.5.** *Let  $\mathcal{W}$  be a potential wall. If there exists an isotropic class  $\mathbf{w} \in \mathcal{H}_{\mathcal{W}}$  with  $(\mathbf{w}, \mathbf{v}) = 1$ , then  $\mathcal{W}$  is a totally semistable wall.*

*Proof.* Note that by Lemma 8.1, the primitive class  $\mathbf{w}$  is automatically effective. Let  $\sigma_0 \in \mathcal{W}$  be a generic stability condition. If  $M_{\sigma_0}^{st}(\mathbf{w}) \neq \emptyset$ , then we can assume  $\mathbf{w} = (0, 0, 1)$ . In this case  $-\mathbf{v}$  has rank one,  $M_{\sigma_+}(\mathbf{v})$  is the Hilbert scheme, and  $\mathcal{W}$  is the Hilbert-Chow wall discussed in [BM12, Example 9.1]; in particular, it is totally semistable.

Otherwise,  $M_{\sigma_0}^{st}(\mathbf{w}) = \emptyset$ ; hence, in the notation of Lemma 8.1, we are in the case  $\mathbf{w} = \mathbf{w}_1$ , and there exists a  $\sigma_0$ -stable spherical object  $S$ , with Mukai vector  $\mathbf{s}$ , such that  $(\mathbf{s}, \mathbf{w}_1) < 0$ .

Write  $\mathbf{w}_1 = \mathbf{w}_0 + r\mathbf{s}$ , where  $r = (\mathbf{s}, \mathbf{w}_0) \in \mathbb{Z}_{>0}$ . Then

$$1 = (\mathbf{v}, \mathbf{w}_1) = (\mathbf{v}, \mathbf{w}_0) + r(\mathbf{v}, \mathbf{s}).$$

By Lemma 8.1,  $(\mathbf{v}, \mathbf{w}_0)$  is strictly positive, and so  $(\mathbf{v}, \mathbf{s}) \leq 0$ . If the inequality is strict, Proposition 6.8 applies. Otherwise,  $(\mathbf{v}, \mathbf{s}) = 0$  and  $(\mathbf{v}, \mathbf{w}_1) = (\mathbf{v}, \mathbf{w}_0) = 1$ ; thus we are again in the case of the Hilbert-Chow wall, and  $\mathcal{W}$  is a totally semistable wall for  $\mathbf{v}$ .  $\square$

**Divisorial contractions.** We now deal with divisorial contractions for isotropic walls. The case of a flopping wall, a fake wall, and no wall will be examined in Section 9.

**Proposition 8.6.** *Let  $\mathcal{W}$  be a wall inducing a divisorial contraction. Assume that  $(\mathbf{v}, \mathbf{w}) \neq 1, 2$ , for all isotropic vectors  $\mathbf{w} \in \mathcal{H}$ . Then there exists an effective spherical class  $\mathbf{s} \in \mathcal{H}$  with  $(\mathbf{s}, \mathbf{v}) = 0$ .*

*Proof.* The proof is similar to the one of Lemma 7.2: in particular, we are going to use Theorem 3.6. Let  $D \subset M_{\sigma_+}(\mathbf{v})$  be an irreducible divisor contracted by  $\pi^+ : M_{\sigma_+}(\mathbf{v}) \rightarrow \overline{M}$ . We know that  $\dim \pi^+(D) = \mathbf{v}^2$ . Consider the rational map

$$\mathrm{HN}_D : D \dashrightarrow M_{\sigma_-}(\mathbf{a}_1) \times \cdots \times M_{\sigma_-}(\mathbf{a}_l)$$

induced by the relative HN filtration with respect to  $\sigma_-$ . We let  $I \subset \{1, \dots, l\}$  be the subset of indices  $i$  with  $\mathbf{a}_i^2 > 0$ , and  $\mathbf{a} = \sum_{i \in I} \mathbf{a}_i$ .

**Step 1.** There exists an effective spherical class  $\mathbf{s} \in \mathcal{H}$ .

Indeed, assume for a contradiction that it does not exist. Then we can write  $\mathbf{v} = n_0 \mathbf{w}_0 + n_1 \mathbf{w}_1 + \mathbf{a}$ , with  $(\mathbf{w}_0, \mathbf{w}_1) > 0$  and  $(\mathbf{v}, \mathbf{w}_i) \geq 3$ , for  $i = 0, 1$ . We have

$$\begin{aligned} \mathbf{a}^2 &= (\mathbf{v} - n_0 \mathbf{w}_0 - n_1 \mathbf{w}_1)^2 = \mathbf{v}^2 - 2n_0(\mathbf{v}, \mathbf{w}_0) - 2n_1(\mathbf{v}, \mathbf{w}_1) + 2n_0n_1(\mathbf{w}_0, \mathbf{w}_1) \\ &= \mathbf{v}^2 - n_0(2(\mathbf{v}, \mathbf{w}_0) - (n_1 \mathbf{w}_1, \mathbf{w}_0)) - n_1(2(\mathbf{v}, \mathbf{w}_1) - (n_0 \mathbf{w}_0, \mathbf{w}_1)). \end{aligned}$$

But  $(\_, \mathbf{w}_i)$  is positive on effective classes, for  $i = 0, 1$ . Hence,  $(\mathbf{v}, \mathbf{w}_0) > (n_1 \mathbf{w}_1, \mathbf{w}_0)$  and  $(\mathbf{v}, \mathbf{w}_1) > (n_0 \mathbf{w}_0, \mathbf{w}_1)$ . Therefore,

$$\mathbf{a}^2 \leq \mathbf{v}^2 - n_0((\mathbf{v}, \mathbf{w}_0) + 1) - n_1((\mathbf{v}, \mathbf{w}_1) + 1) \leq \mathbf{v}^2 - 4n_0 - 4n_1.$$

So the dimension of  $M_{\sigma_-}(\mathbf{a}_1) \times \cdots \times M_{\sigma_-}(\mathbf{a}_l)$  is bounded above by

$$\mathbf{a}^2 + 2 + 2n_0 + 2n_1 \leq \mathbf{v}^2 - 2n_0 - 2n_1 < \mathbf{v}^2,$$

a contradiction.

**Step 2.** We have  $(\mathbf{s}, \mathbf{v}) \leq 0$ .

Indeed, assume for a contradiction that  $(\mathbf{s}, \mathbf{v}) > 0$ . Then  $\mathbf{w}_1 = \rho_{\mathbf{s}}(\mathbf{w}_0)$  and we can write  $\mathbf{v} = a\mathbf{s} + b\mathbf{w}_0 + \mathbf{a}$ . We have, as before,

$$\begin{aligned} \mathbf{a}^2 &= (\mathbf{v} - a\mathbf{s} - b\mathbf{w}_0)^2 = \mathbf{v}^2 - 2a(\mathbf{v}, \mathbf{s}) - 2a^2 - 2b(\mathbf{v}, \mathbf{w}_0) + 2ab(\mathbf{s}, \mathbf{w}_0) \\ &= \mathbf{v}^2 - 2a(\mathbf{v}, \mathbf{s}) - 2a^2 - 2b((\mathbf{v}, \mathbf{w}_0) - (a\mathbf{s}, \mathbf{w}_0)) \\ &\leq \mathbf{v}^2 - 2a(\mathbf{v}, \mathbf{s}) - 2a^2 - 2b. \end{aligned}$$

Hence, the dimension of  $M_{\sigma_-}(\mathbf{a}_1) \times \cdots \times M_{\sigma_-}(\mathbf{a}_l)$  is bounded above by

$$\mathbf{a}^2 + 2 + 2b \leq \mathbf{v}^2 - 2a(\mathbf{v}, \mathbf{s}) - 2a^2 + 2 < \mathbf{v}^2,$$

a contradiction.

**Step 3.** We have  $(\mathbf{s}, \mathbf{v}) = 0$ .

Indeed, assume for a contradiction that  $(\mathbf{s}, \mathbf{v}) < 0$ . By Proposition 6.8,  $\mathcal{W}$  is a totally semistable wall for  $\mathbf{v}$ . We consider  $\mathbf{v}' = \rho_{\mathbf{s}}(\mathbf{v})$  as in Lemma 8.3. The wall  $\mathcal{W}$  induces a divisorial contraction for  $\mathbf{v}$  if and only if it induces one for  $\mathbf{v}'$ . But, since  $(\mathbf{v}, \mathbf{w}) \neq 1, 2$ , for all  $\mathbf{w}$  isotropic, then  $(\mathbf{v}', \mathbf{w}) \neq 1, 2$  as well. Moreover,  $(\mathbf{s}, \mathbf{v}') > 0$ . This is a contradiction, by Step 2.  $\square$

The converse of Proposition 8.6 is a consequence of the following three lemmata:

**Lemma 8.7.** *Let  $\mathcal{W}$  be a potential wall. Assume that  $(\mathbf{v}, \mathbf{w}_0) = 2$ . Then  $\mathcal{W}$  induces a divisorial contraction.*

*Proof.* By identifying  $M_{\sigma_+}(\mathbf{v})$  with  $M_{\omega, \beta}(-\mathbf{v})$ , this follows if there is a divisor of  $\omega$ -Gieseker stable sheaves which are torsion-free but not locally-free. We can describe explicitly a contracted divisor of  $\sigma_0$ -semistable objects as follows. Since  $(\mathbf{v}, \mathbf{w}_0) = 2$ , we can assume that  $\mathbf{v}$  has the form  $-(2, D, s)$ , with  $D$  an integral divisor which is either primitive or  $D = 0$ . We can consider the vector  $\mathbf{v}' = -(2, D, s+1)$ . By assumption,  $(\mathbf{v}')^2 \geq -2$ , and so  $M_{\omega, \beta}(-\mathbf{v}') \neq \emptyset$  by Theorem 2.13. Given any  $\omega$ -Gieseker stable sheaf  $F$  with vector  $\mathbf{v}'$  and a point  $x \in X$ , the surjections  $F \twoheadrightarrow k(x)$  induce extensions

$$k(x) \rightarrow E[1] \rightarrow F[1] \rightarrow k(x)[1]$$

of objects in  $M_{\sigma_+}(\mathbf{v})$  that are  $S$ -equivalent with respect to  $\sigma_0$ . Dimension counting shows that they sweep out a divisor.  $\square$

**Lemma 8.8.** *Let  $\mathcal{W}$  be a potential wall. Assume that there exists an effective spherical class  $\mathbf{s} \in \mathcal{H}$  such that  $(\mathbf{v}, \mathbf{s}) = 0$ . Then  $\mathcal{W}$  induces a divisorial contraction.*

*Proof.* Let  $S \in M_{\sigma_0}(\mathbf{s})$  be the unique  $\sigma_0$ -stable spherical object with Mukai vector  $\mathbf{s}$ . Consider the vector  $\mathbf{a} = \mathbf{v} - \mathbf{s}$ . Then we have

$$\begin{aligned} \mathbf{a}^2 &= (\mathbf{v} - \mathbf{s})^2 = \mathbf{v}^2 - 2 \\ (\mathbf{a}, \mathbf{s}) &= -\mathbf{s}^2 = 2. \end{aligned}$$

If  $\mathbf{v}^2 > 2$ , then  $\mathbf{a}^2 > 0$ . Since  $\mathbf{w}_1 = b\mathbf{s} + \mathbf{w}_0$ , with  $b > 0$ , we have  $(\mathbf{w}_1, \mathbf{a}) > (\mathbf{w}_0, \mathbf{a})$ . If  $(\mathbf{w}_0, \mathbf{a}) \geq 2$ , then  $\mathcal{W}$  is not a totally semistable wall for  $\mathbf{a}$ . Hence, given  $A \in M_{\sigma_0}(\mathbf{a})$ , all the extensions

$$S \rightarrow E \rightarrow A$$

give a divisor  $D \subset M_{\sigma_+}(\mathbf{v})$ , which is a  $\mathbb{P}^1$ -fibration over  $M_{\sigma_0}^{st}(\mathbf{a})$  and which gets contracted by crossing the wall  $\mathcal{W}$ . If  $(\mathbf{w}_0, \mathbf{a}) = 1$ , then also  $(\mathbf{w}_0, \mathbf{s}) = 1$ . Hence,  $-\mathbf{v}$  has rank 2. Then  $\mathcal{W}$  induces a divisorial contraction by Lemma 8.7.

Finally, assume that  $\mathbf{v}^2 = 2$ . Then  $\mathbf{a}$  is an isotropic vector with  $(\mathbf{a}, \mathbf{v}) = (\mathbf{a}, \mathbf{s}) = 2$ . But this implies that  $(\mathbf{w}_0, \mathbf{v}) = 1, 2$ . The case  $(\mathbf{w}_0, \mathbf{v}) = 2$  is again Lemma 8.7; and if  $(\mathbf{w}_0, \mathbf{v}) = 1$ , then  $-\mathbf{v}$  has rank 1, and we are in the case of the Hilbert-Chow wall.  $\square$

**Lemma 8.9.** *Let  $\mathcal{W}$  be a potential wall. If there exists an isotropic class  $\mathbf{w}$  such that  $(\mathbf{v}, \mathbf{w}) \in \{1, 2\}$ , then  $\mathcal{W}$  induces a divisorial contraction.*

*Proof.* By Lemma 8.1, the class  $\mathbf{w}$  is automatically effective. By Lemma 8.7, the only remaining case is  $\mathbf{w} = \mathbf{w}_1$ , with  $\mathbf{w}_1 = a\mathbf{s} + \mathbf{w}_0$  and  $a > 0$ . By Lemma 8.8, we can assume that  $(\mathbf{s}, \mathbf{v}) \neq 0$ .

If  $(\mathbf{s}, \mathbf{v}) > 0$ , then

$$(\mathbf{w}_1, \mathbf{v}) = a(\mathbf{s}, \mathbf{v}) + (\mathbf{w}_0, \mathbf{v}) \in \{1, 2\}.$$

Since  $(\mathbf{w}_0, \mathbf{v}) > 0$  and  $a > 0$ , this is possible only if  $(\mathbf{w}_0, \mathbf{v}) = 1$ , which corresponds to the Hilbert-Chow contraction.

Hence, we can assume  $(\mathbf{s}, \mathbf{v}) < 0$ . By Proposition 6.8,  $\mathcal{W}$  is a totally semistable wall for  $\mathbf{v}$ , and  $\mathcal{W}$  induces a divisorial contraction with respect to  $\mathbf{v}$  if and only if it induces one with respect to  $\mathbf{v}' = \rho_{\mathbf{s}}(\mathbf{v})$ . But then  $(\mathbf{v}', \mathbf{w}_0) = (\mathbf{v}, \mathbf{w}_1) \in \{1, 2\}$ . Again, we can use Lemma 8.7 to finish the proof.  $\square$

## 9. FLOPPING WALLS

This section deals with the remaining case of a potential wall  $\mathcal{W}$ : assuming that  $\mathcal{W}$  does not correspond to a divisorial contraction, we describe in which cases it is a flopping wall, a fake wall, or not a wall. This is the content of Propositions 9.1 and 9.4.

**Proposition 9.1.** *Assume that  $\mathcal{W}$  does not induce a divisorial contraction. If either*

- (a)  *$\mathbf{v}$  can be written as the sum  $\mathbf{v} = \mathbf{a}_1 + \mathbf{a}_2$  of two positive classes  $\mathbf{a}_1, \mathbf{a}_2 \in P_{\mathcal{H}} \cap \mathcal{H}$ , or*
- (b) *there exists a spherical class  $\tilde{\mathbf{s}} \in \mathcal{W}$  with  $0 < (\tilde{\mathbf{s}}, \mathbf{v}) \leq \frac{\mathbf{v}^2}{2}$ ,*

*then  $\mathcal{W}$  induces a small contraction.*

**Lemma 9.2.** *Let  $M$  be a lattice of rank two, and  $C \subset M \otimes \mathbb{R}^2$  be a convex cone not containing a line. If a primitive lattice element  $\mathbf{v} \in M \cap C$  can be written as the sum  $\mathbf{v} = \mathbf{a} + \mathbf{b}$  of two classes in  $\mathbf{a}, \mathbf{b} \in M \cap C$ , then it can be written as a sum  $\mathbf{v} = \mathbf{a}' + \mathbf{b}'$  of two classes  $\mathbf{a}', \mathbf{b}' \in m \cap C$  in such a way that the parallelogram with vertices  $0, \mathbf{a}', \mathbf{v}, \mathbf{b}'$  does not contain any other lattice point besides its vertices.*

*Proof.* If the parallelogram  $0, \mathbf{a}, \mathbf{v}, \mathbf{b}$  contains an additional lattice point  $\mathbf{a}'$ , we may replace  $\mathbf{a}$  by  $\mathbf{a}'$  and  $\mathbf{b}$  by  $\mathbf{v} - \mathbf{a}'$ . This procedure terminates.  $\square$

**Lemma 9.3.** *Let  $\mathbf{a}, \mathbf{b}, \mathbf{v} \in \mathcal{H} \cap C_{\mathcal{W}}$  be effective classes with  $\mathbf{v} = \mathbf{a} + \mathbf{b}$ . Assume that the following conditions are satisfied:*

- *The phases of  $\mathbf{a}, \mathbf{b}$  satisfy  $\phi^+(\mathbf{a}) < \phi^+(\mathbf{b})$ .*
- *The objects  $A, B$  are  $\sigma_+$ -stable with  $\mathbf{v}(A) = \mathbf{a}, \mathbf{v}(B) = \mathbf{b}$ .*
- *The parallelogram in  $\mathcal{H} \otimes \mathbb{R}$  with vertices  $0, \mathbf{a}, \mathbf{v}, \mathbf{b}$  does not contain any other lattice point.*
- *The extension  $A \hookrightarrow E \twoheadrightarrow B$  satisfies  $\text{Hom}(B, E) = 0$ .*

*Then  $E$  is  $\sigma_+$ -stable.*

*Proof.* Let  $\mathbf{a}_i$  be the Mukai vector of a Harder-Narashimhan filtration factor of  $E$ . By Proposition 5.1 part (c) and Remark 5.3, we have  $\mathbf{a}_i \in \mathcal{H}$ . We have  $E \in \mathcal{P}_+([\phi^+(\mathbf{a}), \phi^+(\mathbf{b})])$ , and hence  $\mathbf{a}_i$  is contained in the cone generated by  $\mathbf{a}, \mathbf{b}$ . Since the same holds for  $\mathbf{v} - \mathbf{a}_i = \sum_{j \neq i} \mathbf{a}_j$ ,  $\mathbf{a}_i$  is in fact contained in the parallelogram with vertices  $0, \mathbf{a}, \mathbf{v}, \mathbf{b}$ . Since it is also a lattice point, the assumption on the parallelogram implies  $\mathbf{a}_i \in \{\mathbf{a}, \mathbf{b}, \mathbf{v}\}$ .

Assume that  $E$  is not  $\sigma_+$ -stable, and let  $A_1 \subset E$  be the first HN filtration factor. Since  $\phi^+(\mathbf{a}_1) > \phi^+(\mathbf{v})$ , we must have  $\mathbf{a}_1 = \mathbf{b}$ . By the stability of  $A, B$  we have  $\text{Hom}(A_1, A) = 0$ , and  $\text{Hom}(A_1, B) = 0$  unless  $A_1 \cong B$ . Either of these is a contradiction.  $\square$

*Proof of Proposition 9.1.* We first consider case (a), so  $\mathbf{v} = \mathbf{a}_1 + \mathbf{a}_2$  with  $\mathbf{a}_1, \mathbf{a}_2 \in P_{\mathcal{H}}$ . Using Lemma 9.2, we may assume that the parallelogram with vertices  $0, \mathbf{a}_1, \mathbf{v}, \mathbf{a}_2$  does not contain an interior lattice point. In particular,  $\mathbf{a}_1, \mathbf{a}_2$  are primitive. We may also assume that  $\phi^+(\mathbf{a}_1) < \phi^+(\mathbf{a}_2)$ . By the signature of  $\mathcal{H}$  (see the proof of Lemma 6.4), we have  $(\mathbf{a}_1, \mathbf{a}_2) > 2$ . By Theorem 2.13, there exist  $\sigma_+$ -stable objects  $A_i$  of class  $\mathbf{v}(A_i) = \mathbf{a}_i$ . The inequality for the Mukai pairing implies  $\text{ext}^1(A_2, A_1) > 2$ . By Lemma 9.3, any extension

$$0 \rightarrow A_1 \hookrightarrow E \twoheadrightarrow A_2 \rightarrow 0$$

of  $A_2$  by  $A_1$  is  $\sigma_+$ -stable of class  $\mathbf{v}$ . As all these extensions are S-equivalent to each other with respect to  $\sigma_0$ , we obtain a projective space of dimension at least two that gets contracted by  $\pi^+$ .

Now consider case (b). First assume that  $\tilde{s}$  is an effective class. Note that  $(\mathbf{v} - \tilde{s})^2 \geq -2$ . Consider the parallelogram with vertices  $0, \tilde{s}, \mathbf{v}, \mathbf{v} - \tilde{s}$ . If it contains additional lattice point  $\mathbf{a}$ , then a simple computation shows  $\mathbf{a}^2 > -2$ , i.e.,  $\mathbf{a}^2 \geq 0$ ; thus  $\mathbf{v}$  can be written as the sum of positive classes, and the claim follows from the previous paragraph. Otherwise, observe that  $(\tilde{s}, \mathbf{v} - \tilde{s}) = (\tilde{s}, \mathbf{v}) + 2 > 2$ . If  $\tilde{S}$  is the  $\sigma_+$ -stable object of class  $\tilde{s}$ , and  $F$  any  $\sigma_+$ -stable object of class  $\mathbf{v} - \tilde{s}$ , then again  $\text{ext}^1(\tilde{S}, F) = \text{ext}^1(F, \tilde{S}) > 2$ . Thus, with the same arguments we obtain a family of  $\sigma_+$ -stable objects parametrized by a projective space that gets contracted by  $\pi^+$ .

We are left with the case where  $\tilde{s}$  is not effective. Set  $\tilde{\mathbf{t}} = -\tilde{s}$ , which is an effective class. With the same reasoning as above, we may assume that the parallelogram with vertices  $0, \tilde{\mathbf{t}}, \mathbf{v}, \mathbf{v} - \tilde{\mathbf{t}}$  contains no additional lattice points. Set  $\mathbf{v}' = \rho_{\tilde{\mathbf{t}}}(\mathbf{v}) - \tilde{\mathbf{t}} = \mathbf{v} - ((\tilde{s}, \mathbf{v}) + 1)\tilde{\mathbf{t}}$ . We have  $\mathbf{v}'^2 \geq -2$  and  $(\tilde{\mathbf{t}}, \mathbf{v}') = (\tilde{\mathbf{t}}, \mathbf{v}) + 2 > 2$ . The lattice points in the parallelogram with vertices  $0, ((\tilde{\mathbf{t}}, \mathbf{v}) + 1)\tilde{\mathbf{t}}, \mathbf{v}, \mathbf{v}'$  are given by  $k\tilde{\mathbf{v}}$  and  $\mathbf{v}' + k\tilde{\mathbf{v}}$  for  $k \in \mathbb{Z}, 0 \leq k \leq (\tilde{\mathbf{t}}, \mathbf{v}) + 1$  (otherwise, already the parallelogram with vertices  $0, \tilde{\mathbf{t}}, \mathbf{v}, \mathbf{v} - \tilde{\mathbf{t}}$  would contain additional lattice points).

Let  $\tilde{T}$  and  $F$  be  $\sigma_+$ -stable objects of class  $\tilde{\mathbf{t}}$  and  $\mathbf{v}'$ , respectively. Let us assume  $\phi^+(\tilde{\mathbf{t}}) > \phi^+(\mathbf{v})$ , the other case being analogous. Any subspace  $\mathfrak{V} \subset \text{Ext}^1(\tilde{T}, F)$  of dimension  $(\tilde{\mathbf{t}}, \mathbf{v}) + 1$  defines an extension

$$0 \rightarrow F \hookrightarrow E \twoheadrightarrow \tilde{T} \otimes \mathfrak{V} \rightarrow 0$$

such that  $E$  is of class  $\mathbf{v}(E) = \mathbf{v}$ , and satisfies  $\text{Hom}(\tilde{T}, E) = 0$ . If  $E$  were not  $\sigma_+$ -stable, then the class of the maximal destabilizing subobject  $A$  would have to be a lattice point in the parallelogram with vertices  $0, ((\tilde{\mathbf{t}}, \mathbf{v}) + 1)\tilde{\mathbf{t}}, \mathbf{v}, \mathbf{v}'$ ; therefore,  $\mathbf{v}(A) = k\tilde{\mathbf{t}}$ . The only  $\sigma_+$ -semistable object of this class is  $\tilde{T}^{\oplus k}$ , and we get a contradiction. Thus, we have constructed a family of  $\sigma_+$ -stable objects of class  $\mathbf{v}$  parametrized by the Grassmannian  $\text{Gr}((\tilde{\mathbf{t}}, \mathbf{v}) + 1, \text{ext}^1(\tilde{T}, F))$  that become S-equivalent with respect to  $\sigma_0$ .  $\square$

It remains to prove the converse of Proposition 9.1:

**Proposition 9.4.** *Assume that  $\mathcal{W}$  does not induce a divisorial contraction. Assume that  $\mathbf{v}$  can't be written as the sum of two positive classes in  $P_{\mathcal{H}}$ , and that there is no spherical class  $\mathbf{s} \in \mathcal{H}$  with  $0 < (\mathbf{s}, \mathbf{v}) \leq \frac{\mathbf{v}_0^2}{2}$ . Then  $\mathcal{W}$  is either a fake wall, or not a wall.*

*Proof.* First consider the case where  $\mathbf{v} = \mathbf{v}_0$  is the minimal class in its orbit  $G_{\mathcal{H}} \cdot \mathbf{v}$ . We will prove that every  $\sigma_+$ -stable object  $E$  of class  $\mathbf{v}_0$  is also  $\sigma_0$ -stable. Assume otherwise, that  $E$  is  $\sigma_-$ -unstable. Let  $\mathbf{a}_1, \dots, \mathbf{a}_l$  be the Mukai vectors of the HN filtration factors of  $E$  with respect to  $\sigma_-$ . If all classes  $\mathbf{a}_i$  are positive,  $\mathbf{a}_i \in P_{\mathcal{H}}$ , then we have an immediate contradiction to the assumptions.

Otherwise,  $E$  must have a spherical destabilizing subobject, or a spherical destabilizing quotient. Let  $\tilde{\mathbf{s}}$  be the class of this spherical object. If there is only one  $\sigma_0$ -stable spherical object, then it is easy to see that  $\mathbf{v}_0 - \tilde{\mathbf{s}}$  is in the positive cone; therefore,  $(\tilde{\mathbf{s}}, \mathbf{v}_0) < \frac{\mathbf{v}_0^2}{2}$  in contradiction to our assumption.

If there are two  $\sigma_0$ -stable spherical objects of classes  $\mathbf{s}, \mathbf{t}$ , consider the two vectors  $\mathbf{v}_0 - \mathbf{s}$  and  $\mathbf{v}_0 - \mathbf{t}$ . The assumptions imply  $(\mathbf{v}_0 - \mathbf{s})^2 < -2$  and  $(\mathbf{v}_0 - \mathbf{t})^2 < -2$ ; on the other hand,  $\mathbf{v} - \tilde{\mathbf{s}}$  is effective; using Lemma 6.2, this implies that  $\mathbf{v}_0 - \mathbf{s}$  or  $\mathbf{v}_0 - \mathbf{t}$  must be effective. We claim that this leads to a simple numerical contradiction. Indeed,  $(\mathbf{v}_0 - \mathbf{t})^2 < -2$  constrains  $\mathbf{v}_0$  to lie below a concave down hyperbola, and  $(\mathbf{v}_0 - \mathbf{s})^2 < -2$  to lie above a concave up hyperbola; the two hyperbolas intersect at the points  $0$  and  $\mathbf{s} + \mathbf{t}$ . Therefore, if we write






FIGURE 5. The three hyperbolas in the proof of Proposition 9.4

$\mathbf{v}_0 = xs + yt$ , we have  $x, y < 1$ . Thus, neither  $\mathbf{v}_0 - s$  nor  $\mathbf{v}_0 - t$  can be effective (see Figure 5).

In the case where  $\mathbf{v}$  is not minimal,  $\mathbf{v} \neq \mathbf{v}_0$ , let  $\Phi$  be the sequence of spherical twists given by Proposition 6.8. Since the assumptions of our proposition are invariant under the  $G_{\mathcal{H}}$ -action, they are also satisfied by  $\mathbf{v}_0$ . By the previous case, we know that every  $\sigma_+$ -stable objects  $E_0$  of class  $\mathbf{v}_0$  is also  $\sigma_0$ -stable. Thus  $\Phi$  induces a morphism  $\Phi_*: M_{\sigma_+}(\mathbf{v}_0) \rightarrow M_{\sigma_+}(\mathbf{v})$ ; since  $\Phi_*$  is injective and the two spaces are smooth projective varieties of the same dimension, it is an isomorphism. The S-equivalence class of  $\Phi(E_0)$  is determined by that of  $E_0$ ; since S-equivalence is a trivial equivalence relation on  $M_{\sigma_+}(\mathbf{v}_0)$ , the same holds for  $M_{\sigma_+}(\mathbf{v})$ , and thus  $\pi^+$  is an isomorphism.  $\square$

Proposition 9.4 finishes the proof of Theorem 5.7.

## 10. MAIN THEOREMS

We will first complete the proof of Theorem 1.1.

*Proof of Theorem 1.1, part (b).* We consider a wall  $\mathcal{W}$  with nearby stability conditions  $\sigma_{\pm}$ , and  $\sigma_0 \in \mathcal{W}$ . Since  $M_{\sigma_{\pm}}$  are  $K$ -trivial varieties, it is sufficient to find an open subset  $U \subset M_{\sigma_{\pm}}(\mathbf{v})$  with complement of codimension two, and an (anti-)autoequivalence  $\Phi_{\mathcal{W}}$  of  $D^b(X, \alpha)$ , such that  $\Phi_{\mathcal{W}}(E)$  is  $\sigma_-$ -stable for all  $E \in U$ .

We will distinguish cases according to Theorem 5.7. First consider the case when  $\mathcal{W}$  corresponds to a flopping contraction, or when  $\mathcal{W}$  is a fake wall. If  $\mathcal{W}$  does not admit an effective spherical class  $s \in \mathcal{H}_{\mathcal{W}}$  with  $(s, \mathbf{v}) < 0$  then we can choose  $U$  to be the open subset of  $\sigma_0$ -stable objects; its complement has codimension two, and there is nothing to prove. Otherwise, there exists a spherical object destabilizing every object in  $M_{\sigma_+}(\mathbf{v})$ . Let  $\mathbf{v}_0 \in \mathcal{H}_{\mathcal{W}}$  be the minimal class of the  $G_{\mathcal{H}}$ -orbit of  $\mathbf{v}$ , in the sense of Definition 6.6. The subset  $U$  of  $\sigma_0$ -stable objects in  $M_{\sigma_0}(\mathbf{v}_0)$  has complement of codimension two. Then the sequence of spherical twists of Proposition 6.8, applied for  $\sigma_+$  and  $\sigma_-$ , identifies  $U$  with subsets of  $M_{\sigma_+}(\mathbf{v})$  and

$M_{\sigma_-}(\mathbf{v})$  via derived equivalences  $\Phi^+, \Phi^-$ ; then the composition  $\Phi^- \circ (\Phi^+)^{-1}$  has the desired property.

Next assume that  $\mathcal{W}$  induces a divisorial contraction. We have three cases to consider:

**Brill-Noether:** Again, we first assume that there is no effective spherical class  $\mathbf{s}$  with  $(\mathbf{s}, \mathbf{v}) < 0$ . The contracted divisor is described in Proposition 7.1, and the Harder-Narasimhan filtration of the destabilized objects in Lemma 7.4. We may assume that we are in the case where the Brill-Noether divisor in  $M_{\sigma_+}(\mathbf{v})$  is described by  $\text{Hom}(\tilde{S}, \_) \neq 0$ . Now consider the spherical twist  $\text{ST}_{\tilde{S}}$  at  $\tilde{S}$ , applied to objects  $E \in M_{\sigma_+}(\mathbf{v})$ . Note that by  $\sigma_+$ -stability, we have  $\text{Ext}^2(\tilde{S}, E) = \text{Hom}(E, \tilde{S})^\vee = 0$  for any such  $E$ ; since  $(\mathbf{v}(\tilde{S}), \mathbf{v}(E)) = 0$ , it follows that  $\text{hom}(\tilde{S}, E) = \text{ext}^1(\tilde{S}, E)$ .

If  $E$  does not lie on the Brill-Noether divisor, then  $\mathbf{R}\text{Hom}(\tilde{S}, E) = 0$ , and so  $\text{ST}_{\tilde{S}}(E) = E$ . Also, for generic such  $E$  (away from a codimension two subset), the object  $E$  is also  $\sigma_-$ -stable.

If  $E$  is a generic element of the Brill-Noether divisor, then  $\text{Hom}(\tilde{S}, E) \cong \mathbb{C} \cong \text{Ext}^1(\tilde{S}, E)$ , and hence we have an exact triangle

$$\tilde{S} \oplus \tilde{S}[-1] \rightarrow E \rightarrow \text{ST}_{\tilde{S}}(E).$$

Its long exact cohomology sequence with respect to the t-structure of  $\sigma_0$  induces two short exact sequences

$$\tilde{S} \hookrightarrow E \twoheadrightarrow F \quad \text{and} \quad F \hookrightarrow \text{ST}_{\tilde{S}}(E) \twoheadrightarrow \tilde{S}.$$

By Lemma 7.5, the former is the HN filtration of  $E$  with respect to  $\sigma_-$ ; the latter is the dual extension, which is a  $\sigma_-$ -stable object by [BM12, Lemma 5.9].

Thus, in both cases,  $\text{ST}_{\tilde{S}}(E)$  is  $\sigma_-$ -stable, proving the claim.

If instead there is an effective spherical class  $\mathbf{s}$  with  $(\mathbf{s}, \mathbf{v}) < 0$ , we reduce to the previous case, similarly to the situation of flopping contractions: Let  $\mathbf{v}_0$  again denote the minimal class in the orbit  $G_{\mathcal{H}} \cdot \mathbf{v}$ ; note that  $\mathcal{W}$  also induces a divisorial contraction of Brill-Noether type for  $\mathbf{v}_0$ . In this case, Lemma 7.5 states that the sequence  $\Phi$  of spherical twists identifies an open subset  $U^+ \subset M_{\sigma_+}(\mathbf{v}_0)$  (with complement of codimension two) with an open subset of  $M_{\sigma_+}(\mathbf{v})$ ; similarly for  $U^- \subset M_{\sigma_-}(\mathbf{v}_0)$ . Combined with the single spherical twist identifying a common open subset of  $M_{\sigma_{\pm}}(\mathbf{v}_0)$ , this implies the claim.

**Hilbert-Chow:** As shown in Section 8, we may assume that shift by one identifies  $M_{\sigma_+}(\mathbf{v})$  with the Gieseker-moduli space  $M_{\omega}(-\mathbf{v})$  of stable sheaves of rank one on a twisted K3 surface  $(Y, \alpha')$ . After tensoring with a line bundle, we may assume that objects in  $M_{\sigma_+}(\mathbf{v})$  are exactly the shifts  $I_Z[1]$  of ideal sheaves of 0-dimensional subschemes  $Z \subset Y$ .

In the setting of Proposition 8.2, we have  $\beta = 0$ . Since there are line bundles on  $(Y, \alpha')$ , the Brauer group element  $\alpha'$  is trivial. By the last statement of the same Proposition, the moduli space  $M_{\sigma_-}(\mathbf{v})$  parametrizes the shifts of derived duals ideal sheaf. Thus there is a natural isomorphism  $M_{\sigma_-}(\mathbf{v}) \cong M_{\sigma_+}(\mathbf{v})$  induced by the derived anti-autoequivalence  $(\_)^\vee[2]$ .

**Li-Gieseker-Uhlenbeck:** We will argue along similar lines as in the previous case; unfortunately, the details are more involved. The first difference is that we can't assume  $\beta = 0$ . Instead, first observe that  $M_{\sigma_+}(\mathbf{v}) = M_{\omega, \beta}(-\mathbf{v})$  is parametrizing  $\beta$ -twisted Gieseker-stable sheaves  $F$  of rank  $2 = (\mathbf{v}, \mathbf{w})$ , and of slope  $\mu_{\omega}(F) = \omega \cdot \beta$ . If we assume  $\omega$  to be generic, then Gieseker-stability is independent of the choice of  $\beta$ ; we can consider  $M_{\sigma_+}(\mathbf{v}) = M_{\omega}(-\mathbf{v})$  to be the moduli space of shifts  $F[1]$  of  $\omega$ -Gieseker-stable sheaves  $F$ .

Since  $(Y, \alpha')$  admits rank two vector bundles, the order of  $\alpha'$  in the Brauer group is one or two; in both cases, we can identify  $(Y, \alpha')$  with  $(Y, -\alpha')$ , and thus the derived dual  $E \mapsto E^\vee$  defines an anti-autoequivalence of  $D^b(Y, \alpha')$ .

Write  $-\mathbf{v}$  as  $-\mathbf{v} = (2, c, d)$ , and let  $\mathcal{L}$  be the unique line bundle with  $c_1(L) = c$ . From the previous discussion it follows that  $\Phi(\_) = (\_)^\vee \otimes \mathcal{L}[2]$  is the desired functor:

Indeed, any object in  $M_{\sigma_+}(\mathbf{v})$  is of the form  $F[1]$  for a  $\omega$ -Gieseker stable sheaf  $F$  of class  $\mathbf{v}$ . Then  $\Phi(F[1]) = F^\vee \otimes \mathcal{L}[1]$  the derived dual of a Gieseker-stable sheaf, and has class  $\mathbf{v}$ . By Proposition 8.2, this is an object of  $M_{\sigma_-}(\mathbf{v})$ .  $\square$

Consider two adjacent chamber  $\mathcal{C}^+, \mathcal{C}^-$  separated by a wall  $\mathcal{W}$ ; as always, we pick stability conditions  $\sigma_\pm \in \mathcal{C}^\pm$ , and a stability condition  $\sigma_0 \in \mathcal{W}$ . By the identification of Néron-Severi groups induced by Theorem 1.1, we can think of the corresponding maps  $\ell_\pm$  of equation (5) as maps

$$\ell_\pm: \mathcal{C}^\pm \rightarrow \text{NS}(M_{\sigma_\pm}(\mathbf{v})).$$

They can be written as the following composition of maps

$$\text{Stab}(X, \alpha) \xrightarrow{\mathcal{Z}} H_{\text{alg}}^*(X, \alpha, \mathbb{Z}) \otimes \mathbb{C} \xrightarrow{I} \mathbf{v}^\perp \xrightarrow{\theta_{\mathcal{C}^\pm}} \text{NS}(M_{\sigma_\pm}(\mathbf{v}))$$

where  $\mathcal{Z}$  is the map defined in Theorem 2.8,  $I$  is given by  $I(\Omega_Z) = \Im \frac{\Omega_Z}{-(\Omega_Z, \mathbf{v})}$ , and where  $\theta_{\mathcal{C}^\pm}$  are the Mukai morphisms, as reviewed in Remark 2.14.

Our next goal is to show that these two maps behave as nicely as one could hope; we will distinguish two cases according to the behaviour of the contraction morphism

$$\pi^+: M_{\sigma_+}(\mathbf{v}) \rightarrow \overline{M}^+$$

induced by  $\mathcal{W}$  via Theorem 2.16:

**Lemma 10.1.** *The maps  $\ell^+, \ell^-$  agree on the wall  $\mathcal{W}$  (when extended by continuity).*

- (a) (Fake or flopping walls) When  $\pi^+$  is an isomorphism, or a small contraction, then the maps  $\ell_+, \ell_-$  are analytic continuations of each other.
- (b) (Bouncing walls) When  $\pi^+$  is a divisorial contraction, then the analytic continuations of  $\ell^+, \ell^-$  differ by the reflection  $\rho_D$  in  $\text{NS}(M_{\sigma_+}(\mathbf{v}))$  at the divisor  $D$  contracted by  $\ell_{\sigma_0}$ .

Here “reflection at  $D$ ” denotes the linear involution leaving  $D^\perp$  fixed, and sending  $D$  to  $-D$ . Markman proved in [Mar09] that such a reflection is an integral linear transformation, for any irreducible exceptional divisor on a hyperkähler variety; of course in our situation, this statement can easily be deduced from the classification of divisorial contractions.

As a consequence, walls of the former type (a) are fake walls when  $\pi^+$  is an isomorphism, or induce a flop when  $\pi^+$  is a divisorial contraction; for walls of the latter type (b), corresponding to a divisorial contraction, the moduli spaces  $M_{\sigma_+}(\mathbf{v}), M_{\sigma_-}(\mathbf{v})$  for the two adjacent chambers are isomorphic.

*Proof.* We have to prove  $\theta_{\mathcal{C}^-} = \theta_{\mathcal{C}^+}$  in case (a), and  $\theta_{\mathcal{C}^-} = \rho_D \circ \theta_{\mathcal{C}^+}$  in case (b). We will always assume for simplicity that the two moduli spaces admit universal families; the arguments apply identically to quasi-universal families.

Consider case (a). If the wall is not totally semistable, then the two moduli spaces  $M_{\mathcal{C}^\pm}(\mathbf{v})$  share a common open subset, with complement of codimension two, on which the two universal families agree. By the projectivity of the moduli spaces, the maps  $\theta_{\mathcal{C}^\pm}$  are determined

by their restriction to curves contained in this subset; this proves the claim. If the wall is instead totally semistable, we additionally have to use Proposition 6.8. Let  $\Phi^+$  and  $\Phi^-$  be the two sequences of spherical twists, sending  $\sigma_0$ -stable objects of class  $\mathbf{v}_0$  to  $\sigma_+$ - and  $\sigma_-$ -stable objects of class  $\mathbf{v}$ , respectively. The autoequivalence inducing the birational map  $M_{\sigma_+}(\mathbf{v}) \dashrightarrow M_{\sigma_-}(\mathbf{v})$  is given by  $\Phi^- \circ (\Phi^+)^{-1}$ . As the classes of the spherical objects occurring in  $\Phi^+$  and  $\Phi^-$  are identical, this does not change the class of the universal family in the  $K$ -group; therefore, the Mukai morphisms  $\theta_{\mathcal{C}^+}, \theta_{\mathcal{C}^-}$  agree.

Now consider the case of a Brill-Noether divisorial contraction; we first assume that there is no effective spherical class  $\mathbf{s}' \in \mathcal{H}_{\mathcal{W}}$  with  $(\mathbf{s}', \mathbf{v}) < 0$ . The contraction induced by a spherical object  $S$  with Mukai vector  $\mathbf{s} := \mathbf{v}(S) \in \mathbf{v}^\perp$ . The class of the contracted divisor is  $\theta(\mathbf{s})$ . The universal families differ (up to a subset of codimension two) by the spherical twist  $\mathrm{ST}_S(\_)$ . This induces the reflection at  $\mathbf{s}$  in  $H_{\mathrm{alg}}^*(X, \alpha, \mathbb{Z})$ ; thus the Mukai morphisms differ by reflection at  $\theta(\mathbf{s})$ , as claimed.

If in addition to  $\mathbf{s} \in \mathbf{v}^\perp$ , there does exist an effective spherical class  $\mathbf{s}' \in \mathcal{H}_{\mathcal{W}}$  with  $(\mathbf{s}', \mathbf{v}) < 0$ , we have to rely on the constructions of Lemma 7.5, as in the proof of Theorem 1.1. We have a common open subset  $U \subset M_{\sigma_\pm}(\mathbf{v}_0)$ , such that the two universal families  $\mathcal{E}^\pm|_U$  are related by the spherical twist at a spherical object  $S_0$  of class  $\mathbf{s}_0$ . Let  $\Phi^\pm$  be the sequences of spherical twists obtained from Lemma 7.5, applied to  $\sigma_+$  or  $\sigma_-$ , respectively. Their induced maps  $\Phi^\pm: H_{\mathrm{alg}}^*(X, \alpha, \mathbb{Z}) \rightarrow H_{\mathrm{alg}}^*(X, \alpha, \mathbb{Z})$  on the Mukai lattice are identical, as they are obtained by twists of spherical objects of the same classes; it sends  $\mathbf{v}_0$  to  $\mathbf{v}$ , and thus  $\mathbf{s}_0$  to  $\pm \mathbf{s}$ . Therefore, the composition  $\Phi^- \circ \mathrm{ST}_{S_0} \circ (\Phi^+)^{-1}$  induces the reflection at  $\mathbf{s}$ , as claimed.

It remains to consider divisorial contractions of Hilbert-Chow and Li-Gieseker-Uhlenbeck type. We may assume  $M_{\sigma_+}(v)$  is the Hilbert scheme, or a moduli space of Giesker-stable sheaves of rank two.

By the proof of Theorem 1.1, there is a line bundle  $\mathcal{L}$  on  $X$  such that  $\mathbf{R}\mathcal{H}om_{M_{\sigma_\pm} \times X}(\mathcal{E}, (p_X)^* \mathcal{L}[2])$  is a universal family with respect to  $\sigma_-$  on  $M_{\sigma_-}(v) = M_{\sigma_+}(v)$ . We will compare  $\theta_{\mathcal{C}^\pm}$  by evaluating its degree on a test curve  $C \subset M_{\sigma_\pm}$ . Let  $i$  denote the inclusion  $i: C \times X \hookrightarrow M_{\sigma_\pm} \times X$ , and by  $p$  the projection  $p: C \times X \rightarrow X$ . This yields the following chain of equalities for  $\mathbf{a} \in \mathbf{v}^\perp$ :

$$(18) \quad \theta_{\mathcal{C}^-}(\mathbf{a}).C = \left( \mathbf{a}, \mathbf{v}(p_* i^*(\mathcal{E}^-)) \right) = \left( \mathbf{a}, \mathbf{v}(p_* \mathbf{R}\mathcal{H}om_{C \times X}(i^* \mathcal{E}, \mathcal{O}_C \boxtimes \mathcal{L}[2])) \right)$$

$$(19) \quad = \left( \mathbf{a}, \mathbf{v}(p_* \mathbf{R}\mathcal{H}om_{C \times X}(i^* \mathcal{E}, \omega_C[2] \boxtimes \mathcal{L})) \right)$$

$$(20) \quad = \left( \mathbf{a}, \mathbf{v}(\mathbf{R}\mathcal{H}om_X(p_* i^* \mathcal{E}, \mathcal{L})) \right)$$

$$(21) \quad = \left( \mathbf{a}^\vee \cdot \mathrm{ch}(\mathcal{L}), \mathbf{v}(p_* i^* \mathcal{E}) \right) = \theta_{\mathcal{C}^+}(\mathbf{a}^\vee \cdot \mathrm{ch}(\mathcal{L})).C$$

Here we used compatibility of duality with base change in (18),  $\mathbf{a} \in \mathbf{v}^\perp$  in (19), and Grothendieck duality in (20). In (21), we wrote  $\mathbf{a}^\vee$  for the class corresponding to  $\mathbf{a}$  under duality  $(\_)^\vee$ , i.e., the class that agrees with  $\mathbf{a}$  except for the sign of the component in the divisor part  $\mathrm{NS}(X)$ .

In the Hilbert-Chow case, with  $\mathbf{v} = -(1, 0, 1 - n)$ , the class of the contracted divisor  $D$  is proportional to  $(1, 0, n - 1)$ , and we have  $\mathcal{L} \cong \mathcal{O}_X$ ; in the Li-Gieseker-Uhlenbeck case, we can write  $\mathbf{v} = (2, c, d)$ , the class of the contracted divisor as a multiple of  $(2, c, \frac{c^2}{2} - d)$ , and  $c_1(\mathcal{L}) = c$ . In both cases, a direct verification shows that the reflection  $\rho_D$  is compatible with the above chain of equalities:  $\rho_D(\mathbf{a}) = \mathbf{a}^\vee \cdot \mathrm{ch}(\mathcal{L})$ .  $\square$

*Proof of Theorem 1.2, (a), (b), (c).* Lemma 10.1 proves part (a). Part (c) follows directly from the positivity  $\ell_{\mathcal{C}}(\mathcal{C}) \subset \text{Amp}M_{\mathcal{C}}(\mathbf{v})$  once we have established part (b).

Consider a big class in the movable cone, given as  $\theta_{\sigma}(\mathbf{a})$  for some class  $\mathbf{a} \in \mathbf{v}^{\perp}$ ,  $\mathbf{a}^2 > 0$ ; we have to show that it is in the image of  $\ell$ . Recall the definition of  $\mathcal{P}_0^+(X, \alpha)$  given in the discussion preceding Theorem 2.8. If we set  $\Omega' = i\mathbf{a} - \frac{\mathbf{v}}{\mathbf{v}^2} \in H_{\text{alg}}^*(X, \mathbb{Z}) \otimes \mathbb{C}$ , then clearly  $\Omega' \in \mathcal{P}(X, \alpha)$ . In case there is a spherical class  $\mathbf{s} \in H_{\text{alg}}^*(X, \alpha, \mathbb{Z})$  with  $(\Omega', \mathbf{s}) = 0$ , we modify  $\Omega'_{\mathbf{a}}$  by a small real multiple of  $\mathbf{s}$  to obtain  $\Omega \in \mathcal{P}_0(X, \alpha)$ , otherwise we set  $\Omega_{\mathbf{a}} = \Omega'_{\mathbf{a}}$ ; in either case, we have  $\Omega_{\mathbf{a}} \in \mathcal{P}_0(X, \alpha)$  with  $(\Omega_{\mathbf{a}}, \mathbf{v}) = -1$  and  $\Im \Omega_{\mathbf{a}} = \mathbf{a}$ . In addition, the fact that  $\theta(\mathbf{a})$  is contained in the positive cone gives  $\Omega \in \mathcal{P}_0^+(X, \alpha)$ .

Let  $\Omega_{\sigma} \in \mathcal{P}_0^+(X, \alpha)$  be the central charge for the chosen basepoint  $\sigma \in \text{Stab}(X, \alpha)$ . Then there is a path  $\gamma: [0, 1] \rightarrow \mathcal{P}_0^+(X, \alpha)$  starting at  $\Omega_{\sigma}$  and ending at  $\Omega_{\mathbf{a}}$  with the following additional property: for all  $t \in [0, 1]$ , the class  $-\frac{\theta_{\sigma}(\Im \gamma(t))}{(\gamma(t), \mathbf{v})}$  is contained in the movable cone of  $M_{\sigma}(\mathbf{v})$ .

By Theorem 2.8, there is a lift  $\sigma: [0, 1] \rightarrow \text{Stab}^{\dagger}(X, \alpha)$  of  $\gamma$  starting at  $\sigma(0) = \sigma$ . By the above assumption on  $\gamma$ , this will never hit a wall of the movable cone corresponding to a divisorial contraction; by Lemma 10.1, the map  $\ell$  extends analytically, with  $\theta_{\sigma} = \theta_{\sigma(0)} = \theta_{\sigma(1)}$ . Therefore,

$$\ell_{\sigma(1)}(\sigma(1)) = \theta_{\sigma(1)}(\mathbf{a}) = \theta_{\sigma}(\mathbf{a})$$

as claimed.  $\square$

In fact, following [Mar11, Section 6], one can use a Weyl group action on the positive cone to give a global description of the map  $\ell$ . As in [Mar11, Definition 6.8], we denote by  $W_{\text{Exc}} \subset \text{Aut}(\text{NS}(M_{\sigma}(\mathbf{v})))$  the hyperbolic reflection group generated by the reflections  $\rho_D$  at exceptional divisors  $D$  of divisorial contractions.

We consider its action on the cone  $\text{Pos}(M_{\sigma}(\mathbf{v}))$  of strictly positive divisors. The *exceptional chamber* of this Weyl group action is defined by  $(D, \_ ) \geq 0$  for all exceptional divisors  $D$ . As explained by Markman, the general theory of hyperbolic reflection groups shows that this is a fundamental domain for  $W_{\text{Exc}}$ . On the other hand, it coincides with the image of  $\ell$ ; by Theorem 1.2, the exceptional chamber is equal to the intersection of the movable cone with the big cone. This recovers [Mar11, Lemma 6.22].

The exceptional chamber of a hyperbolic reflection group intersects every  $W_{\text{Exc}}$ -orbit exactly once. Thus there is a map

$$W: \text{Pos}(M_{\sigma}(\mathbf{v})) \rightarrow \overline{\text{Mov}}(M_{\sigma}(\mathbf{v}))$$

sending any class to the intersection of its  $W_{\text{Exc}}$ -orbit with the fundamental domain. Then Lemma 10.1 and Theorem 1.2 immediately give the following:

**Theorem 10.2.** *The map  $\ell$  of Theorem 1.2 can be given as the composition of the following maps:*

$$\text{Stab}^{\dagger}(X, \alpha) \xrightarrow{\mathcal{Z}} H_{\text{alg}}^*(X, \alpha, \mathbb{Z}) \otimes \mathbb{C} \xrightarrow{I} \mathbf{v}^{\perp} \xrightarrow{\theta_{\sigma}} \text{Pos}(M_{\sigma}(\mathbf{v})) \xrightarrow{W} \overline{\text{Mov}}(M_{\sigma}(\mathbf{v})).$$

To finish the proof of Theorem 1.2, (d), we have the following observation:

**Proposition 10.3.** *Let  $\mathcal{C} \subset \text{Stab}^{\dagger}(X, \alpha)$  be a chamber of the chamber decomposition with respect to  $\mathbf{v}$ . Then the image of  $\ell_{\mathcal{C}}(\mathcal{C}) \subset \text{NS}(M_{\mathcal{C}}(\mathbf{v}))$  of the chamber  $\mathcal{C}$  is exactly the ample cone of the corresponding moduli space  $M_{\mathcal{C}}(\mathbf{v})$ .*

*Proof.* In light of Theorems 2.15 and 1.2, (a), (b), (c), the only potential problem is given by walls  $\mathcal{W} \subset \partial\mathcal{C}$  that do not get mapped to walls of the nef cone of the moduli space. These are totally semistable fake walls induced by an effective spherical class  $\mathbf{s} \in \mathcal{H}_{\mathcal{W}}$  with  $(\mathbf{s}, \mathbf{v}) < 0$ . The idea is that there is always a potential wall  $\mathcal{W}'$ , with the same lattice  $\mathcal{H}_{\mathcal{W}'} = \mathcal{H}_{\mathcal{W}}$ , for which all effective spherical classes have positive pairing with  $\mathbf{v}$ . By Theorem 5.7,  $\mathcal{W}'$  is not a wall, and it will have the same image in the nef cone of  $M_{\mathcal{C}}(\mathbf{v})$  as the wall  $\mathcal{W}$ .

Let  $\sigma_0 = (Z_0, \mathcal{P}_0) \in \mathcal{W}$  be a very general stability condition on the given wall: this means we can assume that  $\mathcal{H}_{\mathcal{W}}$  contains all integral classes  $\mathbf{a} \in H_{\text{alg}}^*(X, \alpha, \mathbb{Z})$  with  $\Im Z_0(\mathbf{a}) = 0$ . If we write  $Z_0(\_) = (\Omega_0, \_)$  as in Theorem 2.8, we may assume that  $\Omega_0$  is normalized by  $(\Omega_0, \mathbf{v}) = -1$  and  $\Omega_0^2 = 0$ , i.e.,  $(\Re\Omega_0, \Im\Omega_0) = 0$  and  $(\Re\Omega_0)^2 = (\Im\Omega_0)^2$  (see [Bri08, Section 10]). We will now replace  $\sigma_0$  by a stability condition whose central charge has real part give by  $(-\mathbf{v}, \_)$ , and identical imaginary part.

To this end, let  $\sigma_1 \in \mathcal{C}$  be a stability condition nearby  $\sigma_0$ , whose central charge is defined by  $\Omega_1 = \Omega_0 + i\epsilon$ , where  $\epsilon \in H_{\text{alg}}^*(X, \alpha, \mathbb{Z}) \otimes \mathbb{R}$  is a sufficiently small vector with  $(\epsilon, \mathbf{v}) = 0$ ; we may also assume that multiples of  $\mathbf{v}$  are the only integral classes  $\mathbf{a} \in H_{\text{alg}}^*(X, \alpha, \mathbb{Z})$  with  $(\Im\Omega_1, \tilde{\mathbf{a}}) = 0$ . Let  $\Omega_2 = -\mathbf{v} + i\Im\Omega_1$ ; then a straight-forward computation shows that the straight path connection  $\Omega_1$  with  $\Omega_2$  lies completely within  $\mathcal{P}_0^+(X, \alpha)$ . Finally, let  $\Omega_3 = -\mathbf{v} + \Im\Omega_0$ ; by Theorem 5.7, there are no spherical classes  $\tilde{\mathbf{s}} \in \mathcal{H}_{\mathcal{W}}$  with  $(\mathbf{v}, \mathbf{s}) = 0$ , implying that the straight path from  $\Omega_2$  to  $\Omega_3$  is also contained in  $\mathcal{P}_0^+(X, \alpha)$ .

By Theorem 2.8, there is a lift of the path  $\Omega_0 \mapsto \Omega_1 \mapsto \Omega_2 \mapsto \Omega_3$  to  $\text{Stab}(X, \alpha)$ ; let  $\sigma_2$  and  $\sigma_3$  the stability conditions corresponding to  $\Omega_2$  and  $\Omega_3$ , respectively. By choice of  $\epsilon$ , we may assume that the paths  $\sigma_0 \mapsto \sigma_1$  and  $\sigma_2 \mapsto \sigma_3$  do not cross any walls. Since  $(\Omega_1, \mathbf{v}) = (\Omega_2, \mathbf{v}) = -1$ , and since the imaginary part on the path  $\Omega_1 \mapsto \Omega_2$  is constant, the same holds for the path  $\sigma_1 \mapsto \sigma_2$ . Hence  $\sigma_3$  is in the closure of the chamber  $\mathcal{C}$ . In particular,  $\sigma_3$  lies on a potential wall of  $\mathcal{C}$  with hyperbolic lattice given by  $\mathcal{H}_{\mathcal{W}}$ ; by construction, any spherical class  $\mathbf{s} \in \mathcal{H}_{\mathcal{W}}$  with  $(\mathbf{v}, \mathbf{s}) < 0$  satisfies  $(\Omega_3, \mathbf{s}) > 0$ , and thus  $\mathbf{s}$  is not effective.

By Theorem 5.7,  $\sigma_3$  does not lie on a wall. Since  $\Im\Omega_3 = \Im\Omega_0$ , the images  $l_{\mathcal{C}}(\sigma_0) = l_{\mathcal{C}}(\sigma_3)$  in the Néron-Severi group of  $M_{\mathcal{C}}(\mathbf{v})$  agree.  $\square$

We conclude this section by proving Corollary 1.3.

*Proof of Corollary 1.3.* We first prove the two implications.

“ $\Leftarrow$ ”: Assume that  $\Psi: D^b(X) \xrightarrow{\sim} D^b(X')$  is a derived (anti-)equivalence such that  $\Psi_*(\mathbf{v}) = \mathbf{v}'$ . Its associated Fourier-Mukai kernel induces a Hodge isometry  $\Psi_*: H^*(X, \mathbb{Z}) \rightarrow H^*(X', \mathbb{Z})$ , sending  $\mathbf{v}$  to  $\mathbf{v}'$ . By Verbitsky’s Torelli Theorem in the  $\text{Hilb}^n(\text{K3})$ -deformation type, the birational class of the moduli space is determined by the embedding  $\mathbf{v}^\perp \hookrightarrow H^*(X)$ , see [Mar11, Corollary 9.5 and Example 9.6].

“ $\Rightarrow$ ”: Now assume that  $M_H(\mathbf{v})$  is birational to  $M_{H'}(\mathbf{v}')$ . Applying the Torelli Theorem again, we obtain a Hodge isometry  $\psi: H^*(X, \mathbb{Z}) \xrightarrow{\sim} H^*(X', \mathbb{Z})$ . Since  $\mathbf{v}$  and  $\mathbf{v}'$  are primitive, this gives  $\psi(\mathbf{v}) = \pm\mathbf{v}'$ . Up to composing with the derived dual functor and the shift functor, we can assume that  $\psi(\mathbf{v}) = \mathbf{v}'$  and  $\psi$  is *orientation-preserving*. By Mukai-Orlov’s Derived Torelli Theorem (see [Orl97, HS06, HMS09]) for K3 surfaces, there exists a derived equivalence  $\Psi: D^b(X) \xrightarrow{\sim} D^b(X')$  such that  $\Psi_* = \psi$ .

To prove the final claim, choose stability conditions  $\sigma \in \text{Stab}(X)$ ,  $\sigma' \in \text{Stab}(X')$  such that  $M_\sigma(\mathbf{v}) = M_H(\mathbf{v})$  and  $M_{\sigma'}(\mathbf{v}') = M_{H'}(\mathbf{v}')$ . By the construction of the derived equivalence  $\Psi$ ,



it is easy to see that  $\Psi_*\sigma$  is in the same connected component as  $\sigma'$ . Thus we can apply Theorem 1.1 to obtain an auto-equivalence  $\Phi$  of  $D^b(X')$  that sends a generic element of  $M_{\Psi_*\sigma}(\mathbf{v}')$  to an element of  $M_{\sigma'}(\mathbf{v}')$ . Then the composition  $\Phi \circ \Psi$  has the desired property.  $\square$

## 11. APPLICATION 1: LAGRANGIAN FIBRATIONS

In this section, we will explain how birationality of wall-crossing implies Theorem 1.5, verifying the Lagrangian fibration conjecture.

One implication in Theorem 1.5 is immediate: if  $f: M_H(\mathbf{v}) \dashrightarrow Z$  is a rational abelian fibration, then the pull-back  $f^*D$  of any ample divisor  $D$  on  $Z$  has volume zero; by equation (6), the self-intersection of  $f^*D$  with respect to the Beauville-Bogomolov form must also equal zero.

We will prove the converse for any moduli space  $M_\sigma(\mathbf{v})$  of Bridgeland-stable objects on a (possibly twisted) K3 surface  $(X, \alpha)$ , under the assumptions that  $\mathbf{v}$  is primitive, and  $\sigma$  generic with respect to  $\mathbf{v}$ . We will first restate carefully the argument establishing part (a) of Conjecture 1.4, which was already sketched in the introduction; then we will explain how to extend the argument to also obtain part (b).

Assume that there is an integral divisor  $D$  on  $M_\sigma(\mathbf{v})$  with  $q(D) = 0$ . Applying the inverse of the Mukai morphism  $\theta_{\mathbf{v}}$  of Theorem 3.5, we obtain a primitive vector  $\mathbf{w} = \theta_{\mathbf{v}}^{-1}(D) \in \mathbf{v}^\perp$  with  $\mathbf{w}^2 = 0$ .

After a small deformation, we may assume that  $\sigma$  is also generic with respect to  $\mathbf{w}$ . As in Section 8, we consider the moduli space  $Y := M_\sigma(\mathbf{w})$  of  $\sigma$ -stable objects, which is a smooth K3 surface. There is a derived equivalence

$$(22) \quad \Phi: D^b(X, \alpha) \xrightarrow{\sim} D^b(Y, \alpha')$$

for the appropriate choice of a Brauer class  $\alpha' \in \text{Br}(Y)$ ; as before, we have  $\Phi_*(\mathbf{w}) = (0, 0, 1)$ .

By definition,  $\Phi$  induces an isomorphism

$$(23) \quad M_\sigma(\mathbf{v}) \cong M_{\Phi_*(\sigma)}(\Phi(\mathbf{v})),$$

where  $\Phi(\sigma)$  is generic with respect to  $\Phi(\mathbf{v})$ .

**Lemma 11.1.** *The Mukai vector  $\Phi_*(\mathbf{v})$  has rank zero.*

*Proof.* This follows directly from  $\Phi_*(\mathbf{w}) = (0, 0, 1)$  and  $(\Phi_*(\mathbf{w}), \Phi_*(\mathbf{v})) = (\mathbf{w}, \mathbf{v}) = 0$ .  $\square$

We write  $\Phi(\mathbf{v}) = (0, C, s)$ , with  $C \in \text{Pic}(Y)$  and  $s \in \mathbb{Z}$ . Since  $\mathbf{v}^2 > 0$  we have  $C^2 > 0$ .

**Lemma 11.2.** *After replacing  $\Phi$  by the composition  $\Psi \circ \Phi$ , where  $\Psi \in \text{Aut}(D^b(Y, \alpha'))$ , we may assume that  $C$  is ample, and that  $s \neq 0$ .*

*Proof.* Up to shift [1], we may assume that  $H' \cdot C > 0$ , for a given ample class  $H'$  on  $Y$ . In particular,  $C$  is an effective class; it is ample unless there is a rational  $-2$ -curve  $D \subset Y$  with  $C \cdot D < 0$ . Applying the spherical twist  $\text{ST}_{\mathcal{O}_D}$  at the structure sheaf<sup>4</sup> of  $D$  replaces  $C$  its image  $C'$  under the reflection at  $D$ , which satisfies  $C' \cdot D > 0$ . This procedure terminates; indeed, the nef cone is the fundamental exceptional chamber of the Weyl group action generated by reflections at  $-2$ -curves.

Since tensoring with an (untwisted) line bundle on  $Y$  induces an autoequivalence of  $D^b(Y, \alpha')$ , we may also assume  $s \neq 0$ .  $\square$

<sup>4</sup>Note that the restriction of  $\alpha$  to any curve vanishes, hence the structure sheaf  $\mathcal{O}_D$  is a coherent sheaf on  $(Y, \alpha')$ .

Let  $H' \in \text{Amp}(Y)$  be a generic polarization with respect to  $\mathbf{w}$ .

**Lemma 11.3.** *The moduli space  $M_{H'}(\Phi_*(\mathbf{v}))$  is non-empty and admits a structure of Lagrangian fibration.*

Of course, this is just a small (and well-known) generalization of Beauville's integrable system [Bea91].

*Proof.* The non-emptiness follows directly from [Yos01, Theorem 8.1]. The structure of Lagrangian fibration is given as follows:

Consider the line bundle  $L'$  on  $M' := M_{H'}(\Phi(\mathbf{v}))$  given by  $\theta_{\mathbf{v}}((0, 0, -1))$ . By an argument of Faltings and Le Potier (see [LP05, Section 1.3]), we can construct sections of  $L'$  as follows: for all  $y \in Y$ , we define a section  $s_y \in H^0(M', L')$  by its zero-locus

$$Z(s_y) := \{E \in M' : \text{Hom}(E, k(y)) \neq 0\}.$$

Whenever  $y$  is not in the support of  $E$ , then  $s_y$  does not vanish at  $E$ ; hence the sections  $\{s_y\}_{y \in Y}$  generate  $L'$ . The image of  $E$  is determined by its set-theoretic support; hence the image of the map induced by  $L'$  is the complete local system of  $C$ ; by Matsushita's theorem [Mat99, Mat01], the map must be a Lagrangian fibration.  $\square$

**Lemma 11.4.** *The stability condition  $\Phi_*\sigma$  on  $D^b(Y, \alpha')$  is contained in the connected component  $\text{Stab}^\dagger(Y, \alpha')$  constructed in [Bri08, HMS08].*

*Proof.* By construction of the equivalence in (22), skyscraper sheaves of points are  $\Phi_*$ -stable; then the statement follows from [Bri08, Proposition 10.3]. In Lemma 11.2, we had to modify  $\Phi$  by a composition of spherical twists at structure sheaves of rational curves; by [Bri08, Section 12], this will not cause us to leave the connected component.  $\square$

By Remark 2.12, there exists a generic stability condition  $\sigma' \in \text{Stab}^\dagger(Y, \alpha')$  with the property that  $M_{H'}(\Phi(\mathbf{v})) = M_{\sigma'}(\Phi(\mathbf{v}))$ . On the other hand, by the birationality of wall-crossing, Theorem 1.1, the moduli spaces  $M_{\sigma'}(\Phi_*(\mathbf{v}))$  and  $M_{\Phi_*(\sigma)}(\Phi_*(\mathbf{v}))$  are birational; combined with the identification (23), this shows that  $M_\sigma(\mathbf{v})$  is birational to a Lagrangian fibration.

It remains to prove part (b), so let us assume that  $D$  is nef. Using the Fourier-Mukai transform  $\Phi$  as above, and after replacing  $\sigma$  by  $\Phi_*\sigma$ , we may also assume that  $\mathbf{v}$  has rank zero, and that  $\mathbf{w} = \theta_\sigma^{-1}(D)$  is the class of skyscraper sheaf of points. Now consider the autoequivalence  $\Psi \in \text{Aut } D^b(Y, \alpha')$  of Lemma 11.2. Except for the possible shift [1], each autoequivalence used in the construction of  $\Psi$  leaves the class  $\mathbf{w}$  invariant. Thus, in the moduli space  $M_{\Psi_*\sigma}(\Psi_*\mathbf{v}) = M_\sigma(\mathbf{v})$ , the divisor class  $D$  is still given by  $D = \pm \theta_{\Psi_*\sigma}(\mathbf{w})$ , up to sign.

Let  $f : M_\sigma(\mathbf{v}) \dashrightarrow M_H(\mathbf{v})$  be the birational map to the Gieseker-moduli space  $M_H(\mathbf{v})$  of torsion sheaves induced by a sequence of wall-crossings as above. The Lagrangian fibration  $M_H(\mathbf{v}) \rightarrow \mathbb{P}^n$  is induced by the divisor  $\theta_H(-\mathbf{w})$ . By Theorem 10.2, the classes  $f_*D$  and  $\theta_H(-\mathbf{w})$  are (up to sign) in the same  $W_{\text{Exc}}$ -orbit. Since they are both nef on a smooth K-trivial birational model, they are in the closure of the movable cone (and in particular, their orbits agree, not just up to sign).

Now recall from the discussion preceding Theorem 10.2 that the exceptional chamber for the action of  $W_{\text{Exc}}$  on the positive cone is a fundamental domain, which intersects every  $W_{\text{Exc}}$ -orbit exactly once. The same holds for the closure of the exceptional chamber and the action on the closure of the positive cone. Therefore, the classes  $f_*D$  and  $\theta_H(-\mathbf{w})$  have to be equal.

Since  $M_\sigma(\mathbf{v})$  and  $M_H(\mathbf{v})$  are isomorphic in codimension two, the section rings of  $D$  and  $f_*D$  agree. In particular,  $D$  is effective. The conclusion now follows, for example, by [Kaw85, Fuj11]. This completes the proof of Theorem 1.5.

**Remark 11.5.** In fact, the above proof shows the following two additional statements:

- (a) If  $D \in \text{NS}(M_\sigma(\mathbf{v}))$  with  $q(D) = 0$  lies in the closure of the movable cone, then there is a birational Lagrangian fibration induced by  $D$ . (In particular,  $D$  is movable.)
- (b) Any  $W_{\text{Exc}}$ -orbit of divisors on  $M_{\sigma(\mathbf{v})}$  satisfying  $q(D) = 0$  contains exactly one movable divisor, which induces a birational Lagrangian fibration.

## 12. APPLICATION 2: MORI CONE, NEF CONE, MOVABLE CONE, EFFECTIVE CONE

Let  $\mathbf{v}$  be a primitive vector with  $\mathbf{v}^2 > 0$ , let  $\sigma$  be a generic stability condition with respect to  $\mathbf{v}$ , and let  $M := M_\sigma(\mathbf{v})$  be the moduli space of  $\sigma$ -semistable objects. In this section, we will completely describe the cones associated to the birational geometry of  $M$  in terms of the Mukai lattice of  $X$ .

Recall that  $\overline{\text{Pos}}(M) \subset \text{NS}(M)_\mathbb{R}$  denotes the (closed) cone of positive classes defined by the Beauville-Bogomolov quadratic form. Let  $\overline{\text{Pos}}(M)_\mathbb{Q} \subset \overline{\text{Pos}}(M)$  be the subcone generated by all rational classes in  $\overline{\text{Pos}}(M)$ ; it is the union of the interior  $\text{Pos}(M)$  with all rational rays in the boundary  $\partial \text{Pos}(M)$ . We fix an ample divisor class  $A$  on  $M$  (which can be obtained from Theorem 2.15).

In the following theorems, we will say that a subcone of  $\overline{\text{Pos}}(M)_\mathbb{Q}$  (or of its closure) is “cut out” by a collection of linear subspaces if it is one of the closed chambers of the wall-and-chamber decomposition of  $\text{Pos}(M)_\mathbb{Q}$  whose walls are the given collection of subspaces. This is easily translated into a more explicit statement as in the formulation of Theorem 12.1 given in the introduction.

**Theorem 12.1.** *The nef cone of  $M$  is cut out in  $\overline{\text{Pos}}$  by all linear subspaces of the form  $\theta(\mathbf{v}^\perp \cap \mathbf{a}^\perp)$ , for all classes  $\mathbf{a} \in H_{\text{alg}}^*(X, \alpha, \mathbb{Z})$  satisfying  $\mathbf{a}^2 \geq -2$  and  $0 \leq (\mathbf{v}, \mathbf{a}) \leq \frac{\mathbf{v}^2}{2}$ .*

Via the Beauville-Bogomolov form we can identify the group  $N_1(M)$  of curves up to numerical equivalences with a lattice in the Néron-Severi group:  $N_1(M)_\mathbb{Q} \cong (N^1(M)_\mathbb{Q})^\vee \cong N^1(M)_\mathbb{Q}$ . In particular, we get an induced rational pairing on  $N_1(M)$ ; we then say that the *cone of positive curves* is the cone of classes  $[C] \in N_1(M)_\mathbb{R}$  with  $(C, C) > 0$  and  $C.A > 0$ . Also, we obtain a dual Mukai isomorphism

$$(24) \quad \theta^\vee: H_{\text{alg}}^*(X, \alpha, \mathbb{Z})/\mathbf{v} \otimes \mathbb{Q} \rightarrow N_1(M)_\mathbb{Q}.$$

As the dual statement to Theorem 12.1, we obtain:

**Theorem 12.2.** *The Mori cone of curves in  $M$  is generated by the cone of positive curves, and by all curve classes  $\theta^\vee(\mathbf{a})$ , for all  $\mathbf{a} \in H_{\text{alg}}^*(X, \alpha, \mathbb{Z})$ ,  $\mathbf{a}^2 \geq -2$  satisfying  $0 \leq (\mathbf{v}, \mathbf{a}) \leq \frac{\mathbf{v}^2}{2}$  and  $\theta^\vee(\mathbf{a}).A > 0$ .*

Some of these classes  $\mathbf{a}$  may not define a wall bordering the nef cone; in this case,  $\theta^\vee(\mathbf{a})$  is in the interior of the Mori cone (as it intersects every nef divisor positively).

**Theorem 12.3.** *The movable cone of  $M$  is cut out in  $\overline{\text{Pos}}(M)_\mathbb{Q}$  by the following two types of walls:*

- (a)  $\theta(\mathbf{s}^\perp \cap \mathbf{v}^\perp)$  for every spherical class  $\mathbf{s} \in \mathbf{v}^\perp$ .
- (b)  $\theta(\mathbf{w}^\perp \cap \mathbf{v}^\perp)$  for every isotropic class  $\mathbf{w} \in H_{\text{alg}}^*(X, \alpha, \mathbb{Z})$  with  $1 \leq (\mathbf{w}, \mathbf{v}) \leq 2$ .

**Theorem 12.4.** *The effective cone of  $M$  is generated by  $\overline{\text{Pos}}(M)_\mathbb{Q}$  along with the following exceptional divisors:*

- (a)  $D := \theta(\mathbf{s})$  for every spherical class  $\mathbf{s} \in \mathbf{v}^\perp$  with  $(D, A) > 0$ , and

- (b)  $D := \theta(\mathbf{v}^2 \cdot \mathbf{w} - (\mathbf{v}, \mathbf{w}) \cdot \mathbf{v})$  for every isotropic class  $\mathbf{w} \in H_{\text{alg}}^*(X, \alpha, \mathbb{Z})$  with  $1 \leq (\mathbf{w}, \mathbf{v}) \leq 2$  and  $(D, A) > 0$ .

Note that only those classes  $D$  whose orthogonal complement  $D^\perp$  is a wall of the movable cone will correspond to irreducible exceptional divisors.

The movable cone has essentially been described by Markman for any hyperkähler variety; more precisely, [Mar11, Lemma 6.22] gives the intersection of the movable cone with the strictly positive cone  $\text{Pos}(M)$ . While our methods give an alternative proof, the only new statement of Theorem 12.3 concerns rational classes  $D$  with  $D^2 = 0$  in the closure of the movable cone; such a  $D$  is movable due to our proof of the Lagrangian fibration conjecture in Theorem 1.5.

Using the divisorial Zariski decomposition of [Bou04], one can show for any hyperkähler variety that the pseudo-effective cone is dual to the closure of the movable cone. In particular, Theorem 12.4 could also be deduced from Markman's results and Theorem 1.5.

*Proof of Theorem 12.1.* Let  $\mathcal{C}$  be the chamber of  $\text{Stab}(X, \alpha)$  containing  $\sigma$ . By Theorem 1.2, the boundary of the ample cone inside the positive cone is equal to the union of the images  $\ell(\mathcal{W})$ , for all walls  $\mathcal{W}$  in the boundary of  $\mathcal{C}$  that induce a non-trivial contraction morphism. (These are walls that are not “fake walls” in the sense of Definition 2.17.) Theorem 5.7 characterizes hyperbolic lattices corresponding to such walls.

For any such hyperbolic lattice  $\mathcal{H}$ , we get a class  $\mathbf{a}$  as in Theorem 12.1 as follows:

- in the cases (b) of divisorial contractions, we let  $\mathbf{a}$  be the corresponding spherical of isotropic class;
- in the subcase of (b) of a flopping contraction induced by a spherical class  $\mathbf{s}$ , we also set  $\mathbf{a} = \mathbf{s}$ ;
- and in the subcase of (b) of a flopping contraction induced by a sum  $\mathbf{v} = \mathbf{a} + \mathbf{b}$ , we may assume  $(\mathbf{v}, \mathbf{a}) \leq (\mathbf{v}, \mathbf{b})$ , which is equivalent to  $(\mathbf{v}, \mathbf{a}) \leq \frac{\mathbf{v}^2}{2}$ .

Stability conditions  $\sigma = (Z, \mathcal{A})$  in the corresponding wall  $\mathcal{W}$  satisfy  $\Im \frac{Z(\mathbf{a})}{Z(\mathbf{v})} = 0$ , or, equivalently,  $\ell(\sigma) \in \theta(\mathbf{v}^\perp \cap \mathbf{a}^\perp)$ .

Conversely, given  $\mathbf{a}$ , we obtain a rank two lattice  $\mathcal{H} := \langle \mathbf{v}, \mathbf{a} \rangle$ . If  $\mathcal{H}$  is hyperbolic, then it is straightforward to check that it conversely induces one of the walls listed in Theorem 5.7. Otherwise,  $\mathcal{H}$  is positive-semidefinite. Then the orthogonal complement  $H^\perp = \mathbf{v}^\perp \cap \mathbf{a}^\perp$  does not contain any positive classes, and thus its image under  $\theta$  in  $\text{NS}(M)$  does not intersect the positive cone and can be ignored.  $\square$

*Proof of Theorem 12.3.* As already discussed in Section 10, the intersection  $\text{Mov}(M) \cap \text{Pos}(M)$  follows directly from Theorem 1.2; the statement of Theorem 12.3 is just an explicit description of the exceptional chamber of the Weyl group action.

A movable class  $D$  in the boundary of the positive cone, with  $(D, D) = 0$ , automatically has to be rational. Conversely, by our proof of Theorem 1.5, if we have a rational divisor with  $(D, D) = 0$  that is in the closure of the movable cone, then there is a Lagrangian fibration induced by  $D$  on a smooth birational model of  $M$ ; in particular,  $D$  is movable.  $\square$

*Proof of Theorem 12.4.* As indicated above, the pseudo-effective cone is dual to the movable cone; thus we just need to verify that the Theorem gives the correct description of the boundary.

For exceptional divisors, with  $(D, D) < 0$ , this follows from our classification and construction of divisorial contractions. For classes with  $(D, D) = 0$ , this again follows from Theorem 1.5.  $\square$

**Relation to Hassett-Tschinkel's conjecture on the Mori cone.** Hassett and Tschinkel gave a conjectural description of the nef and Mori cones via intersection numbers of extremal rays in [HT10]. While their conjecture turned out to be incorrect (see [BM12, Remark 9.4] and [CK12, Remark 8.10]), we will now explain that it is in fact very closely related to Theorem 12.2.

We first recall their conjecture. Via the identification  $N_1(M)_\mathbb{Q} \cong N^1(M)_\mathbb{Q}$  explained above, the Beauville-Bogomolov extends to a quadratic form on  $N_1(M)$  with values in  $\mathbb{Q}$ ; we will also denote it by  $q(\_)$ . The following lemma follows immediately from this definition, and the definition of  $\theta^\vee$ :

**Lemma 12.5.** *Consider the isomorphism  $\mathbf{v}_\mathbb{Q}^\perp \cong N_1(M)_\mathbb{Q}$  induced by the dual Mukai morphism  $\theta^\vee$  of (24). This isomorphism respects the quadratic form on either side.*

Let  $2n$  be the dimension of  $M$ , and as above let  $A$  be an ample divisor. Let  $\mathcal{C} \subset N_1(M)_\mathbb{R}$  be the cone generated by all integral curve classes  $R \in N_1(M)_\mathbb{Z}$  that satisfy  $q(R) \geq -\frac{n+3}{2}$  and  $R.A \geq 0$ . In [HT10, Conjecture 1.2], the authors conjectured that for of any hyperkähler variety  $M$  deformation equivalent to the Hilbert scheme of a K3 surface, the cone  $\mathcal{C}$  is equal to the Mori cone.

Our first observation shows that the Mori cone is contained in  $\mathcal{C}$ :

**Proposition 12.6.** *Let  $R$  be the generator of an extremal ray of the Mori cone of  $M$ . Then  $(R, R) \geq -\frac{n+3}{2}$ .*

*Proof.* Note that it is enough to prove the inequality for some effective curve on the extremal ray. Let  $\mathcal{W}$  be a wall inducing the extremal contraction corresponding to the ray generated by  $R$ , and  $\mathcal{H}_\mathcal{W} \subset H_{\text{alg}}^*(X, \mathbb{Z})$  its associated hyperbolic lattice. Let  $\sigma_+$  be a nearby stability condition in the chamber of  $\sigma$ , and  $\sigma_0 \in \mathcal{W}$ . Let  $\mathbf{a} \in \mathcal{H}_\mathcal{W}$  be a corresponding class satisfying the assumptions in Theorem 12.2:  $\mathbf{a}^2 \geq -2$  and  $0 \leq (\mathbf{v}, \mathbf{a}) \leq \frac{\mathbf{v}^2}{2}$ .

We first claim that there exists a contracted curve whose integral class is given by  $\theta^\vee(\mathbf{a})$ . For simplicity we assume that  $\mathcal{W}$  is not a totally semistable wall for any class in  $\mathcal{H}_\mathcal{W}$ ; the general case can be reduced to this one with the same methods as in the previous sections. By assumptions, we have both  $\mathbf{a}^2 \geq -2$  and  $(\mathbf{v} - \mathbf{a})^2 \geq -2$ ; therefore, we can choose  $\sigma_0$ -stable objects  $A$  and  $B$  of class  $\mathbf{a}$  and  $\mathbf{v} - \mathbf{a}$ , respectively. Since  $\mathcal{H}_\mathcal{W}$  is hyperbolic, the assumptions also imply  $(\mathbf{a}, \mathbf{v}) < \mathbf{a}^2$ . Therefore,  $\text{ext}^1(B, A) = \text{ext}^1(A, B) = (\mathbf{a}, \mathbf{v} - \mathbf{a}) > 0$ ; in fact, in all the cases of Theorem 5.7 we have  $(\mathbf{a}, \mathbf{v} - \mathbf{a}) \geq 2$ .

Assume that  $\phi^+(\mathbf{a}) < \phi^+(\mathbf{v}) < \phi^+(\mathbf{v} - \mathbf{a})$ ; the opposite case follows similarly. Varying the extension class in  $\text{Ext}^1(B, A)$  produces curves of objects in  $M_{\sigma_+}(\mathbf{v})$  that are S-equivalent with respect to  $\sigma_0$ ; in order to compute its class, we have to make the construction explicit. Let  $\mathbb{P}(\text{Ext}^1(B, A))$  be the projective space of one-dimensional subspaces of  $\text{Ext}^1(B, A)$ . Choose a parametrized line  $\mathbb{P}^1 \hookrightarrow \mathbb{P}(\text{Ext}^1(B, A))$ , corresponding to a section  $\nu$  of

$$H^0(\mathbb{P}^1, \mathcal{O}(1) \otimes \text{Ext}^1(B, A)) = \text{Ext}_{\mathbb{P}^1 \times X}^1(\mathcal{O}_{\mathbb{P}^1} \boxtimes B, \mathcal{O}_{\mathbb{P}^1}(1) \boxtimes A).$$

Let  $\mathcal{E} \in D^b(\mathbb{P}^1 \times X)$  be the extension  $\mathcal{O}_{\mathbb{P}^1} \boxtimes B \rightarrow \mathcal{E} \rightarrow \mathcal{O}_{\mathbb{P}^1}(1) \boxtimes A$  given by  $\nu$ . By Lemma 6.9, every fiber of  $\mathcal{E}$  is  $\sigma_+$ -stable. Thus we have produced a rational curve  $R \subset M_{\sigma_+}(\mathbf{v})$  of S-equivalent objects.

To compute its class, it is sufficient to compute the intersection product  $\theta(D).R$  with a divisor  $\theta(D)$ , for any  $D \in \mathbf{v}^\perp$ . We have

$$\theta(D).R = (D, \mathbf{v}(\Phi(\mathcal{O}_R))) = (D, \mathbf{v}(B) + 2\mathbf{v}(A)) = (D, \mathbf{v} + \mathbf{a}) = (D, \mathbf{a}) = \theta(D).\theta^\vee(\mathbf{a}),$$

where  $\Phi: D^b(M_{\sigma^+}(\mathbf{v})) \rightarrow D^b(X)$  denotes the Fourier-Mukai transform, and where we used  $D \in \mathbf{v}^\perp$  in the second-to-last equality.

Let  $\mathbf{a}_0 \in H_{\text{alg}}^*(X, \mathbb{Z})$  denote the projection of  $\mathbf{a}$  to the orthogonal complement of  $\mathbf{v}$ . By Lemma 12.5, we have  $(R, R) = \mathbf{a}_0^2$ , and for the latter we obtain:

$$(\mathbf{a}_0, \mathbf{a}_0) = \left( \mathbf{a} - \frac{(\mathbf{v}, \mathbf{a})}{\mathbf{v}^2} \mathbf{v}, \mathbf{a} - \frac{(\mathbf{v}, \mathbf{a})}{\mathbf{v}^2} \mathbf{v} \right) = \mathbf{a}^2 - \frac{(\mathbf{v}, \mathbf{a})^2}{\mathbf{v}^2} \geq -2 - \frac{\mathbf{v}^2}{4} = -\frac{n+3}{2}.$$

□

**Remark 12.7.** When  $M$  is the Hilbert scheme of points on  $X$ , we can make the comparison to Hassett-Tschinkel's conjecture even more precise: in this case, it is easy to see that  $\theta^\vee$  induces an isomorphism

$$H_{\text{alg}}^*(X, \mathbb{Z})(X, \mathbb{Z})/\mathbf{v} \rightarrow N_1(M)$$

of lattices, respecting the integral structures. Given a class  $R \in N_1(M)$  satisfying the inequality  $(R, R) \geq -\frac{n+3}{2}$  of [HT10], let  $\mathbf{a}_0 \in \mathbf{v}_\mathbb{Q}^\perp$  be the (rational) class with  $\theta^\vee(\mathbf{a}_0) = R$ . Let  $k$  be any integer satisfying  $k \leq n-1$  and  $k^2 \geq (2n-2)(-2 - \mathbf{a}_0^2)$ ; by the assumptions,  $k = n-1$  is always an example satisfying both inequalities. Then  $\mathbf{a} := \mathbf{a}_0 + \frac{k}{2n-2} \mathbf{v}$  is a rational class in the algebraic Mukai lattice that satisfies the assumptions appearing in Theorem 12.2. In addition, it has integral pairing with both  $\mathbf{v}$ , and with every integral class in  $\mathbf{v}^\perp$ ; thus, it is potentially an integral class. The Hassett-Tschinkel conjecture holds if and only if for every extremal ray of  $\mathcal{C}$ , there is a choice of  $k$  such that  $\mathbf{a}$  is an integral class.

If we are given the lattice  $\mathbf{v}^\perp$ , then the algebraic Mukai lattice of  $X$  can be any lattice in  $\mathbf{v}_\mathbb{Q}^\perp \oplus \mathbb{Q} \cdot \mathbf{v}$  containing both  $\mathbf{v}^\perp$  and  $\mathbf{v}$ , as long as  $\mathbf{v}$  is primitive. In general, the Hassett-Tschinkel conjecture will hold for some of these lattices, but not for others. The question is thus closely related to the fact that a strong global Torelli statement needs the embedding  $H^2(M) \hookrightarrow H^*(X)$ , rather than just  $H^2(M)$ .

### 13. EXAMPLES OF NEF CONES AND MOVABLE CONES

In this section we examine examples of cones of divisors.

**K3 surfaces with Picard number 1...** Let  $X$  be a K3 surface such that  $\text{Pic}(X) \cong \mathbb{Z} \cdot H$ , with  $H^2 = 2d$ . We let  $M := \text{Hilb}^n(X)$ , for  $n \geq 2$ , and  $\mathbf{v} = (1, 0, 1-n)$ . In this case, everything is determined by certain Pell's equations. We recall that a basis of  $\text{NS}(M)$  is given by  $\tilde{H} = \theta(0, -H, 0)$ , the big and nef divisor given by the symmetric power of  $H$ , and  $B = \theta(-1, 0, 1-n)$ ; the exceptional divisor of the Hilbert-Chow morphism has class  $2B$ .

By Theorem 5.7, divisorial contractions can be divided in three cases:

**Brill-Noether:** If there exists a spherical class  $\mathbf{s}$  with  $(\mathbf{s}, \mathbf{v}) = 0$ .

**Hilbert-Chow:** If there exists an isotropic class  $\mathbf{w}$  with  $(\mathbf{w}, \mathbf{v}) = 1$ .

**Li-Gieseker-Uhlenbeck:** If there exists an isotropic class  $\mathbf{w}$  with  $(\mathbf{w}, \mathbf{v}) = 2$ .

The case of BN-contraction depends on the following Pell's equation

$$(25) \quad (n-1)X^2 - dY^2 = 1,$$

where  $\mathbf{s} = (r, cH, (n-1)r)$ ,  $X = r$  and  $Y = c$ .

The case of HC-contractions is governed by the Pell's equation

$$(26) \quad X^2 - d(n-1)Y^2 = 1,$$

where  $\mathbf{w} = (r, cH, (n-1)r-1)$ ,  $X = 2(n-1)r-1$  and  $Y = 2c$ .



Finally, for the case of LGU-contractions, the Pell's equation is again (26), but now  $\mathbf{w} = (r, cH, (n-1)r-2)$ ,  $X = (n-1)r-1$  and  $Y = c$ . This already gives the structure of the movable cone:

**Proposition 13.1.** *Assume  $\text{Pic}(X) \cong \mathbb{Z} \cdot H$ . The movable cone of the Hilbert scheme  $M = \text{Hilb}^n(X)$  has the following form:*

(a) *If  $d = \frac{k^2}{h^2}(n-1)$ , with  $k, h \geq 1$ ,  $(k, h) = 1$ , then*

$$\text{Mov}(M) = \langle \tilde{H}, h\tilde{H} - kB \rangle,$$

*where  $q(h\tilde{H} - kB) = 0$ , and it induces a (rational) Lagrangian fibration on  $M$ .*

(b) *If  $d(n-1)$  is not a perfect square, and (25) has a solution, then*

$$\text{Mov}(M) = \langle \tilde{H}, \tilde{H} - d\frac{y_1}{x_1(n-1)}B \rangle,$$

*where  $(x_1, y_1)$  is the non-trivial solution to (25) with smallest possible  $x_1 > 0$  and  $y_1 > 0$ .*

(c) *If  $d(n-1)$  is not a perfect square, and (25) has no solution, then*

$$\text{Mov}(M) = \langle \tilde{H}, \tilde{H} - d\frac{y'_1}{x'_1}B \rangle,$$

*where  $(x'_1, y'_1)$  is the non-trivial solution to (26) with smallest possible  $x'_1 > 0$  and  $y'_1 > 0$ .*

*Proof.* Part (a) follows directly from Theorem 1.5. To prove part (b) and part (c), we first notice that (26) has always solutions. If (25) has a solution, by taking  $(x_1, y_1)$  the non-trivial solution with the smallest  $x_1 > 0$  and  $y_1 > 0$ , then an easy computation shows that the divisor on  $M$  associated is

$$D := \tilde{H} - d\frac{y_1}{x_1(n-1)}B.$$

Hence, the fact that  $x_1$  is the smallest possible, guarantees that  $D$  has the smallest slope, with respect to the divisors associated to the other solutions to (25).

Since we know that a wall for the movable cone is  $\tilde{H}$ , which is of Hilbert-Chow type and that  $D$  is associated to some divisorial contractions (on a certain birational model of  $M$ ),  $D$  must be the other wall.

The case in which (25) has no solution follows similarly.  $\square$

**Example 13.2.** If  $d = n-2$ , then

$$\text{Mov}(M) = \langle \tilde{H}, \tilde{H} - \frac{n-2}{n-1}B \rangle.$$

To fully understand the structure of the nef cone, we start with the easy case  $n = 2$ . Consider the Pell's equation

$$(27) \quad X^2 - dY^2 = 5.$$

The associated spherical class is  $\mathbf{s} = (r, cH, r-1)$ ,  $X = 2r-1$  and  $Y = 2c$ .

**Lemma 13.3.** *Let  $M = \text{Hilb}^2(X)$ . The nef cone of  $M$  has the following form:*

(a) *If (27) has no solutions, then*

$$\text{Nef}(M) = \text{Mov}(M).$$

(b) If (27) has solutions, we let  $(x_1, y_1)$  be a solution with the smallest  $x_1 > 0$  and  $y_1 > 0$  even. Then

$$\text{Nef}(M) = \langle \tilde{H}, \tilde{H} - d \frac{y_1}{x_1} B \rangle.$$

*Proof.* The only possibility in which the movable cone is not equal to the nef cone is if there is a flop induced by a spherical class. By Theorem 5.7, this happens if and only if (27) has a solution.  $\square$

**Example 13.4.** Let  $d = 31$ . Then the nef cone for  $M = \text{Hilb}^2(X)$  is

$$\text{Nef}(M) = \langle \tilde{H}, \tilde{H} - \frac{3658}{657} B \rangle.$$

In particular, this gives a negative answer to [CK12, Question 8.4].

Indeed, (27) has the smallest solution given by  $x_1 = 657$  and  $y_1 = 118$ . This gives a  $(-2)$ -class  $s = (329, -59 \cdot H, 328)$ , which induces a flop, by Lemma 13.3.

For higher  $n > 2$  the situation is more complicated, since the number of Pell's equations to consider is higher. But, in any case, everything is completely determined.

**Example 13.5.** Consider the case in which  $d = 1$  and  $n = 7$ ,  $M = \text{Hilb}^7(X)$ . This example exhibits a flop of “higher degree”, in the following sense: it is induced by a decomposition  $\mathbf{v} = \mathbf{a} + \mathbf{b}$ , with  $\mathbf{a}^2, \mathbf{b}^2 > 0$ , and it is not induced by a spherical or isotropic class. Indeed,  $\mathbf{v} = (1, 0, -6)$ ,  $\mathbf{a} = (1, -H, 0)$  and  $\mathbf{b} = (0, H, -6)$  give the wanted example. We also notice that the rank two hyperbolic lattice associated to this wall contains no spherical or isotropic classes. The full list of walls in the movable cone is as follows. We consider the divisor class  $\tilde{H} - \Gamma B$ , for  $\Gamma \in \mathbb{Q}_{>0}$ . The walls in the movable cone of  $M$  are given by the following table:

$\Gamma$	$\mathbf{a}$	$(\mathbf{v}, \mathbf{a})$	Type
0	$(0, 0, -1)$	1	HC divisorial contraction
$\frac{1}{4}$	$(1, -H, 2)$	4	BN flop
$\frac{2}{7}$	$(1, -H, 1)$	5	LGU flop
$\frac{1}{3}$	$(1, -H, 0)$	6	higher degree flop
$\frac{6}{17}$	$(2, -3H, 5)$	7	fake wall
$\frac{4}{11}$	$(1, -2H, 5)$	1	BN flop
$\frac{3}{8}$	$-(1, -3H, 10)$	4	BN flop
$\frac{2}{5}$	$(1, -2H, 4)$	2	LGU divisorial contraction

**...and higher Picard number.** Let  $X$  be a K3 surface such that  $\text{Pic}(X) \cong \mathbb{Z} \cdot \xi_1 \oplus \mathbb{Z} \cdot \xi_2$ .

**Example 13.6.** We let  $M := \text{Hilb}^2(X)$ , and  $\mathbf{v} = (1, 0, -1)$ . We assume that the intersection form (with respect to the basis  $\xi_1, \xi_2$ ) is given by

$$q = \begin{pmatrix} 28 & 0 \\ 0 & -4 \end{pmatrix}.$$

Such a K3 surface exists, see [Mor84, Kov94]. We have:

$$\text{NS}(M) = \mathbb{Z} \cdot \mathbf{s} \oplus \text{NS}(X),$$

where  $\mathbf{s} = (1, 0, 1)$ . Our first claim is

$$(28) \quad \text{Nef}(M) = \text{Mov}(M).$$

Indeed, by Theorem 5.7, a flopping contraction would have to come from a class  $\mathbf{a}$  with  $\mathbf{a}^2 \geq -2$  and  $(\mathbf{v}, \mathbf{a}) = 1$ ; also, the corresponding lattice  $\mathcal{H} = \langle \mathbf{v}, \mathbf{a} \rangle$  has to be hyperbolic, which implies  $\mathbf{a}^2 \leq 0$ . In addition,  $\mathbf{a}^2 = 0$  would correspond to the Hilbert-Chow divisorial contraction, and thus  $\mathbf{a}^2 = -2$  is the only possibility. If we write  $\mathbf{a} = (r, D, r+1)$  with  $D = a\xi_1 + b\xi_2$ , this gives

$$-2r(r+1) + 28a^2 - 4b^2 = -2.$$

This equation has no solutions modulo 4.

The structure of the nef cone is thus determined by divisorial contractions. These are controlled by the quadratic equation

$$(29) \quad X^2 - 2(7a^2 - b^2) = 1,$$

where  $X = r$ ,  $\mathbf{a} = (r, D, r)$ . For example, the Hilbert-Chow contraction corresponds to the solution  $a = b = 0$  and  $X = 1$  to (29). Other contractions arise, for example, at  $a = 4, b = 2, X = 15$ , or  $a = 2, b = 2, X = 7$ , etc. The nef cone will be a non-round non-finitely generated cone. Its walls have infinitely many accumulations points at the boundary of the positive cone: these come from solutions of

$$X^2 - 2(7a^2 - b^2) = 0,$$

corresponding to Lagrangian fibrations.

To simplify the computations and obtain examples of a round nef (or movable) cone, we consider a twist by a Brauer class  $\alpha \in \text{Br}(X)$ . Given  $X$  as before, we can assume that  $\alpha$  admits a  $B$ -field lift  $B$  with the properties that

$$B \cdot \text{NS}(X) = 0 \quad \text{and} \quad B^2 = 0.$$

(See [HMS08] for more details; in particular, the fact that such K3 surface exists follows as in [HMS08, Lemma 3.22].)

**Example 13.7.** We assume that the (twisted) intersection form on

$$H_{\text{alg}}^*(X, \alpha, \mathbb{Z}) = \text{NS}(X) \oplus \mathbb{Z} \cdot (2, 2B, 0) \oplus \mathbb{Z} \cdot (0, 0, -1)$$

takes the form

$$q = \begin{pmatrix} 4 & 0 & 0 & 0 \\ 0 & -4 & 0 & 0 \\ 0 & 0 & 0 & 2 \\ 0 & 0 & 2 & 0 \end{pmatrix}.$$

Consider the primitive vector  $\mathbf{v} = (0, \xi_1, 0)$ , and let  $M := M_H(\mathbf{v})$  be the moduli space of  $\alpha$ -twisted  $H$ -Gieseker semistable sheaves on  $X$ , for  $H$  a generic polarization on  $X$ . Then:

- (a)  $\text{Nef}(M) = \text{Mov}(M)$ ;

(b)  $\text{Nef}(M)$  is a rational circular cone.

To prove the above statements, observe that  $\mathbf{v}^2 = 4$  and  $(\mathbf{v}, \mathbf{a}) \in 4\mathbb{Z}$  for all  $\mathbf{a} \in H_{\text{alg}}^*(X, \alpha, \mathbb{Z})$ . According to Theorem 5.7, the only possible wall in this situation would be given by a Brill-Noether divisorial contraction, coming from a spherical class  $\mathbf{s} \in \mathbf{v}^\perp$ . But the above lattice admits no spherical classes, and thus there are no walls.

Thus the nef cone and the closure of the movable cone are both equal to the positive cone. Since  $M$  obviously admits Lagrangian fibrations, the cone is rational.

Modifying slightly the previous example, we obtain a moduli space with circular movable cone and locally polyhedral nef cone:

**Example 13.8.** We assume that the (twisted) intersection form on

$$H_{\text{alg}}^*(X, \alpha, \mathbb{Z}) = \text{NS}(X) \oplus \mathbb{Z} \cdot (3, 3B, 0) \oplus \mathbb{Z} \cdot (0, 0, -1)$$

takes the form

$$q = \begin{pmatrix} 6 & 0 & 0 & 0 \\ 0 & -6 & 0 & 0 \\ 0 & 0 & 0 & 3 \\ 0 & 0 & 3 & 0 \end{pmatrix}.$$

Consider the primitive vector  $\mathbf{v} = (0, \xi_1, 1)$ , and let  $M := M_H(\mathbf{v})$ . Then:

- (a)  $\text{Nef}(M)$  is a rational locally-polyhedral cone;
- (b)  $\text{Mov}(M)$  is a rational circular cone.

Indeed, (b) follows exactly as in Example 13.7: there are no spherical classes, and, for all  $\mathbf{a} \in H_{\text{alg}}^*(X, \alpha, \mathbb{Z})$ ,  $(\mathbf{a}, \mathbf{v}) \in 3\mathbb{Z}$ . However, flopping contractions are induced by solutions to the quadratic equation

$$a^2 - b^2 - 2as + s = 0,$$

where we set  $D = a\xi_1 + b\xi_2$ , and  $\mathbf{a} = (3(2a - 1), a\xi_1 + b\xi_2 + 3(2a - 1)B, s)$ . This has infinitely many solutions. It is an easy exercise to deduce (a) from this.

#### 14. THE GEOMETRY OF FLOPPING CONTRACTIONS

One can also refine the analysis leading to Theorem 5.7 to give a precise description of the geometry of the flopping contraction associated to a flopping wall  $\mathcal{W}$ .

As in Section 5, we let  $\sigma_0 \in \mathcal{W}$  be a stability condition on the wall, and  $\sigma_+ \notin \mathcal{W}$  be sufficiently close to  $\sigma_0$ . For simplicity, let us assume throughout this section that the hyperbolic lattice  $\mathcal{H}_{\mathcal{W}}$  associated to  $\mathcal{W}$  via Definition 5.2 does not admit spherical or isotropic classes; in particular,  $\mathcal{W}$  is not a totally semistable wall for any class  $\mathbf{a} \in \mathcal{H}$ , and does not induce a divisorial contraction.

Let  $\mathfrak{P}$  be the set of unordered partitions  $P = [\mathbf{a}_i]_i$  of  $\mathbf{v}$  into a sum  $\mathbf{v} = \mathbf{a}_1 + \cdots + \mathbf{a}_m$  of positive classes  $\mathbf{a}_i \in \mathcal{H}$ . We say that a partition  $P$  is a refinement of another partition  $Q = [\mathbf{b}_i]_i$  if it can be obtained by choosing partitions of each  $\mathbf{b}_i$ . This defines a natural partial order on  $\mathfrak{P}$ , with  $P \prec Q$  if  $P$  is a refinement of  $Q$ . The trivial partition is the maximal element of  $\mathfrak{P}$ .

Given  $P = [\mathbf{a}_i]_i \in \mathfrak{P}$ , we let  $M_P \subset M_{\sigma_+}(\mathbf{v})$  be the subset of objects  $E$  such that the Mukai vectors of the Jordan-Hölder factors  $E_i$  of  $E$  with respect to  $\sigma_0$  are given by  $\mathbf{a}_i$  for all  $i$ . Using openness of stability and closedness of semistability in families, one easily proves:

**Lemma 14.1.** *The disjoint union  $M_{\sigma_+}(\mathbf{v}) = \coprod_{P \in \mathfrak{P}} M_P$  defines a stratification of  $M_{\sigma_+}(\mathbf{v})$  into locally closed subsets, such that  $M_P$  is contained in the closure of  $M_Q$  if and only if  $P \prec Q$ .*

In addition, our simplifying assumptions on  $\mathcal{H}_{\mathcal{W}}$  give the following:

**Lemma 14.2.** *Assume that  $P = [\mathbf{a}_1, \mathbf{a}_2]$  is a two-element partition of  $\mathbf{v}$ . Then  $M_P \subset M_{\sigma_+}(\mathbf{v})$  is non-empty, and of codimension  $(\mathbf{a}_1, \mathbf{a}_2) - 1$ .*

*Proof.* Since  $\mathbf{v}$  is primitive, we may assume that  $\mathbf{a}_1$  has smaller phase than  $\mathbf{a}_2$  with respect to  $\sigma_+$ . By assumption on  $\mathcal{H}_{\mathcal{W}}$  and by Theorem 2.13, the generic element  $A_i \in M_{\sigma_+}(\mathbf{a}_i)$  is  $\sigma_0$ -stable for  $i = 1, 2$ . In particular,  $\text{Hom}(A_1, A_2) = \text{Hom}(A_2, A_1) = 0$ , and therefore  $\dim \text{Ext}^1(A_2, A_1) = (\mathbf{a}_1, \mathbf{a}_2)$ . By [BM11, Lemma 5.9], any non-trivial extension  $A_1 \hookrightarrow E \rightarrow A_2$  is  $\sigma_+$ -stable. Using Theorem 2.13 again, one computes the dimension of the space of such extensions as

$$\mathbf{a}_1^2 + 2 + \mathbf{a}_2^2 + 2 + (\mathbf{a}_1, \mathbf{a}_2) - 1 = \mathbf{v}^2 + 2 - ((\mathbf{a}_1, \mathbf{a}_2) - 1).$$

□

For  $P$  as above, the flopping contraction  $\pi^+$  contracts  $M_P$  to the product of moduli spaces  $M_{\sigma_0}^{st}(\mathbf{a}_1) \times M_{\sigma_0}^{st}(\mathbf{a}_2)$  of  $\sigma_0$ -stable objects. The contracted locus of  $\pi^+$  is the union of  $M_P$  for all non-trivial partitions  $M_P$ . In particular, when there is more than one way to write  $\mathbf{v}$  as a sum of two positive classes, our stratification is only partially ordered; this leads to a generalization of Markman's notion of *stratified Mukai flops* introduced in [Mar01] (where the stratification is indexed by a totally ordered set).

For any given  $m \in \mathbb{Z}_{>0}$ , one can easily construct examples where the locus contracted by  $\pi^+$  has  $m$  irreducible components: by Lemma 14.2, this is equivalent to constructing a lattice  $\mathcal{H}$  such that  $\mathbf{v}$  can be written as a sum  $\mathbf{v} = \mathbf{a}_1 + \mathbf{a}_2$  in exactly  $m$  different ways:

**Example 14.3.** Choose  $M \gg m$  for which  $x^2 + Mxy + y^2 = -1$  does not admit an integral solution. We define the symmetric pairing on  $\mathcal{H} \cong \mathbb{Z}^2$  via the matrix  $\begin{pmatrix} 2 & M \\ M & 2 \end{pmatrix}$ , and let  $\mathbf{v} = \begin{pmatrix} 1 \\ m-1 \end{pmatrix}$ . The positive cone is the open cone containing the upper right quadrant that is bordered by the lines of slopes approximately  $-\frac{1}{M}$  and  $-M$ . Since  $M \gg m$  (in fact,  $M > 2m$  is enough), any partition of  $\mathbf{v}$  into positive classes is in fact a partition in  $\mathbb{Z}_{\geq 0}^2$ . Therefore, the two-element partitions are given by  $\mathbf{v} = \begin{pmatrix} 1 \\ k \end{pmatrix} + \begin{pmatrix} 0 \\ m-1-k \end{pmatrix}$  for  $0 \leq k \leq m-1$ . There is a unique minimal partition  $Q$ , given by  $\mathbf{v} = \begin{pmatrix} 1 \\ 0 \end{pmatrix} + (m-1) \begin{pmatrix} 0 \\ 1 \end{pmatrix}$ . The corresponding stratum  $M_Q$  is contained in the closure of any other stratum.

Similarly, one can construct flopping contractions with arbitrarily many *connected* components:

**Example 14.4.** Let  $m$  be an odd positive integer. Choose  $M \gg m$  and define the lattice  $\mathcal{H}$  by the matrix  $\begin{pmatrix} -4 & 2M \\ 2M & 4 \end{pmatrix}$ . The positive cone lies between the lines of slope approximately  $+\frac{1}{M}$  and  $-M$ . We let  $\mathbf{v} = \begin{pmatrix} m \\ 2 \end{pmatrix}$ . Any summand in a partition of  $\mathbf{v}$  must be of the form  $\begin{pmatrix} x \\ y \end{pmatrix}$  with  $x \geq 0$  and  $y > 0$ , and therefore  $y = 1$ . Besides the trivial element, the only partitions

occurring in  $\mathfrak{P}$  are therefore of the form  $A_k = \left[ \binom{k}{1}, \binom{m-k}{1} \right]$ , for  $0 \leq k < \frac{m}{2}$ . Each corresponding strata  $M_{A_k}$  is one connected component of the exceptional locus of  $\pi^+$ .

**Remark 14.5.** To show that the lattices  $\mathcal{H}$  as above occur as lattice associated to some wall  $\mathcal{H}_{\mathcal{W}}$  on some K3 surface, we only have to find a K3 surface  $X$  such that  $\mathcal{H}$  embeds into its Mukai lattice  $H_{\text{alg}}^*(X, \mathbb{Z})$ . In particular, we may choose  $\text{Pic}(X) \cong \mathcal{H}$ , and the class  $\mathbf{v}$  to be of the form  $\mathbf{v} = (0, c, 0)$  for a corresponding curve class  $c$ . In particular, Example 14.3 occurs in a relative Jacobian of curves on special double covers  $X \rightarrow \mathbb{P}^2$ , and Example 14.4 in special quartics  $X \subset \mathbb{P}^3$ . This wall-crossing already occurs for Giesker stability with respect to a non-generic polarization  $H$ . The morphism  $\pi^+$  contracts sheaves supported on reducible curves  $C = C_1 \cup C_2$  in the corresponding linear system; it forgets the gluing data at the intersection points  $C_1 \cap C_2$ . The flopping transformation preserves the Lagrangian fibration given by the Beauville integrable system.

## 15. LE POTIER'S STRANGE DUALITY FOR ISOTROPIC CLASSES

In this section, we will explain a relation of Theorem 1.5 to Le Potier's Strange Duality Conjecture for K3 surfaces. We thank Dragos Oprea for pointing us to this application.

We first recall the basic construction from [LP05, MO08]. Let  $(X, \alpha)$  be a twisted K3 surface and let  $\sigma \in \text{Stab}(X, \alpha)$  be a generic stability condition. Let  $\mathbf{v}, \mathbf{w} \in H_{\text{alg}}^*(X, \alpha, \mathbb{Z})$  be primitive Mukai vectors with  $\mathbf{v}^2, \mathbf{w}^2 \geq 0$ . We denote by  $L_{\mathbf{w}}$  (resp.,  $L_{\mathbf{v}}$ ) the line bundle  $\mathcal{O}_{M_{\sigma}(\mathbf{v})}(-\theta_{\mathbf{v}}(\mathbf{w}))$  (resp.,  $\mathcal{O}_{M_{\sigma}(\mathbf{w})}(-\theta_{\mathbf{w}}(\mathbf{v}))$ ). We assume:

- $(\mathbf{v}, \mathbf{w}) = 0$ , and
- for all  $E \in M_{\sigma}(\mathbf{v})$  and all  $F \in M_{\sigma}(\mathbf{w})$ ,  $\text{Hom}^2(E, F) = 0$ .

Then the locus

$$\Theta = \{(E, F) \in M_{\sigma}(\mathbf{v}) \times M_{\sigma}(\mathbf{w}) : \text{Hom}(E, F) \neq 0\}$$

gives rise to a section of the line bundle  $L_{\mathbf{v}, \mathbf{w}} := L_{\mathbf{w}} \boxtimes L_{\mathbf{v}}$  on  $M_{\sigma}(\mathbf{v}) \times M_{\sigma}(\mathbf{w})$  (which may or may not vanish). We then obtain a morphism, well-defined up to scalars,

$$\text{SD}: H^0(M_{\sigma}(\mathbf{v}), L_{\mathbf{w}})^{\vee} \longrightarrow H^0(M_{\sigma}(\mathbf{w}), L_{\mathbf{v}}).$$

The two basic questions are:

- When is  $h^0(M_{\sigma}(\mathbf{v}), L_{\mathbf{w}}) = h^0(M_{\sigma}(\mathbf{w}), L_{\mathbf{v}})$ ?
- If equality holds, is the map SD an isomorphism?

We answer the two previous questions in the case where one of the two vectors is isotropic:

**Proposition 15.1.** *Let  $(X, \alpha)$  be a twisted K3 surface and let  $\sigma \in \text{Stab}(X, \alpha)$  be a generic stability condition. Let  $\mathbf{v}, \mathbf{w} \in H_{\text{alg}}^*(X, \alpha, \mathbb{Z})$  be primitive Mukai vectors with  $(\mathbf{v}, \mathbf{w}) = 0$ ,  $\mathbf{v}^2 \geq 2$  and  $\mathbf{w}^2 = 0$ .*

*We assume that  $-\theta_{\mathbf{v}}(\mathbf{w}) \in \text{Mov}(M_{\sigma}(\mathbf{v}))$  and  $-\theta_{\mathbf{w}}(\mathbf{v}) \in \text{Nef}(M_{\sigma}(\mathbf{w}))$ . Then*

- (a)  $h^0(M_{\sigma}(\mathbf{v}), L_{\mathbf{w}}) = h^0(M_{\sigma}(\mathbf{w}), L_{\mathbf{v}})$ , and
- (b) *the morphism SD is either zero or an isomorphism.*

*Proof.* Let  $Y := M_{\sigma}(\mathbf{w})$ . By [Muk87a, Că102, Yos06], there exist an element  $\beta \in \text{Br}(Y)$  and a derived equivalence  $\Phi: D^b(X, \alpha) \xrightarrow{\sim} D^b(Y, \beta)$ . Replacing  $(X, \alpha)$  by  $(Y, \beta)$ , we may assume that  $\mathbf{w} = (0, 0, 1)$  and  $\mathbf{v} = (0, D, s)$ , for some  $s \in \mathbb{Z}$  and  $D \in \text{NS}(X)$ , and that  $X = M_{\sigma}(\mathbf{w})$  is the moduli space of skyscraper sheaves. Moreover,  $D = -\theta_{\mathbf{w}}(\mathbf{v}) \in \text{Nef}(X)$



is effective, by assumption. By stability and Serre duality, for all  $E \in M_\sigma(\mathbf{v})$  and all  $x \in X$ ,  $\mathrm{Hom}^2(E, k(x)) = \mathrm{Hom}(k(x), E)^\vee = 0$ , and the locus  $\Theta$  gives a section of  $L_{\mathbf{w}} \boxtimes L_{\mathbf{v}}$ .

By Remark 11.5, there exists a chamber  $\mathcal{L}_\infty$  in the interior of the movable cone  $\mathrm{Mov}(M_\sigma(\mathbf{v}))$  whose boundary contains  $-\theta_{\mathbf{v}}(\mathbf{w})$ . Moreover, there exist a polarization  $H$  on  $X$  and a chamber  $\mathcal{C}_\infty \subset \mathrm{Stab}(X, \alpha)$  such that  $\ell(\mathcal{C}_\infty) = \mathcal{L}_\infty$ ,  $M_H(\mathbf{v}) = M_{\mathcal{C}_\infty}(\mathbf{v})$ , and the Lagrangian fibration induced by  $\mathbf{w}$  is the Beauville integrable system on  $M_H(\mathbf{v})$ .

The argument in [MO08, Example 8] shows that  $h^0(M_H(\mathbf{v}), L_{\mathbf{w}}) = h^0(X, \mathcal{O}(D))$  and the morphism SD is an isomorphism. Since  $M_H(\mathbf{v})$  is connected to  $M_\sigma(\mathbf{v})$  by a sequence of flops, which do not change the dimension of the spaces of sections of  $L_{\mathbf{w}}$ , we obtain immediately (a).

To prove (b), we need to study the behavior of the morphism SD under wall-crossing. We pick a stability condition  $\sigma_\infty \in \mathcal{C}_\infty$ . Both  $\sigma$  and  $\sigma_\infty$  belong to the open subset  $U(X, \alpha)$  of Theorem 2.7. By Theorem 10.2, we can find a path in  $U(X, \alpha)$  connecting  $\sigma$  and  $\sigma_\infty$  which crosses only fake or flopping walls with respect to  $\mathbf{v}$ . If it crosses no totally semistable walls, then the morphism SD is compatible with the wall-crossing; since it induces an isomorphism at  $\sigma_\infty$ , it induces an isomorphism at  $\sigma$ .

Assume instead that there is a totally semistable wall. We write  $\sigma = \sigma_{\omega, \beta}$ . The straight path from  $\sigma_\infty$  to  $\sigma_{t\omega, \beta}$ , for  $t \gg 0$ , corresponds to a change of polarization for Gieseker stability, and thus does not cross any totally semistable wall. Therefore, we may replace  $\sigma_\infty$  with  $\sigma_{t\omega, \beta}$ , for  $t \gg 0$ .

We claim that all objects  $E$  in  $M_\sigma(\mathbf{v})$  must be actual complexes, with  $\mathrm{rk}(H^0(E)) > 0$ . Indeed, if there exists a sheaf  $E$  in  $M_\sigma(\mathbf{v})$ , then the generic element is a sheaf. Moreover, since  $D$  is nef and big, it is globally generated, and we can assume that the support of  $E$  is a smooth integral curve. Stability in  $U(X, \alpha)$  for torsion sheaves implies, in particular, that the sheaf is actually stable on the curve. But then  $E$  would be stable for  $t \rightarrow \infty$ . This shows that we crossed no totally semistable wall.

Hence, since  $E$  is an actual complex,  $\mathrm{Hom}(E, k(x)) \neq 0$ , for all  $x \in X$ . This shows that  $\Theta$  is nothing but the zero-section of  $L_{\mathbf{v}, \mathbf{w}}$  and the induced map SD is the zero map.  $\square$

By applying the previous proposition to stability conditions near the large volume limit, we deduce the corresponding strange duality statement for moduli of stable sheaves (with respect to a generic polarization on  $X$ ).

**Example 15.2.** Let  $X$  be a K3 surface such that  $\mathrm{Pic}(X) = \mathbb{Z} \cdot H$ , with  $H^2 = 2$ . Let  $\mathbf{v} = (1, 0, -1)$  and  $\mathbf{w} = -(1, -H, 1)$ . Consider a stability condition  $\sigma_\infty = \sigma_{tH, -2H}$ , for  $t \gg 0$ . Then, as observed in [Bea99, Proposition 1.3],  $\mathrm{Hilb}^2(X) = M_{\sigma_\infty}(\mathbf{v})$  admits a flop to a Lagrangian fibration induced by the vector  $\mathbf{w}$ . The assumptions of Proposition 15.1 are satisfied. In this case, for all  $E[1] \in M_{\sigma_\infty}(\mathbf{w})$ ,  $E \cong I_{pt}(-H)$ , and for all  $\Gamma \in \mathrm{Hilb}^2(X)$ , we have  $\mathrm{Hom}(I_\Gamma, E[1]) \neq 0$ . Hence, the map SD is the zero map.

The following example shows that the assumption in Proposition 15.1 is necessary:

**Example 15.3.** Let  $X$  be a K3 surface with  $\mathrm{NS}(X) = \mathbb{Z} \cdot C_1 \oplus \mathbb{Z} \cdot C_2$  and intersection form

$$q = \begin{pmatrix} -2 & 4 \\ 4 & -2 \end{pmatrix}.$$

We assume the two rational curves  $C_1$  and  $C_2$  generate the cone of effective divisors on  $X$ . Let  $\mathbf{v} = (0, 3C_1 + C_2, 1)$  and  $\mathbf{w} = (0, 0, 1)$ . Then  $\mathbf{v}^2 = 4$ . Pick a generic ample divisor  $H$  on  $X$ . We have

$$H^0(M_H(\mathbf{v}), \theta_{\mathbf{v}}(\mathbf{w})) \cong \mathbb{C}^{\oplus 4}.$$

Indeed, for example, one can argue by flopping at the spherical vector  $(0, C_1, 0)$ . Then we have a birational map  $M_H(\mathbf{v}) \dashrightarrow M_H(\mathbf{v}')$ , where  $\mathbf{v}' = (0, C_1 + C_2, 1)$  and a chain of isomorphisms

$$H^0(M_H(\mathbf{v}), \theta_{\mathbf{v}}(\mathbf{w})) \cong H^0(M_H(\mathbf{v}'), \theta_{\mathbf{v}'}(\mathbf{w})) \cong H^0(\mathbb{P}^3, \mathcal{O}_{\mathbb{P}^3}(1)) \cong \mathbb{C}^{\oplus 4}.$$

On the other side, we have

$$H^0(M_H(\mathbf{w}), \theta_{\mathbf{w}}(\mathbf{v})) \cong H^0(X, \mathcal{O}_X(3C_1 + C_2)) \cong \mathbb{C}^{\oplus 5}.$$

The last isomorphism follows from the exact sequence

$$0 \rightarrow \mathcal{O}_X(2C_1 + C_2) \rightarrow \mathcal{O}_X(3C_1 + C_2) \rightarrow \mathcal{O}_{\mathbb{P}^1}(-2) \rightarrow 0,$$

since  $\mathcal{O}_X(2C_1 + C_2)$  is big and nef and thus has no higher cohomology.

## REFERENCES

- [AB13] Daniele Arcara and Aaron Bertram. Bridgeland-stable moduli spaces for  $K$ -trivial surfaces. *J. Eur. Math. Soc. (JEMS)*, 15(1):1–38, 2013. With an appendix by Max Lieblich. arXiv:0708.2247.
- [ABCH13] Daniele Arcara, Aaron Bertram, Izzet Coskun, and Jack Huizenga. The minimal model program for the Hilbert scheme of points on  $\mathbb{P}^2$  and Bridgeland stability. *Adv. Math.*, 235:580–626, 2013. arXiv:1203.0316.
- [Ame11] Ekaterina Amerik. A remark on a question of Beauville about lagrangian fibrations, 2011. arXiv:1110.2852.
- [AP06] Dan Abramovich and Alexander Polishchuk. Sheaves of  $t$ -structures and valuative criteria for stable complexes. *J. Reine Angew. Math.*, 590:89–130, 2006. arXiv:math/0309435.
- [BC12] Aaron Bertram and Izzet Coskun. The birational geometry of the Hilbert scheme of points on surfaces, 2012. Available at: <http://homepages.math.uic.edu/~coskun/>.
- [BCHM10] Caucher Birkar, Paolo Cascini, Christopher D. Hacon, and James McKernan. Existence of minimal models for varieties of log general type. *J. Amer. Math. Soc.*, 23(2):405–468, 2010. arXiv:math/0610203.
- [Bea83] Arnaud Beauville. Variétés Kähleriennes dont la première classe de Chern est nulle. *J. Differential Geom.*, 18(4):755–782 (1984), 1983.
- [Bea91] A. Beauville. Systèmes hamiltoniens complètement intégrables associés aux surfaces  $K3$ . In *Problems in the theory of surfaces and their classification (Cortona, 1988)*, Sympos. Math., XXXII, pages 25–31. Academic Press, London, 1991.
- [Bea99] Arnaud Beauville. Counting rational curves on  $K3$  surfaces. *Duke Math. J.*, 97(1):99–108, 1999. arXiv:alg-geom/9701019.
- [Bea10] Arnaud Beauville. Holomorphic symplectic geometry: a problem list, 2010. arXiv:1002.4321.
- [BJ11] Benjamin Bakker and Andrei Jorza. Lagrangian hyperplanes in holomorphic symplectic varieties, 2011. arXiv:1111.0047.
- [BM11] Arend Bayer and Emanuele Macrì. The space of stability conditions on the local projective plane. *Duke Math. J.*, 160(2):263–322, 2011. arXiv:0912.0043.
- [BM12] Arend Bayer and Emanuele Macrì. Projectivity and birational geometry of Bridgeland moduli spaces, 2012. arXiv:1203.4613.
- [BMT11] Arend Bayer, Emanuele Macrì, and Yukinobu Toda. Bridgeland stability conditions on threefolds I: Bogomolov-Gieseker type inequalities, 2011. To appear in *J. Alg. Geom.* arXiv:1103.5010.
- [BMW13] Aaron Bertram, Cristian Martinez, and Jie Wang. The birational geometry of moduli space of sheaves on the projective plane, 2013. arXiv:1301.2011.
- [Bou04] Sébastien Boucksom. Divisorial Zariski decompositions on compact complex manifolds. *Ann. Sci. École Norm. Sup. (4)*, 37(1):45–76, 2004.
- [Bri07] Tom Bridgeland. Stability conditions on triangulated categories. *Ann. of Math. (2)*, 166(2):317–345, 2007. arXiv:math/0212237.
- [Bri08] Tom Bridgeland. Stability conditions on  $K3$  surfaces. *Duke Math. J.*, 141(2):241–291, 2008. arXiv:math/0307164.
- [Că100] Andrei Căldăraru. *Derived categories of twisted sheaves on Calabi-Yau manifolds*. ProQuest LLC, Ann Arbor, MI, 2000. Thesis (Ph.D.)—Cornell University.

- [Că102] Andrei Căldăraru. Nonfine moduli spaces of sheaves on  $K3$  surfaces. *Int. Math. Res. Not.*, 20:1027–1056, 2002. arXiv:math/0108180.
- [CK12] Ciro Ciliberto and Andreas Leopold Knutsen. On  $k$ -gonal loci in Severi varieties on general  $K3$  surfaces and rational curves on hyperkähler manifolds, 2012. arXiv:1204.4838.
- [DH98] Igor V. Dolgachev and Yi Hu. Variation of geometric invariant theory quotients. *Inst. Hautes Études Sci. Publ. Math.*, 87:5–56, 1998. With an appendix by Nicolas Ressayre. arXiv:alg-geom/9402008.
- [EG95] Geir Ellingsrud and Lothar Göttsche. Variation of moduli spaces and Donaldson invariants under change of polarization. *J. Reine Angew. Math.*, 467:1–49, 1995. arXiv:alg-geom/9410005.
- [FQ95] Robert Friedman and Zhenbo Qin. Flips of moduli spaces and transition formulas for Donaldson polynomial invariants of rational surfaces. *Comm. Anal. Geom.*, 3(1-2):11–83, 1995. arXiv:alg-geom/9410007.
- [FS10] Maksym Fedorchuk and David Ishii Smyth. Alternate compactifications of moduli spaces of curves, 2010. arXiv:1012.0329.
- [Fuj11] Osamu Fujino. On Kawamata’s theorem. In *Classification of algebraic varieties*, EMS Ser. Congr. Rep., pages 305–315. Eur. Math. Soc., Zürich, 2011. arXiv:0910.1156.
- [GHJ03] M. Gross, D. Huybrechts, and D. Joyce. *Calabi-Yau manifolds and related geometries*. Universitext. Springer-Verlag, Berlin, 2003. Lectures from the Summer School held in Nordfjordeid, June 2001.
- [GLR11a] Daniel Greb, Christian Lehn, and Sönke Rollenske. Lagrangian fibrations on hyperkähler manifolds - On a question of Beauville, 2011. arXiv:1105.3410.
- [GLR11b] Daniel Greb, Christian Lehn, and Sönke Rollenske. Lagrangian fibrations on hyperkähler manifolds: intersections of Lagrangians,  $l$ -reduction, and fourfolds, 2011. arXiv:1110.2680.
- [HHT12] David Harvey, Brendan Hassett, and Yuri Tschinkel. Characterizing projective spaces on deformations of Hilbert schemes of  $K3$  surfaces. *Comm. Pure Appl. Math.*, 65(2):264–286, 2012. arXiv:1011.1285.
- [HL10] Daniel Huybrechts and Manfred Lehn. *The geometry of moduli spaces of sheaves*. Cambridge Mathematical Library. Cambridge University Press, Cambridge, second edition, 2010.
- [HMS08] Daniel Huybrechts, Emanuele Macrì, and Paolo Stellari. Stability conditions for generic  $K3$  categories. *Compos. Math.*, 144(1):134–162, 2008. arXiv:math/0608430.
- [HMS09] Daniel Huybrechts, Emanuele Macrì, and Paolo Stellari. Derived equivalences of  $K3$  surfaces and orientation. *Duke Math. J.*, 149(3):461–507, 2009. arXiv:0710.1645.
- [HS05] Daniel Huybrechts and Paolo Stellari. Equivalences of twisted  $K3$  surfaces. *Math. Ann.*, 332(4):901–936, 2005. arXiv:math/0409030.
- [HS06] Daniel Huybrechts and Paolo Stellari. Proof of Căldăraru’s conjecture. In *Moduli spaces and arithmetic geometry*, volume 45 of *Adv. Stud. Pure Math.*, pages 31–42. Math. Soc. Japan, Tokyo, 2006. arXiv:math/0411541.
- [HT01] B. Hassett and Y. Tschinkel. Rational curves on holomorphic symplectic fourfolds. *Geom. Funct. Anal.*, 11(6):1201–1228, 2001. arXiv:math/9910021.
- [HT09] Brendan Hassett and Yuri Tschinkel. Moving and ample cones of holomorphic symplectic fourfolds. *Geom. Funct. Anal.*, 19(4):1065–1080, 2009. arXiv:0710.0390.
- [HT10] Brendan Hassett and Yuri Tschinkel. Intersection numbers of extremal rays on holomorphic symplectic varieties. *Asian J. Math.*, 14(3):303–322, 2010. arXiv:0909.4745.
- [Hui12] Jack Huizenga. Effective divisors on the Hilbert scheme of points in the plane and interpolation for stable bundles, 2012. arXiv:1210.6576.
- [Huy99] Daniel Huybrechts. Compact hyper-Kähler manifolds: basic results. *Invent. Math.*, 135(1):63–113, 1999. arXiv:alg-geom/9705025.
- [Huy11] Daniel Huybrechts. A global Torelli theorem for hyperkähler manifolds (after Verbitsky), 2011. arXiv:1106.5573.
- [HW12] Jun-Muk Hwang and Richard M. Weiss. Webs of Lagrangian tori in projective symplectic manifolds, 2012. arXiv:1201.2369.
- [Hwa08] Jun-Muk Hwang. Base manifolds for fibrations of projective irreducible symplectic manifolds. *Invent. Math.*, 174(3):625–644, 2008. arXiv:0711.3224.
- [Kaw85] Y. Kawamata. Pluricanonical systems on minimal algebraic varieties. *Invent. Math.*, 79(3):567–588, 1985.
- [KLS06] D. Kaledin, M. Lehn, and Ch. Sorger. Singular symplectic moduli spaces. *Invent. Math.*, 164(3):591–614, 2006. arXiv:math/0504202.
- [Kov94] Sándor J. Kovács. The cone of curves of a  $K3$  surface. *Math. Ann.*, 300(4):681–691, 1994.

- [KS08] Maxim Kontsevich and Yan Soibelman. Stability structures, motivic Donaldson-Thomas invariants and cluster transformations, 2008. arXiv:0811.2435.
- [Li93] Jun Li. Algebraic geometric interpretation of Donaldson’s polynomial invariants. *J. Differential Geom.*, 37(2):417–466, 1993.
- [Lie07] Max Lieblich. Moduli of twisted sheaves. *Duke Math. J.*, 138(1):23–118, 2007. arXiv:math/0411337.
- [Lo12] Jason Lo. On some moduli of complexes on  $K3$  surfaces, 2012. arXiv:1203.1558.
- [LP05] Joseph Le Potier. Dualité étrange, sur les surfaces, 2005.
- [LQ11] Jason Lo and Zhenbo Qin. Mini-walls for Bridgeland stability conditions on the derived category of sheaves over surfaces, 2011. arXiv:1103.4352.
- [Mac12] Antony Maciocia. Computing the walls associated to Bridgeland stability conditions on projective surfaces, 2012. arXiv:1202.4587.
- [Mar01] Eyal Markman. Brill-Noether duality for moduli spaces of sheaves on  $K3$  surfaces. *J. Algebraic Geom.*, 10(4):623–694, 2001. arXiv:math/9901072.
- [Mar06] Dimitri Markushevich. Rational Lagrangian fibrations on punctual Hilbert schemes of  $K3$  surfaces. *Manuscripta Math.*, 120(2):131–150, 2006. arXiv:math/0509346.
- [Mar09] Eyal Markman. Prime exceptional divisors on holomorphic symplectic varieties and monodromy-reflections, 2009. arXiv:0912.4981.
- [Mar11] Eyal Markman. A survey of Torelli and monodromy results for holomorphic-symplectic varieties. *Proceedings of the conference “Complex and Differential Geometry”, Springer Proceedings in Mathematics*, 2011:Volume8,257–322, 2011. arXiv:1101.4606.
- [Mar13] Eyal Markman. Lagrangian fibrations of holomorphic-symplectic varieties of  $K3^{[n]}$ -type, 2013. arXiv:1301.6584.
- [Mat99] Daisuke Matsushita. On fibre space structures of a projective irreducible symplectic manifold. *Topology*, 38(1):79–83, 1999. arXiv:alg-geom/9709033.
- [Mat01] Daisuke Matsushita. Addendum: “On fibre space structures of a projective irreducible symplectic manifold” [Topology **38** (1999), no. 1, 79–83]. *Topology*, 40(2):431–432, 2001. arXiv:math/9903045.
- [Mat12a] Daisuke Matsushita. On almost holomorphic Lagrangian fibrations, 2012. arXiv:1209.1194.
- [Mat12b] Daisuke Matsushita. Talk given at “Hyper-Kähler Geometry”, Simons Center for Geometry and Physics, 2012. Video available at: <http://scgp.stonybrook.edu/archives/3434>.
- [MM11] Antony Maciocia and Ciaran Meachan. Rank one Bridgeland stable moduli spaces on a principally polarized abelian surface, 2011. arXiv:1107.5304.
- [MM12] Eyal Markman and Sukhendu Mehrotra. Hilbert schemes of  $K3$  surfaces are dense in moduli, 2012. arXiv:1201.0031.
- [MO08] Alina Marian and Dragos Oprea. A tour of theta dualities on moduli spaces of sheaves. In *Curves and abelian varieties*, volume 465 of *Contemp. Math.*, pages 175–201. Amer. Math. Soc., Providence, RI, 2008. arXiv:0710.2908.
- [Mor84] D. R. Morrison. On  $K3$  surfaces with large Picard number. *Invent. Math.*, 75(1):105–121, 1984.
- [Muk81] Shigeru Mukai. Duality between  $D(X)$  and  $D(\hat{X})$  with its application to Picard sheaves. *Nagoya Math. J.*, 81:153–175, 1981.
- [Muk84] Shigeru Mukai. Symplectic structure of the moduli space of sheaves on an abelian or  $K3$  surface. *Invent. Math.*, 77(1):101–116, 1984.
- [Muk87a] S. Mukai. On the moduli space of bundles on  $K3$  surfaces. I. In *Vector bundles on algebraic varieties (Bombay, 1984)*, volume 11 of *Tata Inst. Fund. Res. Stud. Math.*, pages 341–413. Tata Inst. Fund. Res., Bombay, 1987.
- [Muk87b] Shigeru Mukai. Fourier functor and its application to the moduli of bundles on an abelian variety. In *Algebraic geometry, Sendai, 1985*, volume 10 of *Adv. Stud. Pure Math.*, pages 515–550. North-Holland, Amsterdam, 1987.
- [MW97] Kenji Matsuki and Richard Wentworth. Mumford-Thaddeus principle on the moduli space of vector bundles on an algebraic surface. *Internat. J. Math.*, 8(1):97–148, 1997. arXiv:alg-geom/9410016.
- [MY11a] Hiroki Minamide, Shintarou Yanagida, and Kōta Yoshioka. Fourier-mukai transforms and the wall-crossing behavior for Bridgeland’s stability conditions, 2011. arXiv:1106.5217.
- [MY11b] Hiroki Minamide, Shintarou Yanagida, and Kōta Yoshioka. Some moduli spaces of Bridgeland’s stability conditions, 2011. arXiv:1111.6187.
- [Nam01] Yoshinori Namikawa. Deformation theory of singular symplectic  $n$ -folds. *Math. Ann.*, 319(3):597–623, 2001. arXiv:math/0010113.

- [O'G99] Kieran G. O'Grady. Desingularized moduli spaces of sheaves on a  $K3$ . *J. Reine Angew. Math.*, 512:49–117, 1999. arXiv:alg-geom/9708009.
- [Orl97] D. O. Orlov. Equivalences of derived categories and  $K3$  surfaces. *J. Math. Sci. (New York)*, 84(5):1361–1381, 1997. Algebraic geometry, 7. arXiv:alg-geom/9606006.
- [Pol07] A. Polishchuk. Constant families of  $t$ -structures on derived categories of coherent sheaves. *Mosc. Math. J.*, 7(1):109–134, 167, 2007. arXiv:math/0606013.
- [Saw03] Justin Sawon. Abelian fibred holomorphic symplectic manifolds. *Turkish J. Math.*, 27(1):197–230, 2003. arXiv:math/0404362.
- [Saw07] Justin Sawon. Lagrangian fibrations on Hilbert schemes of points on  $K3$  surfaces. *J. Algebraic Geom.*, 16(3):477–497, 2007. arXiv:math/0509224.
- [ST01] Paul Seidel and Richard Thomas. Braid group actions on derived categories of coherent sheaves. *Duke Math. J.*, 108(1):37–108, 2001. arXiv:math/0001043.
- [Tha96] Michael Thaddeus. Geometric invariant theory and flips. *J. Amer. Math. Soc.*, 9(3):691–723, 1996. arXiv:alg-geom/9405004.
- [Tod08] Yukinobu Toda. Moduli stacks and invariants of semistable objects on  $K3$  surfaces. *Adv. Math.*, 217(6):2736–2781, 2008. arXiv:math.AG/0703590.
- [Ver09] Misha Verbitsky. A global Torelli theorem for hyperkähler manifolds, 2009. arXiv:0908.4121.
- [Ver10] Misha Verbitsky. HyperKähler SYZ conjecture and semipositive line bundles. *Geom. Funct. Anal.*, 19(5):1481–1493, 2010. arXiv:0811.0639.
- [Wie03] Jan Wierzbka. Contractions of symplectic varieties. *J. Algebraic Geom.*, 12(3):507–534, 2003. arXiv:math/9910130.
- [Yos99] Kōta Yoshioka. Irreducibility of moduli spaces of vector bundles on  $K3$  surfaces, 1999. arXiv:math/9907001.
- [Yos01] Kōta Yoshioka. Moduli spaces of stable sheaves on abelian surfaces. *Math. Ann.*, 321(4):817–884, 2001. arXiv:math/0009001.
- [Yos06] Kōta Yoshioka. Moduli spaces of twisted sheaves on a projective variety. In *Moduli spaces and arithmetic geometry*, volume 45 of *Adv. Stud. Pure Math.*, pages 1–30. Math. Soc. Japan, Tokyo, 2006. arXiv:math/0411538.
- [Yos09] Kōta Yoshioka. Fourier-Mukai transform on abelian surfaces. *Math. Ann.*, 345(3):493–524, 2009. arXiv:math/0605190.
- [Yos12] Kōta Yoshioka. Bridgeland's stability and the positive cone of the moduli spaces of stable objects on an abelian surface, 2012. arXiv:1206.4838.
- [YY12] Shintarou Yanagida and Kōta Yoshioka. Bridgeland's stabilities on abelian surfaces, 2012. arXiv:1203.0884.

SCHOOL OF MATHEMATICS, THE UNIVERSITY OF EDINBURGH, JAMES CLERK MAXWELL BUILDING,  
THE KING'S BUILDINGS, MAYFIELD ROAD, EDINBURGH, SCOTLAND EH9 3JZ, UNITED KINGDOM

*E-mail address:* arend.bayer@ed.ac.uk

*URL:* <http://www.maths.ed.ac.uk/~abayer/>

DEPARTMENT OF MATHEMATICS, THE OHIO STATE UNIVERSITY, 231 W 18TH AVENUE, COLUMBUS,  
OH 43210-1174, USA

*E-mail address:* macri.6@math.osu.edu

*URL:* <http://www.math.osu.edu/~macri.6/>