

INVARIANCE OF PONTRJAGIN CLASSES FOR BOTT MANIFOLDS

SUYOUNG CHOI, MIKIYA MASUDA, AND SATOSHI MURAI

ABSTRACT. A Bott manifold is the total space of some iterated $\mathbb{C}P^1$ -bundle over a point. We prove that any graded ring isomorphism between the cohomology rings of two Bott manifolds preserves their Pontrjagin classes. Moreover, we prove that such an isomorphism is induced from a diffeomorphism if the Bott manifolds are $\mathbb{Z}/2$ -trivial, where a Bott manifold is called $\mathbb{Z}/2$ -trivial if its cohomology ring with $\mathbb{Z}/2$ -coefficient is isomorphic to that of a product of $\mathbb{C}P^1$'s.

1. INTRODUCTION

One of the fundamental problems in topology is to classify manifolds (up to diffeomorphism, homeomorphism etc.) by invariants and cohomology rings are not sufficient to classify them in general. For example, the surgery theory tells us that there are infinitely many diffeomorphism types in the family of closed smooth manifolds homotopy equivalent to a complex projective space $\mathbb{C}P^n$ when $n \geq 3$. However, the surgery theory further tells us that they are distinguished by their Pontrjagin classes up to finite ambiguity, and this is true in general for the family of closed smooth manifolds homotopy equivalent to a fixed closed smooth manifold X where X is simply connected and of dimension ≥ 5 .

On the other hand, we have a feeling that most of closed smooth manifolds do not admit an effective smooth S^1 -action. For example, T. Petrie [15] conjectures that if M is a closed smooth manifold homotopy equivalent to $\mathbb{C}P^n$ and M admits an effective smooth S^1 -action, then a homotopy equivalence $f: M \rightarrow \mathbb{C}P^n$ preserves their Pontrjagin classes. Note that $\mathbb{C}P^n$ has an effective smooth action of $(S^1)^n$. The conjecture is not solved but many partial affirmative solutions are obtained. Among them, Petrie [16] shows that the conjecture is true if M admits an effective smooth action of $(S^1)^n$, see [17] for more details.

A complete non-singular toric variety (simply called a toric manifold) of complex dimension n admits an effective algebraic action of $(\mathbb{C}^*)^n$, in particular, an effective smooth action of $(S^1)^n$. The complex projective space $\mathbb{C}P^n$ is a typical example of a toric manifold. Motivated by the Petrie's conjecture and his result mentioned above, the second named author and D. Y. Suh [13] posed a problem which asks whether any cohomology ring isomorphism between toric manifolds preserves their Pontrjagin classes. Little is known about the problem. One of our main purposes is to show that the problem is affirmative for some nice class of toric manifolds called

Date: May 7, 2014.

The first author partially supported by Basic Science Research Program through the National Research Foundation of Korea(NRF) funded by the Ministry of Education(NRF-2011-0024975) and the TJ Park Science Fellowship funded by the POSCO TJ Park Foundation.

The second author was partially supported by JSPS KAKENHI 25400095.

The third author was partially supported by JSPS KAKENHI 25400043.

Bott manifolds. We even show that any cohomology ring isomorphism is induced from a diffeomorphism for a certain subclass of Bott manifolds.

A Bott tower of height n is a sequence of $\mathbb{C}P^1$ -bundles

$$(1.1) \quad B_n \xrightarrow{\pi_n} B_{n-1} \xrightarrow{\pi_{n-1}} \cdots \xrightarrow{\pi_2} B_1 \xrightarrow{\pi_1} B_0 = \{\text{a point}\},$$

where B_j is the projectivization $P(\mathbb{C} \oplus L_j)$ of a trivial complex line bundle \mathbb{C} and a complex line bundle L_j over B_{j-1} , and $\pi_j: B_j \rightarrow B_{j-1}$ is the projection for $j = 1, 2, \dots, n$. It is known that B_n is a toric manifold and B_n is called an n -stage Bott manifold or simply a Bott manifold.

If all the fibrations in (1.1) are trivial, then B_n is diffeomorphic to $(\mathbb{C}P^1)^n$. The 1-stage Bott manifold is $\mathbb{C}P^1$ and 2-stage Bott manifolds are Hirzebruch surfaces. As is well-known, there are only two diffeomorphism types among Hirzebruch surfaces; they are $(\mathbb{C}P^1)^2$ and $\mathbb{C}P^2 \# \overline{\mathbb{C}P^2}$, where $\overline{\mathbb{C}P^2}$ is $\mathbb{C}P^2$ with the opposite orientation. However, there are infinitely many diffeomorphism types among n -stage Bott manifolds when $n \geq 3$, and it is an interesting open question to classify them up to diffeomorphism or homeomorphism ([8]).

Our first main result is the following.

Theorem 1.1. *Any graded ring isomorphism between the cohomology rings with integer coefficients of two Bott manifolds preserves their Pontrjagin classes.*

Since Bott manifolds are simply connected, by combining Theorem 1.1 with a result of Novikov [14], we obtain the following corollary.

Corollary 1.1. *The number of Bott manifolds homotopy equivalent to a given Bott manifold is finite up to diffeomorphism.*

If a graded ring isomorphism between the cohomology rings of two smooth manifolds is induced from a diffeomorphism, then the isomorphism preserves their Pontrjagin classes. Hence, Theorem 1.1 provides a supporting evidence to the following conjecture.

Strong cohomological rigidity conjecture for Bott manifolds. Any graded ring isomorphism between the cohomology rings with integer coefficients of two Bott manifolds is induced from a diffeomorphism.

The conjecture above in particular claims that two Bott manifolds are diffeomorphic if their cohomology rings with integer coefficients are isomorphic as graded rings (in particular, if they are homotopy equivalent), and we call this weaker conjecture *Cohomological rigidity conjecture for Bott manifolds*. No counterexamples are known to these conjectures and some partial affirmative solutions are known; for instance, the strong cohomological rigidity conjecture is affirmative for \mathbb{Q} -trivial Bott manifolds, where a Bott manifold B_n is called \mathbb{Q} -trivial if $H^*(B_n; \mathbb{Q}) \cong H^*((\mathbb{C}P^1)^n; \mathbb{Q})$ as graded rings ([3]). It is also affirmative up to 3-stage Bott manifolds and the cohomological rigidity conjecture is affirmative for 4-stage Bott manifolds ([2]).

We say that a Bott manifold B_n is $\mathbb{Z}/2$ -trivial if $H^*(B_n; \mathbb{Z}/2) \cong H^*((\mathbb{C}P^1)^n; \mathbb{Z}/2)$ as graded rings, where $\mathbb{Z}/2 = \mathbb{Z}/2\mathbb{Z}$. Our second main result is the following.

Theorem 1.2. *The strong cohomological rigidity conjecture is affirmative for $\mathbb{Z}/2$ -trivial Bott manifolds, namely any graded ring isomorphism between the cohomology rings (with integer coefficients) of two $\mathbb{Z}/2$ -trivial Bott manifolds is induced from a diffeomorphism.*

There are infinitely many diffeomorphism types among $\mathbb{Z}/2$ -trivial n -stage Bott manifolds when $n \geq 3$ while there are only finitely many diffeomorphism types among \mathbb{Q} -trivial n -stage Bott manifolds for any n . Therefore, the family of $\mathbb{Z}/2$ -trivial Bott manifolds is much larger than that of \mathbb{Q} -trivial Bott manifolds but the former family does not contain the latter, for instance, a 2-stage Bott manifold $\mathbb{C}P^2 \# \overline{\mathbb{C}P^2}$ is not $\mathbb{Z}/2$ -trivial but \mathbb{Q} -trivial. We remark that, for each n , only $(\mathbb{C}P^1)^n$ is a Bott manifold which is both $\mathbb{Z}/2$ -trivial and \mathbb{Q} -trivial.

The rigidity conjectures mentioned above are posed (as problems) more generally for toric manifolds or some related family of manifolds. The strong cohomological rigidity does not hold for arbitrary toric manifolds while no counterexamples are known to the cohomological rigidity problems for toric manifolds. A real analogue of the rigidity problems is also studied. See survey papers [4] and [13] for details.

This paper is organized as follows. In Section 2 we review some known facts on the cohomology rings of Bott manifolds. In Section 3 we introduce new bases of the cohomology rings and restate some facts mentioned in Section 2. In Section 4 we analyze graded ring isomorphisms between the cohomology rings of two Bott manifolds, where the new bases introduced in Section 3 play a role. We prove Theorem 1.1 in Section 5 and Theorem 1.2 in Section 6.

In Section 7, we will observe that the invariance of Pontrjagin classes under cohomology ring isomorphisms hold in a slightly wider class of manifolds than that of Bott manifolds. Finally, in Section 8 we make some remarks on automorphisms of the cohomology ring of a Bott manifold, which clarifies our difficulty to solve the strong cohomological rigidity conjecture for Bott manifolds completely.

2. COHOMOLOGY RINGS OF BOTT MANIFOLDS

In this section we will recall some known facts on the cohomology rings of Bott manifolds and the quotient construction of Bott manifolds.

We denote by α_j the first Chern class of the complex line bundle L_j used to construct the Bott tower (1.1). It follows from the Borel-Hirzebruch formula ([1]) that $H^*(B_j; \mathbb{Z})$ is a free module over $H^*(B_{j-1}; \mathbb{Z})$ through the map $\pi_j^*: H^*(B_{j-1}; \mathbb{Z}) \rightarrow H^*(B_j; \mathbb{Z})$ on two generators 1 and x_j of degree 0 and 2 respectively, where x_j is the first Chern class of the tautological line bundle γ_j over B_j , and that the ring structure is determined by the single relation

$$x_j^2 = \pi_j^*(\alpha_j)x_j.$$

Using the formula inductively on j and regarding $H^*(B_j; \mathbb{Z})$ as a subring of $H^*(B_n; \mathbb{Z})$ through the projections in (1.1), we see that

$$(2.1) \quad H^*(B_n; \mathbb{Z}) = \mathbb{Z}[x_1, \dots, x_n] / (x_j^2 - \alpha_j x_j \mid j = 1, 2, \dots, n)$$

where $\alpha_1 = 0$. The following lemma easily follows from (2.1).

Lemma 2.1. *Let k be a positive integer less than or equal to n . Then the set*

$$\{x_{i_1}x_{i_2}\dots x_{i_k} \mid 1 \leq i_1 < i_2 < \dots < i_k \leq n\}$$

is an additive basis of $H^{2k}(B_n; \mathbb{Z})$.

The Pontrjagin class of a Bott manifold has a simple expression as is shown in the following lemma.

Lemma 2.2. *The Pontrjagin class $p(B_n)$ of the Bott manifold B_n is given by*

$$p(B_n) = \prod_{j=1}^n (1 + (2x_j - \alpha_j)^2) = \prod_{j=1}^n (1 + \alpha_j^2) \in H^*(B_n; \mathbb{Z})$$

where $(2x_j - \alpha_j)^2 = \alpha_j^2$ because $x_j^2 = \alpha_j x_j$.

Proof. The lemma is known but since there seems no literature which mentions the formula explicitly, we shall give a proof.

Since $\pi_j: B_j \rightarrow B_{j-1}$ is the projectivization of the Whitney sum of the trivial line bundle $\underline{\mathbb{C}}$ and the line bundle L_j over B_{j-1} , the tangent bundle TB_j of B_j splits into

$$(2.2) \quad TB_j = T_f B_j \oplus \pi_j^*(TB_{j-1})$$

where $T_f B_j$ denotes the complex line bundle along the fibers of the fiber bundle $\pi_j: B_j \rightarrow B_{j-1}$ and $\pi_j^*(TB_{j-1})$ is the pullback of the tangent bundle TB_{j-1} of the base space B_{j-1} by the projection π_j . Since x_j is the first Chern class of the tautological line bundle γ_j over B_j and the total Chern class of $\underline{\mathbb{C}} \oplus L_j$ is $1 + \alpha_j$, it follows from [1, (2) in p.515] that the total Chern class of the complex line bundle $T_f B_j$ is given by $1 - 2x_j + \alpha_j$ and hence its total Pontrjagin class is given by $1 + (2x_j - \alpha_j)^2$. This together with (2.2) shows that

$$p(B_j) = (1 + (2x_j - \alpha_j)^2)p(B_{j-1})$$

because $H^*(B_n; \mathbb{Z})$ has no 2-torsion. Then the lemma follows by applying the above formula inductively on j . \square

For $u = \sum_{i=1}^n c_i x_i \in H^2(B_n; \mathbb{Z})$ with $c_i \in \mathbb{Z}$, we define

$$\text{ht}(u) = \max\{i \in [n] = \{1, 2, \dots, n\} \mid c_i \neq 0\}$$

and call it the *height* of u . Note that $\text{ht}(\alpha_j) < \text{ht}(x_j) = j$. We say that a pair of primitive elements in $H^2(B_n; \mathbb{Z})$ is a *primitive vanishing pair* if the product of the elements vanishes. Note that a pair $(x_j, x_j - \alpha_j)$ is a primitive vanishing pair for any $j \in [n]$.

Lemma 2.3 (Lemma 2.2 in [3]). *A primitive vanishing pair is of the form*

$$(ax_j + u, a(x_j - \alpha_j) - u) \text{ or } (ax_j + u, -a(x_j - \alpha_j) + u)$$

where $a \in \mathbb{Z} \setminus \{0\}$, $u \in H^2(B_n; \mathbb{Z})$, $u(u + a\alpha_j) = 0$ and $\text{ht}(u) < j$.

Corollary 2.1 (Corollary 2.1 in [3]). *Primitive square zero elements in $H^2(B_n; \mathbb{Z})$ are of the form $\pm(x_j - \frac{1}{2}\alpha_j)$ if $\alpha_j \equiv 0 \pmod{2}$ and $\pm(2x_j - \alpha_j)$ otherwise.*

We shall review the quotient construction of Bott manifolds ([7, Prop. 3.1]). Remember that $\alpha_1 = 0$ and express

$$(2.3) \quad \alpha_j = \sum_{i=1}^{j-1} A_j^i x_i \quad \text{for } j = 2, 3, \dots, n$$

with $A_j^i \in \mathbb{Z}$. Let S^{2d-1} be the unit sphere of \mathbb{C}^d for $d = 1, 2$. Then the Bott manifold B_n in (1.1) can be obtained as the quotient of $(S^3)^n$ by the free action of

$(S^1)^n$ defined by

$$(2.4) \quad \begin{aligned} & (g_1, g_2, \dots, g_n) \cdot ((z_1, w_1), (z_2, w_2), \dots, (z_n, w_n)) \\ &= ((g_1 z_1, g_1 w_1), ((g_1^{-A_1^1}) g_2 z_2, g_2 w_2), \dots, ((\prod_{k=1}^{n-1} g_k^{-A_n^k}) g_n z_n, g_n w_n)) \end{aligned}$$

where $g_i \in S^1$ and $(z_i, w_i) \in S^3$ for $i = 1, 2, \dots, n$. Projections

$$((z_1, w_1), \dots, (z_n, w_n)) \rightarrow ((z_1, w_1), \dots, (z_{n-1}, w_{n-1})) \rightarrow \dots \rightarrow (z_1, w_1)$$

induce the Bott tower (1.1). The tautological line bundle γ_j over B_j can be described as the quotient of the trivial complex line bundle $(S^3)^n \times \mathbb{C} \rightarrow (S^3)^n$ by the action of $(S^1)^n$ on the total space $(S^3)^n \times \mathbb{C}$ defined as follows:

$$g \cdot ((z, w), u) = (g \cdot (z, w), g_j^{-1} u) \quad (u \in \mathbb{C}),$$

where $g = (g_1, g_2, \dots, g_n)$, $(z, w) = ((z_1, w_1), (z_2, w_2), \dots, (z_n, w_n))$ and $g \cdot (z, w)$ denotes the action defined in (2.4).

3. BASE CHANGE

We set

$$(3.1) \quad y_j = x_j - \frac{1}{2} \alpha_j \quad \text{for } j = 1, 2, \dots, n.$$

Then, we have

$$(3.2) \quad 4y_j^2 = (2x_j - \alpha_j)^2 = \alpha_j^2 \quad \text{in } H^*(B_n; \mathbb{Z}),$$

and it follows from Lemma 2.2 that

$$(3.3) \quad p(B_n) = \prod_{i=1}^n (1 + 4y_i^2).$$

If $\alpha_j \equiv 0 \pmod{2}$, then y_j is integral, that is, an element of $H^2(B_n; \mathbb{Z})$.

Lemma 3.1. *If $\alpha_i \equiv \alpha_j \equiv 0 \pmod{2}$ and $y_i \equiv y_j \pmod{2}$, then $i = j$.*

Proof. Since $\text{ht}(\alpha_i) < i$ and $\text{ht}(\alpha_j) < j$, the assumption $y_i \equiv y_j \pmod{2}$ implies $x_i \equiv x_j \pmod{2}$ and hence $i = j$ because $\{x_1, x_2, \dots, x_n\}$ is an additive basis of $H^2(B_n; \mathbb{Z})$ (cf. Lemma 2.1). \square

Let A_j^i ($i < j$) be the integers defined in (2.3). Setting $A_j^i = 0$ for $i \geq j$, we obtain an integral strictly upper triangular $n \times n$ matrix A with A_j^i as the (i, j) entry. Then it follows from (3.1) that

$$(y_1, \dots, y_n) = (x_1, \dots, x_n) \left(E - \frac{1}{2} A \right),$$

where E is the identity matrix. Here $E - \frac{1}{2} A$ is an upper triangular unipotent matrix and so is its inverse. Therefore, if we denote the (i, j) entry of $(E - \frac{1}{2} A)^{-1}$ by a_j^i , then

$$(3.4) \quad x_j = \sum_{i \leq j} a_j^i y_i \quad \text{and} \quad \frac{1}{2} \alpha_j = x_j - y_j = \sum_{i < j} a_j^i y_i.$$

By (3.2) and (3.4), we have

$$(3.5) \quad y_j^2 = \left(\frac{\alpha_j}{2}\right)^2 = \left(\sum_{i < j} a_j^i y_i\right)^2 \quad \text{in } H^*(B_n; \mathbb{Q}).$$

The following lemma easily follows from (3.5) or Lemma 2.1.

Lemma 3.2. *Let k be a positive integer less than or equal to n . Then the set*

$$\{y_{i_1} y_{i_2} \cdots y_{i_k} \mid 1 \leq i_1 < i_2 < \cdots < i_k \leq n\}$$

is an additive basis of $H^{2k}(B_n; \mathbb{Q})$ over \mathbb{Q} .

4. COHOMOLOGY RING ISOMORPHISMS

Let \mathcal{B}_n be the set of integral strictly upper triangular $n \times n$ matrices. Since the Bott manifold B_n in (1.1) is determined by a matrix $A \in \mathcal{B}_n$, we will denote B_n by $M(A)$. We remark that it happens that $M(A)$ and $M(B)$ are diffeomorphic even if $A, B \in \mathcal{B}_n$ are different.

Henceforth the cohomology elements x_j, y_j, α_j for $M(A)$ will be denoted by x_j^A, y_j^A, α_j^A respectively to avoid confusion.

Proposition 4.1. *Suppose that $\psi: H^*(M(A); \mathbb{Q}) \rightarrow H^*(M(B); \mathbb{Q})$ is a graded ring isomorphism. Then there are non-zero $q_1, \dots, q_n \in \mathbb{Q}$ and a permutation σ on $[n]$ such that*

$$\psi(y_j^A) = q_j y_{\sigma(j)}^B \quad \text{for } j = 1, \dots, n.$$

Proof. We prove the proposition by induction on j . When $j = 1$, $\psi(y_1^A)^2 = 0$ because $(y_1^A)^2 = 0$. Therefore $\psi(y_1^A)$ is a nonzero scalar multiple of $y_{\sigma(1)}^B$ for some $\sigma(1) \in [n]$ by Corollary 2.1.

Suppose that

$$(4.1) \quad \psi(y_j^A) = q_j y_{\sigma(j)}^B \quad \text{for } j < k.$$

Then, since

$$(q_j y_{\sigma(j)}^B)^2 = \psi(y_j^A)^2 = \psi((y_j^A)^2) = \psi\left(\left(\sum_{i < j} a_j^i y_i^A\right)^2\right) = \left(\sum_{i < j} a_j^i q_i y_{\sigma(i)}^B\right)^2,$$

one can inductively see that for $j < k$,

$$(4.2) \quad (y_{\sigma(j)}^B)^2 \text{ is a linear combination of } y_{\sigma(i_1)}^B y_{\sigma(i_2)}^B \text{ with } i_1 < i_2 < j.$$

Set $S = \{\sigma(1), \dots, \sigma(k-1)\}$. Since ψ is an isomorphism, it follows from (4.1) that $\psi(y_k^A)$ is not a linear combination of $y_{\sigma(1)}^B, \dots, y_{\sigma(k-1)}^B$. Therefore, there is $m \in [n] \setminus S$ such that

$$(4.3) \quad \psi(y_k^A) = \sum_{s \leq m \text{ or } s \in S} b_s y_s^B \quad (b_s \in \mathbb{Q} \text{ and } b_m \neq 0).$$

We consider squares at the both sides above. It follows from (3.5) and (4.1) that

$$\psi(y_k^A)^2 = \psi((y_k^A)^2) = \psi\left(\left(\sum_{i < k} a_k^i y_i^A\right)^2\right) = \left(\sum_{i < k} a_k^i q_i y_{\sigma(i)}^B\right)^2$$

and hence

$$(4.4) \quad \psi(y_k^A)^2 \text{ is a linear combination of } y_{\sigma(i_1)}^B y_{\sigma(i_2)}^B \text{ with } i_1 < i_2 < k$$

by (4.2). On the other hand, we claim that if $b_t \neq 0$ for some $t < m$ or $t \in S$ at the right hand side of (4.3), then a non-zero scalar multiple of $y_m^B y_t^B$ appears in the square of the right hand side of (4.3). Indeed, $(y_s^B)^2$ for $s \leq m$ is a linear combination of $y_i^B y_j^B$ with $i < j < m$ by (3.5) and $(y_s^B)^2$ for $s \in S$ is a linear combination of $y_i^B y_j^B$ with $i \neq j \in S$ by (4.2). Therefore the claim holds because $\{y_i^B y_j^B \mid 1 \leq i < j \leq n\}$ is an additive basis of $H^4(M(B); \mathbb{Q})$ by Lemma 3.2. However the claim contradicts (4.4) since $m \notin S$. Therefore, $\psi(y_k^A) = b_m y_m^B$. This completes the induction step and proves the proposition. \square

Lemma 4.1. *Suppose that the graded ring isomorphism ψ in Proposition 4.1 is induced from a graded ring isomorphism from $H^*(M(A); \mathbb{Z})$ to $H^*(M(B); \mathbb{Z})$. Then the rational number q_j in Proposition 4.1 belongs to $\{\pm\frac{1}{2}, \pm 1, \pm 2\}$ for $j = 1, 2, \dots, n$, and*

- (1) $q_j \in \{\pm\frac{1}{2}\}$ if and only if $\alpha_j^A \not\equiv 0 \pmod{2}$ and $\alpha_{\sigma(j)}^B \equiv 0 \pmod{2}$,
- (2) $q_j \in \{\pm 2\}$ if and only if $\alpha_j^A \equiv 0 \pmod{2}$ and $\alpha_{\sigma(j)}^B \not\equiv 0 \pmod{2}$.

Moreover, if $q_j = \pm 1$ for all j , then $\psi(p(M(A))) = p(M(B))$.

Proof. Since $\psi(x_j^A - \frac{1}{2}\alpha_j^A) = q_j(x_{\sigma(j)}^B - \frac{1}{2}\alpha_{\sigma(j)}^B)$ by Proposition 4.1, the former statement in the lemma follows from the fact that ψ restricted to the cohomology rings with integer coefficients sends primitive elements to primitive elements and the latter follows from (3.3) and Proposition 4.1. \square

We fix a graded ring isomorphism $\psi: H^*(M(A); \mathbb{Z}) \rightarrow H^*(M(B); \mathbb{Z})$ by the end of Lemma 4.3. Since $(\psi(x_j^A), \psi(x_j^A - \alpha_j^A))$ is a primitive vanishing pair, it follows from Lemma 2.3 that there are $a \in \mathbb{Z} \setminus \{0\}$, $u \in H^2(M(B); \mathbb{Z})$ and $k \in [n]$ such that

$$(4.5) \quad (\psi(x_j^A), \psi(x_j^A - \alpha_j^A)) = \begin{cases} (ax_k^B + u, a(x_k^B - \alpha_k^B) - u) & \text{or} \\ (ax_k^B + u, -a(x_k^B - \alpha_k^B) + u), \end{cases}$$

where $u(u + a\alpha_k^B) = 0$ and $\text{ht}(u) < k$. Remember that a_j^i is the (i, j) entry of the matrix $(E - \frac{1}{2}A)^{-1}$, in other words, a_j^i is the rational number determined by (3.4).

With this understood, we have

Lemma 4.2. *In the former case of (4.5), $q_j = a$ (hence q_j is an integer), $k = \sigma(j)$ and $a_j^i = 0$ for $\sigma(i) > \sigma(j)$. In the latter case of (4.5), $\alpha_j^A = 2aq_i^{-1}y_i^A$ for some $i < j$.*

Proof. Since $2y_j^A = x_j^A + (x_j^A - \alpha_j^A)$ and $\alpha_j^A = x_j^A - (x_j^A - \alpha_j^A)$, a simple computation using (4.5) shows that

$$(4.6) \quad (\psi(2y_j^A), \psi(\alpha_j^A)) = \begin{cases} (2ay_k^B, 2u + a\alpha_k^B) & \text{in the former case of (4.5),} \\ (2u + a\alpha_k^B, 2ay_k^B) & \text{in the latter case of (4.5).} \end{cases}$$

In the former case of (4.5), we have $\psi(y_j^A) = ay_k^B$ by (4.6). This together with Proposition 4.1 shows that $q_j = a$ and $k = \sigma(j)$. Moreover, since $\psi(\alpha_j^A) = 2u + a\alpha_k^B$ by (4.6) and $\text{ht}(u) < k$, we have

$$\text{ht}(\psi(\alpha_j^A)) < k = \sigma(j)$$

while since $\alpha_j^A = 2 \sum_{i < j} a_j^i y_i^A$ by (3.4), we have

$$\psi(\alpha_j^A) = 2 \sum_{i < j} a_j^i q_i y_{\sigma(i)}^B$$

by Proposition 4.1. These show that $a_j^i = 0$ for $\sigma(i) > \sigma(j)$.

In the latter case of (4.5), $\psi(\alpha_j^A) = 2ay_k^B$ by (4.6). Therefore $\alpha_j^A = 2aq_i^{-1}y_i^A$ for some $i \in [n]$ by Proposition 4.1 and $j > \text{ht}(\alpha_j^A) = \text{ht}(y_i^A) = i$. \square

Note that $2aq_i^{-1}$ in Lemma 4.2 is a nonzero integer because $a \in \mathbb{Z} \setminus \{0\}$ and $q_i \in \{\pm\frac{1}{2}, \pm 1, \pm 2\}$ by Lemma 4.1. We say that α_j is of *exceptional type* if $\alpha_j = cy_i$ for some nonzero integer c and $i < j$ and is of *even exceptional type* if the nonzero integer c is even.

Lemma 4.3. *Suppose that $q_j = \pm\frac{1}{2}$. Then there are $i < j$ and $c \in \mathbb{Z} \setminus \{0\}$ such that $\alpha_j^A = cy_i^A$. If α_j^A is not of even exceptional type, then c is odd and $q_i = \pm 2$. If neither α_j^A nor $\alpha_{\sigma(i)}^B$ are of even exceptional type, then $\alpha_{\sigma(i)}^B = dy_{\sigma(j)}^B$ with some odd integer d .*

Proof. Since $q_j = \pm\frac{1}{2}$, that is not an integer, by assumption, Lemma 4.2 says that the latter case of (4.5) must occur and $\alpha_j^A = 2aq_i^{-1}y_i^A$. Setting $c = 2aq_i^{-1}$, we have

$$(4.7) \quad \alpha_j^A = cy_i^A,$$

where c is a nonzero integer. If α_j^A is not of even exceptional type, then $c = 2aq_i^{-1}$ is odd and hence q_i must be ± 2 (and a is odd).

It remains to prove the last assertion in the lemma. Suppose that neither α_j^A nor $\alpha_{\sigma(i)}^B$ are of even exceptional type. Then $q_i = \pm 2$ as observed above, and since $\psi(y_i^A) = q_i y_{\sigma(i)}^B$ by Proposition 4.1, we have $\psi^{-1}(y_{\sigma(i)}^B) = \pm\frac{1}{2}y_i^A$. Then the same argument as above applied for $\psi^{-1}(y_{\sigma(i)}^B) = \pm\frac{1}{2}y_i^A$ tells us that there are $\ell \in [n]$ and $d \in \mathbb{Z} \setminus \{0\}$ such that

$$(4.8) \quad \alpha_{\sigma(i)}^B = dy_{\ell}^B.$$

Here d is odd because $\alpha_{\sigma(i)}^B$ is not of even exceptional type by assumption. In the sequel, it suffices to prove $\ell = \sigma(j)$.

Since d is odd, it follows from (4.8) that $\alpha_{\ell}^B \equiv 0 \pmod{2}$ while $\alpha_{\sigma(j)}^B \equiv 0 \pmod{2}$ by Lemma 4.1 since $q_j = \pm\frac{1}{2}$. Therefore, it suffices to show $y_{\ell}^B \equiv y_{\sigma(j)}^B \pmod{2}$ by Lemma 3.1, which we shall check below. Since $\pm y_{\sigma(j)}^B = \psi(2y_j^A) = \psi(2x_j^A - \alpha_j^A) = 2\psi(x_j^A) - \psi(\alpha_j^A)$, we have

$$(4.9) \quad y_{\sigma(j)}^B \equiv \psi(\alpha_j^A) \pmod{2}.$$

Similarly, since $\pm\psi(y_i^A) = 2y_{\sigma(i)}^B = 2x_{\sigma(i)}^B - \alpha_{\sigma(i)}^B$, we have

$$(4.10) \quad \psi(y_i^A) \equiv \alpha_{\sigma(i)}^B \pmod{2}.$$

Thus, since c and d are both odd integers, it follows from (4.9), (4.7), (4.10) and (4.8) that

$$y_{\sigma(j)}^B \equiv \psi(\alpha_j^A) \equiv \psi(y_i^A) \equiv \alpha_{\sigma(i)}^B \equiv y_{\ell}^B \pmod{2},$$

proving the desired congruence relation. \square

Lemma 4.4. *For $A \in \mathcal{B}_n$, there is $B \in \mathcal{B}_n$ such that none of $\alpha_1^B, \dots, \alpha_n^B$ are of even exceptional type and there is a graded ring isomorphism $\psi: H^*(M(A); \mathbb{Z}) \rightarrow H^*(M(B); \mathbb{Z})$ such that $\psi(p(M(A))) = p(M(B))$.*

Proof. Suppose that α_j^A is of even exceptional type but α_k^A for $k < j$ is not. In the following we will find $B \in \mathcal{B}_n$ such that α_k^B for $k < j$ is not of even exceptional type but $\text{ht}(\alpha_j^B) < \text{ht}(\alpha_j^A)$ and that $H^*(M(A); \mathbb{Z})$ and $H^*(M(B); \mathbb{Z})$ are isomorphic as graded rings. If α_j^B is still of even exceptional type, then we repeat the argument until we reach B such that α_k^B for $k \leq j$ is not of even exceptional type. Note that this can be achieved because if $\text{ht}(\alpha_j^B) = 0$, then $\alpha_j^B = 0$ which is not of even exceptional type. Doing this procedure inductively on j , we finally reach the desired B in the lemma.

Since α_j^A is of even exceptional type by assumption, we have

$$(4.11) \quad \alpha_j^A = c(x_i^A - \frac{1}{2}\alpha_i^A)$$

with some nonzero even integer c and $i < j$. We define a matrix B of size n by

$$B_k^\ell = \begin{cases} A_k^\ell & \text{if } k \neq j \text{ and } \ell \neq i, \\ A_k^i + \frac{c}{2}A_k^j & \text{if } k \neq j \text{ and } \ell = i, \\ -\frac{c}{2}A_i^\ell & \text{if } k = j. \end{cases}$$

Since $A \in \mathcal{B}_n$ and $i < j$, the matrix B is indeed in \mathcal{B}_n (i.e., $B_k^\ell = 0$ for $\ell \geq k$), and

$$(4.12) \quad \alpha_k^B = \begin{cases} \sum_{\ell=1}^n A_k^\ell x_\ell^B + \frac{c}{2}A_k^j x_i^B & \text{if } k \neq j, \\ -\frac{c}{2} \sum_{\ell=1}^n A_i^\ell x_\ell^B & \text{if } k = j. \end{cases}$$

Note that since $i < j$, $A_i^j = 0$ and hence it follows from (4.12) that

$$(4.13) \quad \alpha_j^B = -\frac{c}{2}\alpha_i^B.$$

We define

$$(4.14) \quad \psi(x_\ell^A) = \begin{cases} x_\ell^B & \text{if } \ell \neq j, \\ x_j^B + \frac{c}{2}x_i^B & \text{if } \ell = j. \end{cases}$$

This clearly induces an isomorphism $\psi: \mathbb{Z}[x_1^A, \dots, x_n^A] \rightarrow \mathbb{Z}[x_1^B, \dots, x_n^B]$ between polynomial rings. We claim that ψ induces a graded isomorphism from $H^*(M(A); \mathbb{Z})$ to $H^*(M(B); \mathbb{Z})$. Indeed, when $k \neq j$, it follows from (4.14) and (4.12) that

$$(4.15) \quad \psi(\alpha_k^A) = \psi\left(\sum_{\ell=1}^n A_k^\ell x_\ell^A\right) = \sum_{\ell=1}^n A_k^\ell x_\ell^B + \frac{c}{2}A_k^j x_i^B = \alpha_k^B$$

and hence

$$(4.16) \quad \psi(x_k^A(x_k^A - \alpha_k^A)) = x_k^B(x_k^B - \alpha_k^B) \quad \text{when } k \neq j.$$

When $k = j$, we have

$$\begin{aligned}
(4.17) \quad \psi(x_j^A(x_j^A - \alpha_j^A)) &= \psi(x_j^A(x_j^A - c(x_i^A - \frac{1}{2}\alpha_i^A))) && \text{by (4.11)} \\
&= (x_j^B + \frac{c}{2}x_i^B)(x_j^B - \frac{c}{2}x_i^B + \frac{c}{2}\alpha_i^B) && \text{by (4.14) and (4.15)} \\
&= x_j^B(x_j^B + \frac{c}{2}\alpha_i^B) - \frac{c^2}{4}x_i^B(x_i^B - \alpha_i^B) \\
&= x_j^B(x_j^B - \alpha_j^B) - \frac{c^2}{4}x_i^B(x_i^B - \alpha_i^B) && \text{by (4.13)}.
\end{aligned}$$

(4.16) and (4.17) together with (2.1) show that ψ induces a graded ring isomorphism from $H^*(M(A); \mathbb{Z})$ to $H^*(M(B); \mathbb{Z})$.

Since $\text{ht}(\alpha_j^A) = i$ by (4.11) and $\text{ht}(\alpha_j^B) = \text{ht}(\alpha_i^B) < i$ by (4.13), we have $\text{ht}(\alpha_j^B) < \text{ht}(\alpha_j^A)$. Moreover, the isomorphism ψ defined in (4.14) is represented as a unipotent upper triangular matrix with respect to the basis x_1^A, \dots, x_n^A of $H^2(M(A); \mathbb{Z})$ and x_1^B, \dots, x_n^B of $H^2(M(B); \mathbb{Z})$, so $q_j = 1$ for any j and hence $\psi(p(M(A))) = p(M(B))$ by Lemma 4.1. \square

Remark 4.1. The graded ring isomorphism ψ in the proof above is actually induced from a diffeomorphism, which follows from Theorem 6.1 mentioned in Section 6. One can also see it using the quotient construction of Bott manifolds.

5. PROOF OF THEOREM 1.1

This section is devoted to the proof of Theorem 1.1 in the Introduction.

Let $\psi: H^*(M(A); \mathbb{Z}) \rightarrow H^*(M(B); \mathbb{Z})$ be a graded ring isomorphism. By (3.3) what we must prove is

$$\psi\left(\prod_{j=1}^n (1 + 4(y_j^A)^2)\right) = \prod_{j=1}^n (1 + 4(y_j^B)^2) \in H^*(M(B); \mathbb{Z}).$$

We may assume that none of $\alpha_1^A, \dots, \alpha_n^A$ and $\alpha_1^B, \dots, \alpha_n^B$ are of even exceptional type by Lemma 4.4. Since $\psi(y_j^A) = q_j y_{\sigma(j)}^B$ by Proposition 4.1, $\psi(1 + 4(y_j^A)^2) = 1 + 4(y_{\sigma(j)}^B)^2$ if $q_j = \pm 1$. Therefore, we shall treat the case where $q_j = \pm \frac{1}{2}$ or ± 2 by Lemma 4.1.

Suppose $q_j = \pm \frac{1}{2}$. Then there is $i < j$ such that

$$(5.1) \quad \alpha_j^A = cy_i^A \quad \text{and} \quad \alpha_{\sigma(i)}^B = dy_{\sigma(j)}^B$$

with some odd integers c, d and $q_i = \pm 2$ by Lemma 4.3. We shall show that

$$(5.2) \quad \psi((1 + 4(y_j^A)^2)(1 + 4(y_i^A)^2)) = (1 + 4(y_{\sigma(j)}^B)^2)(1 + 4(y_{\sigma(i)}^B)^2).$$

It follows from (3.2) and (5.1) that

$$(5.3) \quad 4(y_j^A)^2 = (\alpha_j^A)^2 = c^2(y_i^A)^2$$

and sending the first and last elements in the identity above by ψ , we obtain

$$(5.4) \quad (y_{\sigma(j)}^B)^2 = 4c^2(y_{\sigma(i)}^B)^2$$

since $q_j = \pm \frac{1}{2}$ and $q_i = \pm 2$. Here $4(y_{\sigma(i)}^B)^2 = (\alpha_{\sigma(i)}^B)^2$ by (3.2) and $(\alpha_{\sigma(i)}^B)^2 = d^2(y_{\sigma(j)}^B)^2$ by (5.1). Therefore, (5.4) turns into

$$(5.5) \quad (y_{\sigma(j)}^B)^2 = c^2 d^2 (y_{\sigma(j)}^B)^2.$$

When $(y_{\sigma(i)}^B)^2 \neq 0$, (5.5) implies $c^2 d^2 = 1$ and hence $c^2 = d^2 = 1$ because c, d are integers. It follows from (5.3) and (5.4) that

$$4(y_j^A)^2 = (y_i^A)^2 \quad \text{and} \quad (y_{\sigma(j)}^B)^2 = 4(y_{\sigma(i)}^B)^2.$$

Therefore

$$\begin{aligned} \text{the left hand side of (5.2)} &= \psi((1 + (y_i^A)^2)(1 + 16(y_j^A)^2)) \\ &= (1 + q_i^2 (y_{\sigma(i)}^B)^2)(1 + 16q_j^2 (y_{\sigma(j)}^B)^2) \\ &= \text{the right hand side of (5.2)} \end{aligned}$$

because $q_j = \pm \frac{1}{2}$ and $q_i = \pm 2$. When $(y_{\sigma(i)}^B)^2 = 0$, we have $(y_{\sigma(j)}^B)^2 = 0$ by (5.4) and $(y_j^A)^2 = (y_i^A)^2 = 0$ because $\psi(y_j^A) = y_{\sigma(j)}^B$ and $\psi(y_i^A) = y_{\sigma(i)}^B$ and ψ is an isomorphism. Therefore, (5.2) holds even when $(y_{\sigma(i)}^B)^2 = 0$.

When $q_j = \pm 2$, $\psi^{-1}(y_{\sigma(j)}^B) = \pm \frac{1}{2} y_j^A$. Therefore, the same argument as above applied for ψ^{-1} shows that (5.2) also holds when $q_j = \pm 2$. This completes the proof of the theorem.

6. PROOF OF THEOREM 1.2

The purpose of this section is to prove Theorem 1.2 in the Introduction. We begin with the following lemma.

Lemma 6.1. *Let $A \in \mathcal{B}_n$, σ a permutation on $[n]$ and let P be the permutation matrix of σ^{-1} , that is, the (i, j) entry of P is 1 if $i = \sigma(j)$ and 0 otherwise. If $PAP^{-1} \in \mathcal{B}_n$, then there is a graded ring isomorphism $\psi_\sigma: H^*(M(A); \mathbb{Z}) \rightarrow H^*(M(PAP^{-1}); \mathbb{Z})$ sending x_j^A to $x_{\sigma(j)}^{PAP^{-1}}$ for $j = 1, 2, \dots, n$ and it is induced from a diffeomorphism.*

Proof. Remember the quotient construction of Bott manifolds explained in Section 2. Let $\varphi_\sigma: (S^3)^n \rightarrow (S^3)^n$ be the coordinate change defined by

$$\begin{aligned} &\varphi_\sigma((z_1, w_1), (z_2, w_2), \dots, (z_n, w_n)) \\ &= ((z_{\sigma(1)}, w_{\sigma(1)}), (z_{\sigma(2)}, w_{\sigma(2)}), \dots, (z_{\sigma(n)}, w_{\sigma(n)})) \end{aligned}$$

where we consider the $(S^1)^n$ -action associated to the matrix PAP^{-1} on the source space and the one associated to A on the target space, and let ϕ_σ be the group automorphism of $(S^1)^n$ defined by

$$\phi_\sigma(g_1, g_2, \dots, g_n) = (g_{\sigma(1)}, g_{\sigma(2)}, \dots, g_{\sigma(n)}).$$

Then, φ_σ is ϕ_σ -equivariant, i.e.,

$$(6.1) \quad \varphi_\sigma(g \cdot (z, w)) = \phi_\sigma(g) \cdot \varphi_\sigma((z, w)),$$

where $g = (g_1, \dots, g_n)$ and $(z, w) = ((z_1, w_1), (z_2, w_2), \dots, (z_n, w_n))$. Indeed, the j -th component of the left hand side of (6.1) is

$$(6.2) \quad \left(\left(\prod_{k=1}^{\sigma(j)-1} g_k \right)^{-(PAP^{-1})_{\sigma(j)}^k} \right) g_{\sigma(j)} z_{\sigma(j)}, g_{\sigma(j)} w_{\sigma(j)}$$

while that of the right hand side of (6.1) is

$$(6.3) \quad \left(\left(\prod_{i=1}^{j-1} g_{\sigma(i)} \right)^{-A_{\sigma(i)}^j} \right) g_{\sigma(j)} z_{\sigma(j)}, g_{\sigma(j)} w_{\sigma(j)}.$$

Here, since P is the permutation matrix of σ^{-1} , $(PAP^{-1})_{\sigma(j)}^{\sigma(i)} = A_j^i$ and it is zero for $\sigma(i) \geq \sigma(j)$ or $i \geq j$ since both PAP^{-1} and A belong to \mathcal{B}_n ; so we have

$$\prod_{k=1}^{\sigma(j)-1} g_k^{-(PAP^{-1})_{\sigma(j)}^k} = \prod_{k=1}^n g_k^{-(PAP^{-1})_{\sigma(j)}^k} = \prod_{i=1}^n g_{\sigma(i)}^{-(PAP^{-1})_{\sigma(j)}^{\sigma(i)}} = \prod_{i=1}^n g_{\sigma(i)}^{-A_j^i} = \prod_{i=1}^{j-1} g_{\sigma(i)}^{-A_j^i}.$$

This shows that (6.2) and (6.3) agree, which means that φ_σ is ϕ_σ -equivariant and hence φ_σ induces a diffeomorphism from $M(PAP^{-1})$ to $M(A)$.

It remains to prove that the cohomology ring isomorphism induced by φ_σ maps x_j^A to $x_{\sigma(j)}^{PAP^{-1}}$. Remember that x_j^A and $x_{\sigma(j)}^{PAP^{-1}}$ are the first Chern classes of the complex line bundles γ_j^A and $\gamma_{\sigma(j)}^{PAP^{-1}}$ mentioned in Section 2, where the matrices A and PAP^{-1} are specified to avoid confusion. Therefore, it suffices to find a bundle isomorphism from $\gamma_{\sigma(j)}^{PAP^{-1}}$ to γ_j^A which covers φ_σ . The map f_σ from $(S^3)^n \times \mathbb{C}$ to itself defined by

$$f_\sigma((z, w), u) = (\varphi_\sigma(z, w), u)$$

satisfies

$$f_\sigma(g \cdot (z, w), g_{\sigma(j)}^{-1} u) = (\phi_\sigma(g) \cdot (z, w), g_{\sigma(j)}^{-1} u).$$

Since $g_{\sigma(j)}$ is the j -th component of $\phi_\sigma(g)$, f_σ induces the desired bundle isomorphism from $\gamma_{\sigma(j)}^{PAP^{-1}}$ to γ_j^A . \square

The following theorem due to H. Ishida plays a role in our argument.

Theorem 6.1 ([9]). *Let $A, B \in \mathcal{B}_n$ and $\psi: H^*(M(A); \mathbb{Z}) \rightarrow H^*(M(B); \mathbb{Z})$ be a graded ring isomorphism. If ψ restricted to the degree two cohomology groups is represented as an upper triangular matrix with respect to the basis x_1^A, \dots, x_n^A of $H^2(M(A); \mathbb{Z})$ and x_1^B, \dots, x_n^B of $H^2(M(B); \mathbb{Z})$ defined in Section 2, then ψ is induced from a diffeomorphism.*

Let $\psi: H^*(M(A); \mathbb{Z}) \rightarrow H^*(M(B); \mathbb{Z})$ be a graded ring isomorphism. By Proposition 4.1, there is a permutation σ on $[n]$ such that $\psi(y_j^A) = q_j y_{\sigma(j)}^B$ with some nonzero $q_j \in \mathbb{Q}$ for any j .

Proposition 6.1. *If $q_j \in \{\pm 1\}$ for all j , then the isomorphism ψ above is induced from a diffeomorphism.*

Proof. We may assume that none of $\alpha_1^A, \dots, \alpha_n^A$ are of even exceptional type by Lemma 4.4 and Remark 4.1 (replace A with the matrix given in Lemma 4.4 if necessary). Let a_j^i be the (i, j) entry of $(E - \frac{1}{2}A)^{-1}$. Because none of $\alpha_1^A, \dots, \alpha_n^A$ are of even exceptional type and $q_k \in \{\pm 1\}$ for all k , we have $a_j^i = 0$ for $\sigma(i) > \sigma(j)$ by Lemma 4.2. This means that if P is the permutation matrix of σ^{-1} , then $P(E - \frac{1}{2}A)^{-1}P^{-1}$ is a unipotent upper triangular matrix because the $(\sigma(i), \sigma(j))$ entry of $P(E - \frac{1}{2}A)^{-1}P^{-1}$ is equal to the (i, j) entry a_j^i of $(E - \frac{1}{2}A)^{-1}$. Then, since $P(E - \frac{1}{2}A)P^{-1} = (P(E - \frac{1}{2}A)^{-1}P^{-1})^{-1}$ is a unipotent upper triangular matrix, PAP^{-1} is a strictly upper triangular matrix and hence $PAP^{-1} \in \mathcal{B}_n$.

Let $\psi_\sigma: H^*(M(A); \mathbb{Z}) \rightarrow H^*(M(PAP^{-1}); \mathbb{Z})$ be the graded ring isomorphism in Lemma 6.1. Then the graded ring isomorphism arising from the composition

$$\psi \circ \psi_\sigma^{-1}: H^*(M(PAP^{-1}); \mathbb{Z}) \rightarrow H^*(M(A); \mathbb{Z}) \rightarrow H^*(M(B); \mathbb{Z})$$

satisfies $(\psi \circ \psi_\sigma^{-1})(y_j^{PAP^{-1}}) = q_{\sigma^{-1}(j)} y_j^B$ for any j since $\psi_\sigma(x_i^A) = x_{\sigma(i)}^{PAP^{-1}}$ for any i . This means that $\psi \circ \psi_\sigma^{-1}$ satisfies the assumption in Theorem 6.1 and hence is induced from a diffeomorphism. Moreover, ψ_σ is induced from a diffeomorphism by Lemma 6.1. This shows that ψ is induced from a diffeomorphism. \square

Remember that $M(O) = (\mathbb{C}P^1)^n$ for the zero matrix $O \in \mathcal{B}_n$. We say that a Bott manifold $M(A)$ for $A \in \mathcal{B}_n$ is $\mathbb{Z}/2$ -trivial if $H^*(M(A); \mathbb{Z}/2) \cong H^*(M(O); \mathbb{Z}/2)$ as graded rings.

Lemma 6.2. *Let $A \in \mathcal{B}_n$. Then the following three statements are equivalent.*

- (1) $M(A)$ is $\mathbb{Z}/2$ -trivial.
- (2) $\alpha_j^A \equiv 0 \pmod{2}$ for $j = 1, 2, \dots, n$.
- (3) $A \equiv O \pmod{2}$, i.e., every entry of A is an even integer.

Proof. The equivalence (2) \Leftrightarrow (3) and the implication (2) \Rightarrow (1) are obvious, so it suffices to prove the implication (1) \Rightarrow (2). Suppose that $M(A)$ is $\mathbb{Z}/2$ -trivial. Then the square of any element in $H^2(M(A); \mathbb{Z}/2)$ vanishes. Therefore $\alpha_j^A x_j^A (= x_j^{A^2})$ vanishes in $H^4(M(A); \mathbb{Z}/2)$ for any j . On the other hand, the set $\{x_i^A x_j^A \mid 1 \leq i < j \leq n\}$ is an additive basis of $H^4(M(A); \mathbb{Z}/2)$ because it is an additive basis of $H^4(M(A); \mathbb{Z})$ by Lemma 2.1. Since α_j^A is a linear combination of x_i^A for $1 \leq i < j$, this implies that $\alpha_j^A = 0$ in $H^2(M(A); \mathbb{Z}/2)$ for any j , proving (2). \square

Now we are in a position to prove our second main result stated in the Introduction.

Proof of Theorem 1.2. Let $M(A)$ and $M(B)$ be $\mathbb{Z}/2$ -trivial Bott manifolds and let $\psi: H^*(M(A); \mathbb{Z}) \rightarrow H^*(M(B); \mathbb{Z})$ be any graded ring isomorphism. Since $M(A)$ and $M(B)$ are $\mathbb{Z}/2$ -trivial, $\alpha_j^A \equiv 0 \pmod{2}$ and $\alpha_j^B \equiv 0 \pmod{2}$ for any j by Lemma 6.2. This together with Lemma 4.1 shows that the assumption of Proposition 6.1 is satisfied for the ψ , so the theorem follows from Proposition 6.1. \square

7. COHOMOLOGY BOTT MANIFOLDS

In this section we observe that the invariance of Pontrjagin classes under cohomology ring isomorphisms holds for a slightly wider class of manifolds than the class of Bott manifolds.

As mentioned in the Introduction, Bott manifolds are examples of toric manifolds, where a toric manifold of complex dimension n is a smooth compact algebraic variety of complex dimension n with an algebraic action of $(\mathbb{C}^*)^n$ having an open dense orbit. Several topological analogs of toric manifolds are known and the widest class among them is the class of torus manifolds. A *torus manifold* is a $2n$ -dimensional closed smooth manifold M with an effective smooth action of an n -dimensional torus $T = (S^1)^n$ having a fixed point. A toric manifold with the restricted action of T is a torus manifold. We remark that a torus manifold is not necessarily simply connected while any toric manifold is simply connected.

We review some general facts on torus manifolds from [10] and [11]. Suppose that the cohomology ring of a torus manifold M of real dimension $2n$ is generated by degree two elements. Then the orbit space $Q := M/T$ is a nice manifold with corners, any face is acyclic and any multiple intersection of faces in Q is connected unless the intersection is empty, so Q looks like a simple polytope. Let $p: M \rightarrow Q$ be the quotient map and Q_i ($i = 1, 2, \dots, m$) be the facets of Q . Then $M_i := p^{-1}(Q_i)$ is

a closed smooth codimension-two submanifold of M and fixed pointwise under some circle subgroup of T . We choose orientations on M and M_i 's and let $x_i \in H^2(M; \mathbb{Z})$ be the Poincaré dual to the cycle M_i for $i = 1, 2, \dots, m$. Then

$$(7.1) \quad p(M) = \prod_{i=1}^m (1 + x_i^2).$$

A primitive element $v_i \in H_2(BT; \mathbb{Z})$ is uniquely associated to each M_i . In fact, through a natural identification of $H_2(BT; \mathbb{Z})$ and the group of homomorphisms from S^1 to T , $v_i(S^1)$ is the circle subgroup which fixes M_i pointwise. There are two such primitive elements and the orientations on M and M_i determine v_i uniquely. These v_i 's appear in the presentation of the cohomology ring of M as follows:

$$(7.2) \quad H^*(M; \mathbb{Z}) = \mathbb{Z}[x_1, \dots, x_m] / \mathcal{K}$$

where \mathcal{K} is the ideal generated by the following two types of elements:

- (1) $\prod_{i \in I} x_i$ for $I \subset \{1, 2, \dots, m\}$ such that $\bigcap_{i \in I} Q_i = \emptyset$,
- (2) $\sum_{i=1}^m \langle u, v_i \rangle x_i$ for $u \in H^2(BT; \mathbb{Z})$,

where $\langle \cdot, \cdot \rangle$ denotes the natural pairing between $H^2(BT; \mathbb{Z})$ and $H_2(BT; \mathbb{Z})$. The vectors v_i 's satisfy the following non-singular condition (*):

for any subset I of $\{1, 2, \dots, m\}$ with cardinality n such that $\bigcap_{i \in I} Q_i \neq \emptyset$ (in fact, $\bigcap_{i \in I} Q_i$ is a vertex), the set $\{v_i \mid i \in I\}$ forms a basis of $H_2(BT; \mathbb{Z})$.

We say that a closed smooth manifold is a *cohomology Bott manifold* if its integral cohomology ring is isomorphic to that of some Bott manifold B_n as graded rings. Assume that a torus manifold M is a cohomology Bott manifold. The following proposition shows that the Pontrjagin class of M is also of the same form as that of B_n , so the invariance of Pontrjagin classes under cohomology ring isomorphisms holds for the class of cohomology Bott manifolds.

Proposition 7.1. *Suppose that a torus manifold M of real dimension $2n$ is a cohomology Bott manifold. Then there are elements $x_1, x_2, \dots, x_n \in H^2(M; \mathbb{Z})$ such that*

$$(7.3) \quad H^*(M; \mathbb{Z}) = \mathbb{Z}[x_1, x_2, \dots, x_n] / (x_j^2 - \alpha_j x_j \mid j = 1, 2, \dots, n)$$

where $\alpha_1 = 0$ and α_j for $j = 2, \dots, n$ is a linear combination of x_1, \dots, x_{j-1} over \mathbb{Z} and that $p(M) = \prod_{i=1}^n (1 + \alpha_i^2)$.

Proof. We will freely use the notation above. If M is a genuine Bott manifold, then it is known and easy to see that the orbit space Q is an n -cube. As for a cohomology Bott manifold M , the orbit space Q is not necessarily an n -cube but the argument in [5, Theorem 5.3] or [12, Theorem 5.5] shows that the face poset of Q is isomorphic to that of an n -cube. Therefore, Q has $2n$ facets (i.e. $m = 2n$) and one can order the facets Q_i of Q in such a way that $Q_j \cap Q_{j+n} = \emptyset$ for $j = 1, 2, \dots, n$. Then (1) after (7.2) is nothing but $x_j x_{n+j}$ for $j = 1, 2, \dots, n$. Since $\bigcap_{j=1}^n Q_{n+j}$ is a vertex, the set $\{v_{n+1}, v_{n+2}, \dots, v_{2n}\}$ forms a basis of $H_2(BT; \mathbb{Z})$ by the non-singular condition (*) above. Let $\{u_1, \dots, u_n\}$ be the basis of $H^2(BT; \mathbb{Z})$ such that $\langle u_i, v_{n+j} \rangle = \delta_{ij}$ where δ_{ij} denotes Kronecker delta. Taking $u = u_j$ in (2) after (7.2),

one has $x_{n+j} = -\sum_{i=1}^n \langle u_j, v_i \rangle x_i$ in $H^*(M; \mathbb{Z})$; so (7.2) reduces to

$$(7.4) \quad H^*(M; \mathbb{Z}) = \mathbb{Z}[x_1, x_2, \dots, x_n] / (x_j (\sum_{i=1}^n \langle u_j, v_i \rangle x_i) \mid j = 1, 2, \dots, n).$$

The non-singular condition (*) above implies that every principal minor of an $n \times n$ matrix Λ with $\langle u_j, v_i \rangle$ as (i, j) entry is ± 1 . We may assume that all the diagonal entries of the matrix Λ are -1 if necessary by changing the orientations on M_i for $i = 1, \dots, n$. Then it is shown in [6] that the matrix Λ is conjugate to an upper triangular matrix by a permutation matrix, where we again use the assumption that M is a cohomology Bott manifold. Since the conjugation by a permutation matrix is nothing but a re-ordering of facets, we may assume that Λ is upper-triangular and its diagonal entries are all -1 . This shows that $x_{n+j} = x_j - \alpha_j$ where $\alpha_1 = 0$ and α_j for $j = 2, \dots, n$ is a linear combination of x_1, \dots, x_{j-1} over \mathbb{Z} and (7.4) reduces to (7.3).

Finally, since $x_{n+j} = x_j - \alpha_j$ and $x_j(x_j - \alpha_j) = 0$ in $H^*(M; \mathbb{Z})$ for $j = 1, 2, \dots, n$, it follows from (7.1) that

$$\begin{aligned} p(M) &= \prod_{j=1}^n (1 + x_j^2)(1 + x_{n+j}^2) = \prod_{j=1}^n (1 + x_j^2)(1 + (x_j - \alpha_j)^2) \\ &= \prod_{j=1}^n (1 + x_j^2 + (x_j - \alpha_j)^2) = \prod_{j=1}^n (1 + 2x_j(x_j - \alpha_j) + \alpha_j^2) = \prod_{j=1}^n (1 + \alpha_j^2). \end{aligned}$$

This completes the proof of the proposition. \square

Corollary 7.1. *Let M be a torus manifold and B_n a Bott manifold. Then, any graded ring isomorphism $\varphi: H^*(M; \mathbb{Z}) \rightarrow H^*(B_n; \mathbb{Z})$ preserves their Pontrjagin classes.*

8. CONCLUDING REMARKS

We conclude this paper with some remarks on automorphisms of the cohomology ring of a Bott manifold B_n . Proposition 4.1 and Lemma 4.1 say that given a graded ring automorphism ψ of $H^*(B_n; \mathbb{Z})$, there are a permutation σ on $[n]$ and $q_j \in \{\pm \frac{1}{2}, \pm 1, \pm 2\}$ such that $\psi(y_j) = q_j y_{\sigma(j)}$ for each $j = 1, 2, \dots, n$. Therefore, assigning $(q_j/|q_j|)_{j=1}^n$ together with σ to ψ , we obtain a monomorphism

$$(8.1) \quad \text{Aut}(H^*(B_n; \mathbb{Z})) \hookrightarrow \{\pm 1\}^n \rtimes \mathfrak{S}_n.$$

Here $\text{Aut}(H^*(B_n; \mathbb{Z}))$ denotes the group of graded ring automorphisms of $H^*(B_n; \mathbb{Z})$ and $\{\pm 1\}^n \rtimes \mathfrak{S}_n$ is the signed permutation group on $[n]$, that is, the semidirect product of the n -fold product $\{\pm 1\}^n$ of the order two group $\{\pm 1\}$ and the permutation group \mathfrak{S}_n on $[n]$ where the action of \mathfrak{S}_n on $\{\pm 1\}^n$ is the natural permutation of factors of $\{\pm 1\}^n$. Lemma 4.2 in [9] implies that any automorphism in the subgroup $\{\pm 1\}^n$ of $\{\pm 1\}^n \rtimes \mathfrak{S}_n$ can be realized by diffeomorphisms of B_n and our Lemma 6.1 implies that some elements of \mathfrak{S}_n can also be realized by diffeomorphisms of B_n . However, any q_j is equal to ± 1 for the cohomology automorphisms induced from those diffeomorphisms, so if some q_j is not ± 1 then our results do not prove that the cohomology automorphism is induced from a diffeomorphism. This is the reason why we needed to assume the $\mathbb{Z}/2$ -triviality in Theorem 1.2.

As remarked in the Introduction, the Hirzebruch surface $\mathbb{C}P^2 \sharp \overline{\mathbb{C}P^2}$ is not $\mathbb{Z}/2$ -trivial (although it is \mathbb{Q} -trivial). If we put $B_2 = \mathbb{C}P^2 \sharp \overline{\mathbb{C}P^2}$, then

$$H^*(B_2; \mathbb{Z}) = \mathbb{Z}[x_1, x_2]/(x_1^2, x_2^2 - x_1x_2),$$

so $y_1 = x_1$ and $y_2 = x_2 - \frac{1}{2}x_1$. The map (8.1) above is an isomorphism in this case. In fact, $\text{Aut}(H^*(B_2; \mathbb{Z}))$ is generated as a group by the following three automorphisms:

- (1) $(x_1, x_2) \rightarrow (-x_1, -x_1 + x_2)$, so $(y_1, y_2) \rightarrow (-y_1, y_2)$,
- (2) $(x_1, x_2) \rightarrow (x_1, x_1 - x_2)$, so $(y_1, y_2) \rightarrow (y_1, -y_2)$,
- (3) $(x_1, x_2) \rightarrow (-x_1 + 2x_2, x_2)$, so $(y_1, y_2) \rightarrow (2y_2, \frac{1}{2}y_1)$.

The automorphisms (1) and (2) generate the subgroup $\{\pm 1\}^2$ while the automorphism (3) generates the subgroup \mathfrak{S}_2 . As remarked above, the automorphisms (1) and (2) are induced from diffeomorphisms of B_2 . One can see that the automorphism (3) is also induced from a diffeomorphism of B_2 , in fact, the diffeomorphism of Type 2 in the proof of [3, Lemma 5.2] induces the automorphism (3) up to sign. For the convenience of the reader, we shall review the construction of the diffeomorphism. We take involutions

$$s: [z_1, z_2, z_3] \rightarrow [\bar{z}_1, \bar{z}_2, \bar{z}_3] \text{ on } \mathbb{C}P^2, \quad t: [z_1, z_2, z_3] \rightarrow [z_1, z_2, -z_3] \text{ on } \overline{\mathbb{C}P^2}$$

where $[z_1, z_2, z_3]$ denotes the homogeneous coordinate of $\mathbb{C}P^2$ and \bar{z} denotes the conjugate of a complex number z . The fixed point set by s is $\mathbb{R}P^2$ while that by t consists of a point and $\mathbb{C}P^1$. Choose a point from $\mathbb{R}P^2$ and $\mathbb{C}P^1$ respectively and take the equivariant connected sum of $\mathbb{C}P^2$ and $\overline{\mathbb{C}P^2}$ around the chosen fixed points. Then the resulting involution on $B_2 = \mathbb{C}P^2 \sharp \overline{\mathbb{C}P^2}$ is the desired one, see the proof of [3, Lemma 5.2] for more details. As you see, this construction heavily depends on the description $\mathbb{C}P^2 \sharp \overline{\mathbb{C}P^2}$ and we do not know how to find such a diffeomorphism for a not $\mathbb{Z}/2$ -trivial Bott manifold B_n although such a diffeomorphism is found when B_n is \mathbb{Q} -trivial ([3]). If we overcome this difficulty, then we could solve the strong cohomological rigidity conjecture for Bott manifolds completely.

REFERENCES

- [1] A. Borel and F. Hirzebruch, *Characteristic classes and homogeneous spaces I*, Amer. J. Math., **80** (1958), 458–538.
- [2] S. Choi, *Classification of Bott manifolds up to dimension eight*, to appear in Proc. Edinburgh Math. Soc.; arXiv:1112.2321.
- [3] S. Choi and M. Masuda, *Classification of \mathbb{Q} -trivial Bott manifolds*, J. Symp. Geom. **10** (2012), 447–462.
- [4] S. Choi, M. Masuda, and D. Y. Suh, *Rigidity problems in toric topology, a survey*, Proc. Steklov Inst. Math. **275** (2011), 177–190.
- [5] S. Choi, T. E. Panov and D. Y. Suh, *Toric cohomological rigidity of simple polytopes*, J. London Math. Soc. **82**(2) (2010), 343–360.
- [6] S. Choi and D. Y. Suh, *Properties of Bott manifolds and cohomological rigidity*, Algebr. Geom. Topol. **11**(2), (2011), 1053–1076.
- [7] Y. Civan and N. Ray, *Homotopy decompositions and K-theory of Bott towers*. K-Theory **34** (2005), 1–33.
- [8] D. Crowley and M. Kreck, *Hirzebruch surfaces*, Bulletin of the Manifold Atlas, 2011.
- [9] H. Ishida, *(Filtered) cohomological rigidity of Bott manifolds*, Osaka J. Math. **49** (2012), 515–522.
- [10] M. Masuda, *Unitary toric manifolds, multi-fans and equivariant index*, Tohoku Math. J. (2) **51** (1999), no. 2, 237–265.
- [11] M. Masuda and T. E. Panov, *On the cohomology of torus manifolds*, Osaka J. Math. **43** (2006), no. 3, 711–746.

- [12] M. Masuda and T. E. Panov, *Semi-free circle actions, Bott towers, and quasitoric manifolds*, Mat. Sb. **199** (8) (2008) 95–122.
- [13] M. Masuda and D. Y. Suh, *Classification problems of toric manifolds via topology*, in: Toric topology, Contemp. Math., Amer. Math. Soc. **460**, Providence, RI, 2008, 273–286.
- [14] S. P. Novikov, *Homotopically equivalent smooth manifolds. I.*, Izv. Akad. Nauk SSSR Ser. Mat. **28** (1964), 365–474.
- [15] T. Petrie, *Smooth S^1 actions on homotopy complex projective spaces and related topics*, Bull. Amer. Math. Soc. **78** (1972), 105–153.
- [16] T. Petrie, *Torus actions on homotopy complex projective spaces*, Invent. Math. **20** (1973), 139–146.
- [17] Manifold Atlas Project, http://www.map.mpim-bonn.mpg.de/Petrie_conjecture

DEPARTMENT OF MATHEMATICS, AJOU UNIVERSITY, REPUBLIC OF KOREA

DEPARTMENT OF MATHEMATICS, OSAKA CITY UNIVERSITY, SUMIYOSHI-KU, OSAKA 558-8585, JAPAN

DEPARTMENT OF MATHEMATICAL SCIENCE, FACULTY OF SCIENCE, YAMAGUCHI UNIVERSITY, 1677-1 YOSHIDA, YAMAGUCHI 753-8512, JAPAN